



SHORT COMMUNICATION

New distributional records of the Amazon River Frog *Lithobates palmipes* (Spix, 1824) in Peru

¹Roy Santa-Cruz, ²J. Amanda Delgado C., ³Cinthya Y. Salas, and ⁴Rudolf von May

^{1,3}Museo de Historia Natural, Universidad Nacional de San Agustín de Arequipa, PERÚ ²Museo de Historia Natural, Universidad Nacional de San Antonio Abad del Cusco, Cusco, PERÚ ⁴Department of Ecology and Evolutionary Biology, University of Michigan, 2051 Ruthven Museums Building, 1109 Geddes Avenue., Ann Arbor, Michigan 48109, USA

Key words. Amazonia, lowland rainforest, *Rana*, Ranidae, true frogs

Citation: Santa-Cruz R, Delgado C JA, Salas CY, von May R. 2016. New distributional records of the Amazon River Frog *Lithobates palmipes* (Spix, 1824) in Peru. *Amphibian & Reptile Conservation* 10(1) [Special Section]: 17–20 (e118).

Copyright: © 2016 Santa-Cruz et al. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits unrestricted use for non-commercial and education purposes only, in any medium, provided the original author and the official and authorized publication sources are recognized and properly credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website <amphibian-reptile-conservation.org>.

Received: 25 November 2015; **Accepted:** 12 April 2016; **Published:** 30 May 2016

The Amazon River Frog *Lithobates palmipes* (Spix, 1824) is an aquatic-breeding species that inhabits various types of rainforest throughout the lowlands of northern South America, including both the Amazon and Orinoco basins, part of the Guyana Shield, the Atlantic Forest, the Cerrado and neighboring areas in Brazil (Hillis and De Sá 1988; Acosta-Galvis 1999; Canedo and Bilate 2005; Oliveira et al. 2010; Ferreira and Faria 2011; Ramalho et al. 2011; Santos and Vaz-Silva 2012; Rodrigues et al. 2013; Frost 2015). According to Hillis and De Sá (1988), this species belongs to the complex *Rana palmipes*. Frost et al. (2006) placed this and other closely related species in the genus *Lithobates*, a name originally proposed by Fitzinger in 1843. Differences in the recommended species name vary according to different classification criteria (e.g., *Rana palmipes* vs. *Lithobates palmipes*), and were thoroughly discussed by Hillis (2007). However, this species may contain cryptic species (Hillis and Wilcox 2005). In this report, we use the binomen *Lithobates palmipes* because it is still widely accepted, though we recognize that an equally valid alternative would be to treat *Lithobates* as a subgenus of *Rana* in order to preserve a long-standing taxonomy (Hillis and Wilcox 2005; AmphibiaWeb 2015).

Previous studies documenting the distribution of *L. palmipes* in South America (e.g., Hillis and De Sá 1988; Canedo and Bilate 2005; Rodrigues et al. 2013)

provided records from Loreto Region, northern Peru (a Región in Peru is equivalent to a federal state; it was formerly known as Departamento), but its distribution along the Peruvian Amazon remains poorly known. It is notable that *L. palmipes* had not been detected in other well-studied lowland sites in Peru, such as Panguana Biological Station (Schlüter et al. 2004), Cuzco Amazónico (Duellman 2005), Los Amigos Biological Station (von May et al. 2009, 2010), and Cocha Cashu Biological Station in Manu National Park (Catenazzi et al. 2013), despite intensive surveys conducted at those sites. In this report, we provide new distributional data for *L. palmipes* in Peru and update the map of its known distribution in South America. We used the morphological diagnoses provided by Hillis and De Sá (1988) to identify specimens and took measurements to the nearest 0.1 mm with calipers under a stereomicroscope.

Our report is based on field observations and the collection of voucher specimens from two localities in southern Peru, and an additional observation (with a photographic voucher) from northern Peru (Fig. 1). On 09 April 2009, a juvenile individual of *L. palmipes* was collected at Lechemayo, Carabaya Province, Puno Region (13°15'7.39"S, 70°20'18.44"O, 390 m elevation). This specimen was deposited in the Herpetological Collection of the Museo de Historia Natural, Universidad Nacional de San Antonio Abad del Cusco, Peru, with voucher number

Correspondence. Email: ¹chara53@hotmail.com; ²amanditadc@gmail.com; ³cinthyasalas84@hotmail.com; ⁴rvonmay@umich.edu (corresponding author).




Fig. 1. Known distribution of *Lithobates palmipes* in South America and location of new records in Peru. White circles represent literature data and red circles indicate the new records in San Martín (<http://www.inaturalist.org/observations/2384262>), Madre de Dios (MUSA-3722, MUSA-3723) and Puno (MHNC-7864) regions.

MHNC-7864 (snout-vent length 57.07 mm). On 28 May 2011, two individuals of *L. palmipes* were collected at the Reserva Comunal Amaraakaeri, Manu Province, Madre de Dios Region (12°46'20.26"S, 70°56'44.56"O, 367 m elevation). Both specimens were found on the ground at a slow-moving stream dissecting a middle floodplain forest scattered with bamboo. These specimens were deposited in the Herpetological Collection of the Museo de Historia Natural (MUSA), Universidad Nacional de San Agustín de Arequipa, Peru, with voucher numbers MUSA-3722 and MUSA-3723 (snout-vent length 119.30 mm and 118.10 mm, respectively; see Table 1 for additional morphometric data). The third locality record is supported by a field observation made by Alessandro Catenazzi on 15 July 2002 at Callanayacu, at the border of the Cordillera Azul National Park, San Martín Region, 320 m (photographic voucher: <http://www.inaturalist.org/observations/2384262>). In addition to the new records reported here, we updated the known distribution of *L. palmipes* in Bolivia using georeferenced data published by Reichle (2007).

This report represents an extension of >175 km of the known geographic range *L. palmipes* in southwestern Amazonia. Furthermore, it is worth noting that two other species of *Lithobates* have been recorded in Peru: *L. bwana* and *L. catesbeianus* (Catenazzi and von May 2014). One of these, the American Bullfrog, *L. catesbeianus*, is an exotic species that has invaded various

South American ecosystems and its presence in northern Peru was confirmed recently (Cossios 2010). As such, this exotic species could pose a threat to many native aquatic-breeding frogs including *L. palmipes*. Given that both *L. palmipes* and *L. catesbeianus* may inhabit similar types of water bodies such as slow-moving streams, seasonal ponds, swamps, and flooded forests (Duellman 1978; La Marca et al. 2010), continuous field assessments in areas where these species have been sighted is a priority (Catenazzi and von May 2014).

Acknowledgments.—We thank Evaristo López Tejada, Director of the Museo de Historia Natural, Universidad Nacional de San Agustín de Arequipa (MUSA), and Juan Carlos Chaparro Auza, Curator of Herpetology at the Museo de Historia Natural, Universidad Nacional de San Antonio Abad del Cusco (MHNC), for providing access to the herpetological collections. Research and collecting permits were issued by the Ministry of Agriculture (Resolución Directoral N° 0398-2010-AG-DGFFS-DGEFFS) and the Ministry of the Environment (Resolución Jefatural de la Reserva Comunal Amaraakaeri N° 001-2010-SERNANP-RCA). We thank Alessandro Catenazzi for sharing one locality record reported in this paper. We thank Jessica Deichmann and an anonymous reviewer for providing helpful comments on our manuscript. RvM thanks the National Science Foundation for a Postdoctoral Research Fellowship in Biology (DBI-1103087).




Fig. 2. Individuals of *Lithobates palmipes* recorded in this study. (A) Adult, female (MUSA-3722) from Reserva Comunal Amara-kaeri, Manu Province, Madre de Dios Region, Peru. (B) Juvenile MHNC-7864 from Lechemayo, Carabaya Province, Puno Region, Perú. Photographs by Roy Santa Cruz (A) and Amanda Delgado (B).

Literature Cited

Acosta-Galvis AR. 1999. Distribución, variación y estatus taxonómico de las poblaciones del complejo *Rana palmipes* (Amphibia: Anura: Ranidae) en Colombia. *Revista de la Academia Colombiana de Ciencias Exactas Físicas y Naturales* 23: 214–224.

AmphibiaWeb. 2015. AmphibiaWeb. Information on amphibian biology and conservation. Berkeley, California, USA. Available: <http://amphibiaweb.org> [Accessed: 20 November 2015].

Canedo C, Bilate M. 2005. Geographic distribution. *Rana palmipes*. *Herpetological Review* 36(3): 334.

Catenazzi A, Lehr E, von May R. 2013. The amphibians and reptiles of Manu National Park and its buffer

zone, Amazon basin and eastern slopes of the Andes, Peru. *Biota Neotropica* 13(4): 269–283.

Catenazzi, A, von May R. 2014. Conservation status of amphibians in Peru. *Herpetological Monographs* 28: 1–23.

Cossios ED. 2010. Vertebrados naturalizados en el Perú: Historia y estado del conocimiento. *Revista Peruana de Biología* 17: 179–189.

Duellman WE. 2005. *Cusco Amazónico: The Lives of Amphibians and Reptiles in an Amazonian Rainforest*. Comstock Publishing Associates, Ithaca, New York, USA. 488 p.

Duellman WE. 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. *Miscellaneous Publications of the University of Kansas* 65: 1–352.

Ferreira AS, Faria RG. 2011. First record of *Lithobates palmipes* (Spix, 1824) (Anura: Ranidae) for the State of Sergipe, Brazil. *Herpetology Notes* 4: 139–140.

Fitzinger LJFJ. 1843. *Systema Reptilium*. Fasciculus Primus. Amblyglossae. Braumüller et Seidel, Wien. 106 p.

Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CFB, de Sá RO, Channing A, Wilkinson M, Donnellan SC, Raxworthy CJ, Campbell JA, Blotto BL, Moler PE, Drewes RC, Nussbaum RA, Lynch JD, Green DM, Wheeler WC. 2006. The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297: 1–370.

Frost DR. 2015. Amphibian Species of the World: An Online Reference. Version 6.0. American Museum of Natural History, New York, New York, USA. Available: <http://research.amnh.org/herpetology/amphibia/index.html> [Accessed: 20 November 2015].

Hillis DM, De Sá R. 1988. Phylogeny and taxonomy of the *Rana palmipes* group (Salientia: Ranidae). *Herpetology Monographs* 2: 1–26.

Table 1. Measurements (in mm) of two adult female individuals of *Lithobates palmipes*. SVL = snout–vent length, TL = tibia length, FL = foot length, HL = head length, HW = head width, ED = eye diameter, TY = tympanum diameter, IOD = interorbital distance, EW = upper eyelid width, IND = internarial distance, E-N = eye-nostril distance.

Character	MUSA-3722	MUSA-3723
SVL	114.2	112.57
TL	57.97	57.72
FL	57.7	56.61
HL	47.13	46.53
HW	46.22	46.4
ED	13.27	12.67
TY	10.37	10.93
IOD	10.78	10.65
EW	9.77	8.8
IND	9.83	9.78
E-N	11.41	11.39

- Hillis DM, Wilcox TP. 2005. Phylogeny of the New World true frogs (*Rana*). *Molecular Phylogenetics and Evolution* 34: 299–314.
- Hillis DM. 2007. Constraints in naming parts of the Tree of Life. *Molecular Phylogenetics and Evolution* 42: 331–338.
- La Marca E, Azevedo-Ramos C, Coloma LA, Ron S, Hardy J. 2010. *Lithobates palmipes*. The IUCN Red List of Threatened Species 2010: e.T58689A11812112. Available <http://dx.doi.org/10.2305/IUCN.UK.20102.RLTS.T58689A11812112.en> [Accessed: 02 November 2015].
- Oliveira RM, Maciel NM, Vaz-Silva W. 2010. New state of *Lithobates palmipes* (Spix, 1824) (Anura: Ranidae) in Brazil. *Herpetology Notes* 3: 277–278.
- Ramalho WP, Viana F, Benevides R, Pralon E, Alves R. 2011. First record of *Lithobates palmipes* (Spix, 1824) (Anura, Ranidae) for the state of Piauí, Northeastern Brazil. *Herpetology Notes* 4: 249–251.
- Reichle S. 2007. Distribution, Diversity, and Conservation Status of Bolivian Amphibians. Unpublished Ph.D. thesis. University of Bonn, Germany. 183 p. + Appendix.
- Rodrigues D, Barros AB, Da Costa J, Almeida EJ (2013) New record and distribution map of *Lithobates palmipes* (Spix, 1824) (Anura, Ranidae) in the state of Mato Grosso, Brazil. *Herpetology Notes* 6: 391–393.
- Santos D, Vaz-Silva W. 2012. Amphibia, Anura, Ranidae, *Lithobates palmipes* (Spix, 1824): New record and geographic distribution map in South America. *CheckList* 8(6): 1,331–1,332.
- Schlüter A, Icochea J, Pérez JM. 2004. Amphibians and reptiles of the lower Río Lullapichis, Amazonian Peru: Updated species list with ecological and biogeographical notes. *Salamandra* 40(2): 141–160.
- von May R, Siu-Ting K, Jacobs JM, Medina-Müller M, Gagliardi G, Rodríguez LO, Donnelly MA. 2009. Species diversity and conservation status of amphibians in Madre de Dios, Perú. *Herpetological Conservation and Biology* 4(1): 14–29.
- von May R, Jacobs JM, Santa-Cruz R, Valdivia J, Huamán J, Donnelly MA. 2010. Amphibian community structure as a function of forest type in Amazonian Peru. *Journal of Tropical Ecology* 26(5): 509–519.



Roy Santa Cruz is an amphibian and reptile researcher at the Herpetological Collection of the Museum of Natural History, Universidad Nacional de San Agustín de Arequipa, Perú (MUSA). His current research interests include taxonomy and ecology of amphibian and reptiles in Peru.



J. Amanda Delgado C. is a Peruvian Biologist associated with the Natural History Museum of Universidad Nacional San Antonio Abad del Cusco (MHNC). Her research interests include taxonomy and diversity of amphibians and reptiles. She is also involved in the creation and management of new Regional Conservation Areas in southeastern Peru.



Cinthya Y. Salas is a Peruvian biologist and a researcher at the Herpetological Collection of the Museum of Natural History of the Universidad Nacional de San Agustín de Arequipa, Peru (MUSA). Her research interests include ecology and conservation of amphibians and reptiles.



Rudolf von May is a postdoctoral research fellow at the Department of Ecology and Evolutionary Biology at the University of Michigan. His current research seeks to understand how amphibian and reptile communities are structured across habitats and elevations, taking into account the phylogenetic relatedness among species present in those communities.