



Cerrophidion wilsoni Jadin, Townsend, Castoe, and Campbell, 2012. The Honduran Montane Pitviper is a priority one species with an EVS of 15, placing it in the high vulnerability category (see this paper). This pitviper is distributed primarily in lower montane rainforest at elevations from 1,400 to 3,491 m, but can occur peripherally in premontane rainforest and pine-oak forest as low as 1,220 m (Jadin et al. 2012). As indicated by Jadin et al. (2012: 10), this snake “occurs in at least 13 isolated highland forest areas across Eastern Nuclear Central America...and all known populations...are found within the borders of Honduras and El Salvador.” This juvenile individual was found in Refugio de Vida Silvestre Texíguat, in north-central Honduras. One of the describers of this taxon is the dedicatee of this paper, and the snake was named in honor of one of the authors. *Photo by Josiah H. Townsend.*

DEDICATION

We are happy to dedicate this paper to our friend and colleague, Josiah H. Townsend, Associate Professor of Biology at Indiana University of Pennsylvania, in Indiana, Pennsylvania. Over the last two decades, since he was a student in one of Larry Wilson's classes, Joe has built an imposing reputation as the principal authority on the herpetofauna of the biogeographically significant Chortís Highlands of northern Central America. During this time he amassed important collections, and their study is demonstrating that the herpetofaunal diversity of this region of Mesoamerica has been seriously underestimated, especially among anurans, salamanders, and squamates. Since 2006, Joe has authored or co-authored the descriptions of 21 new taxa from northern Central America, including one anuran, 10 salamanders, four lizards, and six snakes. He also has produced important summary papers on the Mesoamerican herpetofauna, including several coauthored chapters in the 2010 book *Conservation of Mesoamerican Amphibians and Reptiles* (Wilson et al. 2010; Johnson et al. 2010; Townsend and Wilson 2010a, b), a 2014 paper entitled "Characterizing the Chortís Block Biogeographic Province: geological, physiographic, and ecological associations and herpetofaunal diversity," and a 2016 paper entitled "Amphibians of the Cordillera Nombre de Dios, Honduras: COI barcoding suggests underestimated taxonomic diversity in a threatened endemic fauna." Additionally, Joe co-authored the 2006 book *The Amphibians and Reptiles of the Honduran Mosquitia* (McCranie et al. 2006) and the 2008 book *Amphibians and Reptiles of Cusuco National Park, Honduras* (Townsend and Wilson 2008). During his career Joe has collaborated with a sizable number of colleagues and students to underscore the significant biodiversity of northern Central America, as an important component of the overall Mesoamerican herpetofauna. The collections he assembled in remote regions will continue to provide insights into the phylogenetic relationships and phylogeography of this herpetofauna, and we believe his trajectory will position him as one of the most influential herpetologists of his era.



The frog in this photograph is a partially metamorphosed individual of *Rana (Lithobates) lenca*, recently described as a new species from Honduras in the following paper: Luque-Montes, Ileana, James D. Austin, Kayla D. Weinfurter, Larry David Wilson, Erich P. Hofmann, and

Josiah H. Townsend. 2018. An integrative assessment of the taxonomic status of putative hybrid leopard frogs (Anura: Ranidae) from the Chortís Highlands of Central America, with description of a new species. *Systematics and Biodiversity* 2018: 1–17. This paper is an example of the seminal work being conducted by Joe Townsend and his colleagues, which is exposing the underestimated herpetofaunal diversity of the biogeographically significant Chortís Highlands. The frog was photographed in a shallow pond above the Thomas Cabot Biological Station at an elevation of 1,640 m, within Reserva Biológica Cerro Uyuca in the department of Francisco Morazán. The pond is located in pine forest, along a trail that leads from the biological station to the summit of Cerro Uyuca. This frog is an inhabitant of the "weeping woods" or cloud forest, so beautifully described by Archie Carr in the first chapter of his 1953 book *High Jungles and Low*.



Josiah H. Townsend photographed in 2008 along the Río Arcáqual in Parque Nacional Celaque, on a trail between the visitors' center and the summit of Cerro Celaque. At that time Joe was undergoing a marathon period of fieldwork in an effort to assess the composition, distribution, and conservation status of the amphibian herpetofauna of Honduras. Cerro Celaque is the highest mountain in Honduras, and a site that contains substantial herpetofaunal diversity and endemism. Ongoing research in Joe's lab likely will add to the diversity of salamander species found on this mountain. Joe is considered the principal authority on the herpetofauna of the Chortís Highlands of Central America and is one of the leading investigators employing next-generation techniques in molecular systematics to recover underestimated phylogenetic diversity, especially in threatened endemic herpetofaunas. *Photo by Ileana Luque-Montes.*



The endemic herpetofauna of Central America: a casualty of anthropocentrism

¹Vicente Mata-Silva, ²Dominic L. DeSantis, ³Elí García-Padilla, ⁴Jerry D. Johnson,
and ⁵Larry David Wilson

^{1,2,4}Department of Biological Sciences, The University of Texas at El Paso, El Paso, Texas 79968-0500, USA ³Oaxaca de Juárez, Oaxaca 68023, MEXICO ⁵Centro Zamorano de Biodiversidad, Escuela Agrícola Panamericana Zamorano, Departamento de Francisco Morazán, HONDURAS

Abstract.—The endemic herpetofauna of Central America is of global significance, and currently consists of 623 species, 56.9% of a total herpetofauna of 1,095 species. During the last two years 43 endemic species have been added to this total, and one species has been deleted. The endemic herpetofauna of Central America is distributed unevenly among 10 physiographic regions, ranging from six species in the Yucatan Platform to 254 in the Isthmian Central American highlands. The distributions of close to three quarters of the 623 species are limited to a single physiographic region, and our assessment of their conservation status indicates that about nine-tenths of these species lie within the high vulnerability range of the Environmental Vulnerability Score (EVS). We prioritized the conservation significance of the Central American species by combining the data on physiographic distribution with those of the EVS and recognize 14 priority levels. About eight of every 10 endemic species occupy the first two priority levels, i.e., high vulnerability species limited to one or two physiographic regions. Protecting the endemic component of the Central American herpetofauna is the greatest challenge currently facing conservation professionals working in this region. We conclude that this goal will not be reached until humanity, in general, addresses the issues generated by the widespread adoption of the anthropocentric worldview.

Keywords. Anthropocentric worldview, anurans, caudates, caecilians, conservation significance, endemism, extinction risk, squamates, turtles

Resumen.—La herpetofauna endémica de Centroamérica es de importancia global y actualmente consiste de 623 especies, 56.9% de una herpetofauna total de 1,095 especies. Durante los dos últimos años, 43 especies endémicas han sido agregadas a la lista, y una especie ha sido eliminada. La herpetofauna endémica de Centroamérica está distribuida de forma desigual entre 10 regiones fisiográficas, que va de seis especies en la Plataforma de Yucatán, a 254 en las Tierras Altas del Istmo Centroamericano. Las distribuciones de aproximadamente tres cuartos de las 623 especies están limitadas a una sola región fisiográfica, y nuestra evaluación sobre su estatus de conservación indica que alrededor de nueve décimas de estas especies se localizan dentro de la categoría de vulnerabilidad alta del sistema de puntaje de vulnerabilidad ambiental (EVS). Ordenamos la importancia de conservación de las especies centroamericanas combinando los datos sobre su distribución fisiográfica con los de EVS, y reconocemos 14 niveles de prioridad. Aproximadamente ocho de cada 10 especies endémicas ocupan los dos primeros niveles de prioridad, por ejemplo, especies con vulnerabilidad alta limitadas a una o dos regiones fisiográficas. La protección del componente endémico de la herpetofauna de Centroamérica es el mayor reto para los profesionales de la conservación que trabajan en esta región. Concluimos que esta meta no será lograda hasta que la humanidad en general confronte los problemas generados por una visión antropocéntrica del mundo.

Palabras Claves. Visión antropocéntrica del mundo, anuros, caudados, cecilios, importancia de conservación, endemismo, riesgo de extinción, escamosos, tortugas

Citation: Mata-Silva, V, DeSantis DL, García-Padilla E, Johnson JD, Wilson LD. 2019. The endemic herpetofauna of Central America: a casualty of anthropocentrism. *Amphibian & Reptile Conservation* 13(1) [General Section]: 1–64 (e168).

Copyright: © 2019 Mata-Silva et al. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits unrestricted use for non-commercial and education purposes only, in any medium, provided the original author and the official and authorized publication sources are recognized and properly credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website <amphibian-reptile-conservation.org>.

Received: 30 April 2018; **Accepted:** 10 September 2018; **Published:** 20 January 2019

Correspondence. ¹vmata@utep.edu ²dldesantis@miners.utep.edu ³eligarcia_18@hotmail.com ⁴jjohnson@utep.edu ⁵bufodoc@aol.com

“... losses, whether of species, landscapes, or seascape, -a world without-will become simply the new default. But whether we notice them or not, these losses matter because each diminishes our experiences, pleasures, and possibilities...”

David W. Orr (2016)

Introduction

Johnson et al. (2017) examined the biodiversity and conservation status of the endemic herpetofauna of Mexico, which then consisted of 789 species and presently amounts to 792 species. Johnson et al. (2017) concluded that this endemic herpetofauna is of global significance and has become severely imperiled as a consequence of actions by humans. These authors calculated that the endemic Mexican herpetofauna constituted 61.0% of the total of 1,292 herpetofaunal species in the country.

This paper is a companion piece to Johnson et al. (2017), and places the 623 endemic species in Central America into 14 conservation priority levels. Our approach is the same as that stipulated by Johnson et al. (2017) in their Introduction and section on Biodiversity Decline.

Given that the foundation and conclusions of this paper are based on the results of Johnson et al. (2017), we briefly quote the first four conclusions in that paper (see p. 614), as follows:

“A. The complex interplay among the atmosphere, hydrosphere, and lithosphere allows for the existence of life on planet Earth.”

“B. Humans are faced with the consequences of an interrelated amalgam of global problems of their own making, which impact the atmosphere, hydrosphere, lithosphere, and biosphere. These problems are sufficiently grave to threaten the continued existence of life.”

“C. Biodiversity decline is a problem of global dimensions. This decline impacts life at all levels: from the ecosystem, through the species comprising these ecosystems, to the genes prescribing the traits of these species.”

“D. Throughout its history, life has been subjected to a series of mass extinction episodes that have preceded the current sixth episode of humanity’s design.”

In this paper we underscore the significance of the endemic herpetofauna of Central America and assess its conservation status. The continued existence of this endemic herpetofauna hinges on sustaining the region’s life-support systems, i.e., the group of interacting elements in the atmosphere, hydrosphere, and lithosphere that allow for the maintenance of life on the planet. As a rational species, we believe that humans are responsible for protecting and preserving the diversity of life on the planet, as well as improving the

quality of life for humans and other life forms. For this reason, we are diametrically opposed to the ideas recently promulgated by R. Alexander Pyron in a perspective published in *The Washington Post* (2017), that “we don’t need to save endangered species” because “extinction is part of evolution,” and further that, “the only creatures we should go out of our way to protect are *Homo sapiens*.”

Recent Changes to the Central American Herpetofauna

Even while facing an increasing tempo for global biodiversity decline (Ceballos et al. 2017), herpetologists continue to increase the number of known species of amphibians and reptiles from around the world (see the AmphibiaWeb and Reptile Database websites). Johnson et al. (2015) reported 92 additional taxa to the Central American herpetofaunal list, based on the cutoff date established for the list in Wilson and Johnson (2010). For this paper, which considers the endemic members of the Central American herpetofauna, we added 19 taxa of amphibians and 24 of reptiles to the list presented in Johnson et al. (2015). We list these taxa and their supportive publications below.

Incilius mayordomus—Savage et al. 2013. *Copeia* 2013: 8–12. New species.

Hyalinobatrachium diana—Kubicki et al. 2015. *Zootaxa* 3920: 69–84. New species.

Craugastor gabbi—Arias et al. 2016. *Zootaxa* 4132: 347–363. New species.

Diasporus darienensis—Batista et al. 2016a. *Zoological Journal of the Linnean Society* 178: 267–311. New species.

Diasporus majeensis—Batista et al. 2016a. *Zoological Journal of the Linnean Society* 178: 267–311. New species.

Diasporus pequeno—Batista et al. 2016a. *Zoological Journal of the Linnean Society* 178: 267–311. New species.

Diasporus sapo—Batista et al. 2016a. *Zoological Journal of the Linnean Society* 178: 267–311. New species.

Plectrohyla calvata—McCranie. 2017a. *Mesoamerican Herpetology* 4: 389–401. New species.

Smilisca manisorum—McCranie. 2017c. *Mesoamerican Herpetology* 4: 512–526. Resurrected from synonymy of *Smilisca baudinii*.

Lithobates lenca—Luque-Montes et al. 2018. *Systematics and Biodiversity* 2018: 1–17. New species.

Bolitoglossa aurae—Kubicki and Arias. 2016. *Zootaxa* 4184: 329–346. New species.

Bolitoglossa chucutaniensis—Batista et al. 2014. *Mesoamerican Herpetology* 1: 96–121. New species.

- Cryptotriton xucaneborum*—Rovito et al. 2015. *Zoological Journal of the Linnean Society* 175: 150–166. New species.
- Nototriton costaricense*—Arias and Kubicki. 2018. *Zootaxa* 4369: 487–500. New species.
- Nototriton nelsoni*—Townsend. 2016. *Zootaxa* 4196: 511–528. New species.
- Nototriton oreadorum*—Townsend. 2016. *Zootaxa* 4196: 511–528. New species.
- Oedipina berlini*—Kubicki. 2016. *Mesoamerican Herpetology* 3: 819–840. New species.
- Oedipina capitalina*—Solís et al. 2016. *Salamandra* 42: 125–133. New species.
- Oedipina salvadorensis*—Brodie et al. 2012. *Journal of Herpetology* 46: 233–240. Resurrected from synonymy of *Oedipina taylora*.
- Celestus laf*—Lotzkat et al. 2016. *Mesoamerican Herpetology* 3: 962–975. New species.
- Mesaspis cuchumatanus*—Solano-Zavaleta et al. 2016. *Journal of Herpetology* 50: 327–335. New species
- Mesaspis salvadorensis*—Solano-Zavaleta and Nieto-Montes de Oca. 2018. *Molecular Phylogenetics and Evolution* 120: 16–27. Elevated from subspecies to species level.
- Dactyloa brooksi*—Poe and Ryan. 2017. *Amphibian & Reptile Conservation* 11: 1–16. Resurrected from the synonymy of *Dactyloa insignis*.
- Dactyloa kathydayae*—Poe and Ryan. 2017. *Amphibian & Reptile Conservation* 11: 1–16. New species.
- Dactyloa maia*—Batista et al. 2015b. *Zootaxa* 4039: 57–84. New species.
- Dactyloa savagei*—Poe and Ryan. 2017. *Amphibian & Reptile Conservation* 11: 1–16. New species.
- Norops elcopeensis*—Poe et al. 2015. *Amphibian & Reptile Conservation* 9: 1–13. New species.
- Norops mcraniei*—Köhler et al. 2016. *Mesoamerican Herpetology* 3: 8–41. New species.
- Norops oxylophus*—McCranie and Köhler. 2015. *Bulletin of the Museum of Comparative Zoology* SPS(1): 1–292. Recognition as distinct from *Norops lionotus*.
- Norops wilsoni*—Köhler et al. 2016. *Mesoamerican Herpetology* 3: 8–41. New species.
- Lepidoblepharis emberawoundule*—Batista et al. 2015a. *Zootaxa* 3994: 187–221. New species.
- Lepidoblepharis rufigularis*—Batista et al. 2015a. *Zootaxa* 3994: 187–221. New species.
- Lepidoblepharis victormartinezi*—Batista et al. 2015a. *Zootaxa* 3994: 187–221. New species.
- Ameiva fuliginosa*—McCranie and Gotte. 2014. *Proceedings of the Biological Society of Washington* 127: 543–556. Elevation from subspecies to species level.
- Holcusus miadis*—Meza-Lázaro et al. 2015. *Zoological Journal of the Linnean Society* 2015: 1–22. Elevation from subspecies to species level.
- Tantilla berguidoi*—Batista et al. 2016b. *Mesoamerican Herpetology* 3: 949–960. New species
- Tantilla excelsa*—McCranie and Smith. 2017. *Herpetologica* 73: 338–348. New species.
- Tantilla gottei*—McCranie and Smith. 2017. *Herpetologica* 73: 338–348. New species.
- Tantilla stenigrammi*—McCranie and Smith. 2017. *Herpetologica* 73: 338–348. New species.
- Rhadinella lisyae*—McCranie. 2017b. *Mesoamerican Herpetology* 4: 243–253. New species.
- Epictia martinezi*—Wallach. 2016. *Mesoamerican Herpetology* 3: 216–374. New species.
- Epictia pauldwyeri*—Wallach. 2016. *Mesoamerican Herpetology* 3: 216–374. New species.
- Bothriechis nubestris*—Doan et al. 2016. *Zootaxa* 4138: 271–290. New species.

The Distributional Status of *Dipsas viguieri*

Peters (1960) reviewed the taxonomic status of members of the subfamily Dipsadinae, and placed nine species in the *Dipsas articulata* group, collectively distributed from western and southeastern Mexico to northwestern Ecuador. Peters' view of the species-level relationships among the members of the *articulata* group was impacted by the paucity of specimens of each taxon known at that time, and he noted that only minimal scale and color differences separated certain species. Peters (1960) indicated the range of *D. viguieri* as the Pacific coast of Panama.

Pérez Santos and Moreno (1988) reported on a specimen of *D. gracilis* from the Pacific coast of Colombia (see discussion below), and Pérez Santos (1999) noted the occurrence of *D. viguieri* from both versants of Panama, including a specimen from the province of Bocas del Toro in the western part of the country. Subsequently, Köhler (2001; 2003; 2008) noted the range of *Dipsas viguieri* as eastern Panama and western Colombia but did not provide additional information. Cadle (2005), in a paper on the systematics of the *Dipsas oreas* complex, tentatively referred to a specimen (FMNH 74376) from northwestern Colombia near the Panama border as *D. viguieri*, which previously had been identified as *D. gracilis*. Nonetheless, Cadle (2005) stated that these two taxa were not distinguishable by any reported characteristics, and on p. 128 noted that, “Without additional study, I am unable to adequately differentiate *Dipsas viguieri* (eastern Panama and northern Chocó, Colombia) and *D. gracilis* (western Ecuador and extreme northern Peru).” Further, based on an examination of morphological characters, Cadle indicated geography as



Plate 1. *Atelopus varius* (Lichtenstein and Martens, 1856). The Harlequin Frog is a priority ten species with an EVS of 11, distributed on both versants of the cordilleras of Costa Rica and western Panama (Frost 2018). This individual is from one of three known surviving populations of this species, and in Costa Rica it is being surveyed near Uvita, in the province of Puntarenas. *Photo by César Barrio-Amorós.*



Plate 2. *Incilius holdridgei* (Taylor, 1952). Holdridge's Toad is a priority one species with an EVS of 14, which is restricted in distribution to Volcán Barva, Costa Rica (Frost 2018). This individual was located in Alto del Roble, in the province of Heredia. *Photo by Víctor Acosta-Chaves.*



Plate 3. *Incilius melanochlorus* (Cope, 1877). The Wet Forest Toad is a priority eight species with an EVS of 12, with a distribution on the Atlantic versant of Costa Rica and adjacent Panama, and likely in adjacent Nicaragua (Frost 2018). This individual was found in Centro Soltis, San Isidro de Peñas Blancas, in the province of Alajuela, Costa Rica. *Photo by Víctor Acosta-Chaves.*

the only currently reliable means of assigning names to these species. Similarly, in discussing a number of poorly-known *Dipsas* from South America, Harvey (2008) commented that he was unable to distinguish *D. viguieri* from *D. gracilis*, and thus did not include *D. viguieri* in his key but noted that the *D. articulata* complex requires further study.

Jaramillo et al. (2010) and Johnson et al. (2015) regarded *Dipsas viguieri* as endemic to Panama. Wallach et al. (2014: 235), however, considered *D. viguieri* as occupying “Eastern Panama (Canal Zone, Darién, Panamá) and Colombia (? Chocó, Piura), NSL–60 m.” In their *D. gracilis* account, however, these authors noted, “Colombian record doubtful *vide* Cadle (2005: 123): possibly *D. viguieri*. *Dipsas gracilis* and *D. viguieri* possibly conspecific *vide* Harvey (2008: 429).” Wallach et al. (2014: 232), however, apparently confused the information provided by Cadle (2005), as the FMNH specimen tentatively was referred to *D. viguieri* and not *D. gracilis*. Finally, Ray (2017) indicated the range of *D. viguieri* as eastern Panama to northwestern Colombia.

The historical timeline for information on the distribution and taxonomic status of *Dipsas viguieri* has been unclear, as different workers have maintained that this species is endemic to Panama or occurs in both Panama and Colombia. In the absence of a definitive analysis involving morphological and molecular approaches, for the purpose of this paper we are considering *D. viguieri* as not endemic to Panama.

Global Status of the Central American Herpetofauna

As with the Mexican herpetofauna (Johnson et al. 2017), the Central American herpetofauna also is highly diverse, consisting of 60 families, 214 genera, and 1,095 species (Table 1), organized into six orders (Anura, Caudata, Gymnophiona, Crocodylia, Squamata, and Testudines). The level of herpetofaunal diversity in Central America is intermediate between that found in Mexico and North America (United States–Canada). The number of species in the United States–Canada is the same as Johnson et al. (2017) reported, i.e., 650 (Center for North American Herpetology website; accessed 9 December 2017). Johnson et al. (2017) reported the number of species in Mexico as 1,292.

Even though the number of herpetofaunal species occurring in Central America is intermediate between that found in the United States–Canada and Mexico, Central America contains about 8.5 times the number of taxa by area as found in Mexico, and 155.6 times the number found in the United States–Canada. Thus, the relative degree of biodiversity is significantly higher in Central America when compared to that in Mexico and the United States–Canada.

If we consider Central America as a single region in our analysis (i.e., not one divided into seven countries), then its herpetofauna also is significant when compared to that of other areas in Latin America. With respect to amphibians, the 509 species occurring in Central America is the fifth

largest in Latin America (amphibiaweb.com; 15 April 2018), and is closest to that for the country of Ecuador, at 562. The area/species ratio for Ecuador however, is 504.6, compared to 998.0 for Central America.

Considering the numbers of crocodylian, squamate, and turtle species, the 586 species in Central America is comparable to that recorded for the neighboring country of Colombia, which is 611 (reptile-database.org; accessed 29 December 2017). Colombia, however, with an area of 1,141,748 km², is 2.25 times the size of Central America, which contains an area of 507,966 km² (www.Oei.es/historico/cultura2/Colombia/03.htm; accessed 29 December 2017). Thus, the area/species ratio for Colombia is 1,868.7, compared to 868.0 for that of Central America. Only Brazil (799) and Colombia in South America contain more species than Central America (reptile-database.org; accessed 29 December 2017).

Endemism within the Central American Herpetofauna

The proportion of herpetofaunal endemism in Central America is slightly less than in Mexico, the other major segment of Mesoamerica. The percentage in Central America is 56.9 (Table 2) compared to 61.1 in Mexico (Johnson et al. 2017). This percentage in Central America is based on an endemic herpetofauna of 623 species and a total herpetofauna of 1,095 species (Table 2). Both of the comparable figures for the Mexican herpetofauna are higher, i.e., 789 and 1,292 (Johnson et al. 2017). As noted by Johnson et al. (2015: 26), “Mesoamerica is one of the world’s most important biodiversity reservoirs, and Central America contains a substantial component of that region’s herpetofauna.” We illustrate the breakdown of the total and endemic components of the Central American herpetofauna in Fig. 1. This graph shows the close correspondence between the endemic and total number of salamander species, the relatively distant correspondence between the endemic and total number of squamate species, and the intermediate correspondence between the two figures for anurans (Fig. 1).

Of the 60 families represented in Central America, 38 (63.3%) contain endemic species (Table 2). This leaves 22 families with no endemic representation, including the anuran families Aromobatidae, Hemiphractidae, and Rhinophrynidae, the crocodylian families Alligatoridae and Crocodylidae, the squamate families Amphisbaenidae, Hoplocercidae, Polychrotidae, Xenosauridae, Boidae, Charinidae, Loxocemidae, Natricidae, Sibynophiidae, and Tropidophiidae, and the turtle families Cheloniidae, Chelydridae, Dermatemydidae, Dermochelyidae, Emydidae, Staurotypidae, and Testudinidae. In Central America these are small-content families, with species numbers ranging from one to five (Table 2). The families with endemic representation have total numbers ranging from one to 166; the endemic numbers vary from one to 143 (Table 2).




Fig. 1. Graph comparing the endemic and total number of species for the Central American herpetofauna, arranged by order.

Table 1. Diversity of the Central American herpetofauna at the familial, generic, and specific levels.

Orders	Families	Genera	Species
Anura	14	58	328
Caudata	1	7	166
Gymnophiona	2	4	15
Crocodylia	2	2	3
Squamata	32	130	560
Testudines	9	13	23
Totals	60	214	1,095

Table 2. Degree of endemism of the Central American herpetofauna at the species level, arranged by family.

Family	Total Number of Species	Number of Endemic Species	Percentage of Endemism
Aromobatidae	1	—	—
Bufo	40	24	60.0
Centrolenidae	15	4	26.7
Craugastoridae	102	77	75.5
Dendrobatidae	21	14	66.7
Eleutherodactylidae	16	10	62.5
Hemiphractidae	3	—	—
Hylidae	92	52	60.9
Leptodactylidae	9	1	11.1
Microhylidae	9	1	11.1
Phyllomedusidae	7	2	28.6
Pipidae	1	1	100

The endemic herpetofauna of Central America

Table 2 (continued). Degree of endemism of the Central American herpetofauna at the species level, arranged by family.

Family	Total Number of Species	Number of Endemic Species	Percentage of Endemism
Ranidae	11	6	54.5
Rhinophrynidae	1	—	—
Subtotals	328	192	58.5
Plethodontidae	166	143	86.1
Subtotals	166	143	86.1
Caeciliidae	7	3	42.9
Dermophiidae	8	4	50.0
Subtotals	15	7	46.7
Totals	509	342	67.2
Alligatoridae	1	—	—
Crocodylidae	2	—	—
Subtotals	3	—	—
Amphisbaenidae	2	—	—
Anguidae	32	25	78.1
Corytophanidae	9	1	11.1
Dactyloidae	104	74	71.2
Eublepharidae	2	1	50.0
Gymnophthalmidae	14	3	21.4
Helodermatidae	2	1	50.0
Hoplocercidae	2	—	—
Iguanidae	9	7	77.8
Mabuyidae	5	4	80.0
Phrynosomatidae	17	2	11.8
Phyllodactylidae	5	3	60.0
Polychrotidae	1	—	—
Scincidae	3	1	33.3
Sphaerodactylidae	22	13	59.1
Sphenomorphidae	4	1	25.0
Teiidae	16	5	31.3
Xantusiidae	4	2	50.0
Xenosauridae	1	—	—
Anomalepididae	3	1	33.3
Boidae	5	—	—
Charinidae	2	—	—
Colubridae	79	30	38.0
Dipsadidae	145	77	53.1
Elapidae	19	6	31.6
Leptotyphlopidae	6	3	50.0
Loxocemidae	1	—	—
Natricidae	5	—	—
Sibynophiidae	2	—	—
Tropidophiidae	1	—	—
Typhlopidae	5	3	60.0
Viperidae	33	16	48.5
Subtotals	560	279	49.8
Cheloniidae	4	—	—

Table 2 (continued). Degree of endemism of the Central American herpetofauna at the species level, arranged by family.

Family	Total Number of Species	Number of Endemic Species	Percentage of Endemism
Chelydridae	2	—	—
Dermatemydidae	1	—	—
Dermochelyidae	1	—	—
Emydidae	2	—	—
Geoemydidae	5	1	20.0
Kinosternidae	4	1	25.0
Staurotypidae	3	—	—
Testudinidae	1	—	—
Subtotals	23	2	8.7
Totals	586	281	48.0
Sum Totals	1,095	623	56.9

Of the 14 anuran families with representatives in Central America, 11 contain endemic species, which include 192 (58.5%) of the total of 328 species (Table 2). Of these 11 families, the largest numbers of endemics are 77 in the Craugastoridae and 52 in the Hylidae. Other than in the Pipidae, with one total and one endemic species (100%), as might be expected, the percentage of endemism is next highest in the Craugastoridae (75.5%), but the third highest is in the Dendrobatidae (66.7%), and not the Hylidae (60.9%). The value for the Eleutherodactylidae (62.5%) also is higher than that for the Hylidae. The remaining families contain from one to 24 endemic species (Table 2).

A single family of salamanders, the Plethodontidae, occurs in Central America. The percentage of endemism (86.1%) is amazingly high and is the highest in all the 38 families represented (Table 2).

The endemic species of caecilians (seven) make up less than one-half (46.7%) of the total number of 15 in Central America. Three of the endemics are caeciliids and four are dermophiids.

None of the three species of crocodylians in Central America is endemic. *Crocodylus acutus* and *Caiman crocodylus* rather are among the naturally most broadly distributed herpetofaunal species in the Western Hemisphere.

The squamates are the most speciose group of herpetofaunal organisms in Central America, with 560 species distributed among 32 families (Table 2). Only the endemic proportions of turtles (8.7%) and crocodylians (0.0%) are lower than those of squamates (49.8%). The endemic squamates are more or less evenly divided between the lizards (143) and snakes (136). Of the 19 families of lizards with representatives in Central America, 15 contain endemic species (78.9%). The largest numbers of endemic squamate species are found within the families Dactyloidae (74) and Dipsadidae (77). The next largest number of endemic lizards (25) is allocated to the family Anguidae. The remaining 12 lizard families contain only one to 13 endemic species. The percentage of endemism among the lizard families ranges from 11.1% in the Corytophanidae to 80.0% in the Mabuyidae (Table 2). Thirteen families of snakes are represented in Central America, of which seven

contain endemic species (53.8%). The greatest numbers of endemic species are in the families Dipsadidae (77) and Colubridae (30). The next largest number (16) lies within the family Viperidae; the remaining four snake families contain from one to six endemic species (Table 2). The percentage of endemism among the snake families ranges from 31.6% in the Elapidae to 60.0% in the Typhlopidae (Table 2).

The percentage of endemism among turtles is very low, with only two such species among a total of 23 (8.7%), one each in the families Geoemydidae and Kinosternidae (Table 2).

At the ordinal level, the percentage of endemism is highest among the salamanders at 86.1%, the same figure as for the family Plethodontidae, since it is the only family of salamanders found in Central America (Table 2). The percentage is lowest among the crocodylians at 0.0%. Intermediate values are evident for the anurans (58.5%), squamates (49.8%), and caecilians (46.7%).

The level of herpetofaunal endemism in Central America is comparable to that found in the other portions of the North American continent, i.e., Mexico, as well as the United States–Canada (Table 3). The overall level for Central America, however, is slightly lower (56.9%) than for either Mexico (61.5%) or the United States–Canada (61.2%).

Although, as expected, the total number of herpetofaunal species is lower in the United States–Canada than to the south in Mesoamerica (Table 3); even with the greater area of the two northern nations, the level of endemism still is impressive. Of the total of 650 species in the United States–Canada, 398 are endemic, for a percentage of endemism of 61.2%. Notably, this level of endemism is based heavily on the amphibians, especially the salamanders. The proportion of amphibian endemism is more than 10 points higher in the United States–Canada (78.6%) than in Mexico (68.3%) or Central America (67.2%). The percentage of endemism is higher for both anurans and salamanders in the United States–Canada (64.4% and 86.4%, respectively) than for these two groups in either Mexico (60.0% and 82.8%, respectively) or Central America (58.5% and 86.1%, respectively). Significantly, the level of amphibian endemism in all three regions heavily depends on the



Plate 4. *Craugastor sandersoni* (Schmidt, 1941). Sanderson's Rainfrog is a priority two species with an EVS of 19, which is distributed on the "Caribbean slopes of the Maya Mountains in east-central Belize southward to the Montañas del Mico in eastern Guatemala and westward into the Sierra de Santa Cruz, the eastern portion of the Sierra de las Minas, and the foothills of the northern Alta Verapaz" (Frost 2018). This individual was found in Montañas del Mico, Guatemala. *Photo by Sean Michael Rovito.*



Plate 5. *Oophaga granulifera* (Taylor, 1958). The Granular Poison Frog is a priority two species with an EVS of 17, distributed in the Golfo Dulce region of Pacific coastal Costa Rica and presumably in adjacent Panama (Frost 2018). This individual was encountered in Ciudad Cortéz de Osa, in the province of Puntarenas, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 6. *Oophaga pumilio* (Schmidt, 1857). The Strawberry Poison Frog is a priority one species with an EVS of 16, and its distribution extends along the Atlantic versant from "eastern Nicaragua...south through the lowlands of Costa Rica and northwestern Panama (Savage 2002: 388). This individual came from the mainland in the province of Bocas del Toro, Panama. *Photo by Abel Batista.*



Plate 7. *Oophaga vicentei* (Jungfer, Weyoldt, and Juraska, 1996). This dendrobatid frog is a priority two species with an EVS of 16, distributed on the Atlantic versant "of the provinces of Veraguas and Coclé and the upper reaches of Pacific versant in Coclé, central Panama" (Frost 2018). This individual is from Santa Fé National Park, in the province of Veraguas, Panama. *Photo by Abel Batista.*



Plate 8. *Phyllobates lugubris* (Schmidt, 1857). The Lovely Poison Frog is a priority one species with an EVS of 17, which ranges along the Atlantic versant from "extreme southeastern Nicaragua to northwestern Panama; a single record from just west of the Panama Canal" (Savage 2002: 390). This individual was found in the Donoso region, in the province of Colón, Panama. *Photo by Abel Batista.*



Plate 9. *Diasporus hylaeformis* (Cope, 1876). The Pico Blanco Robber Frog is a priority one species with an EVS of 17, with a distribution in the highlands of the cordilleras of Costa Rica and Panama (Frost 2018). This individual came from Alto del Roble, in the province of Heredia, Costa Rica. *Photo by Víctor Acosta-Chaves.*

Table 3. Total number of species, endemic species, and relative endemism within the herpetofaunal groups in Central America, North America (United States–Canada), and Mexico. Data for Central America from this paper, for North America from CNAH (www.cnah.org; accessed 9 January 2018), and for Mexico from updated figures in Johnson et al. (2017).

Herpetofaunal Groups	Total Species in Central America	Endemic Species in Central America	Relative Endemism in Central America (%)	Total Species in North America	Endemic Species in North America	Relative Endemism in North America (%)	Total Species in Mexico	Endemic Species in Mexico	Relative Endemism in Mexico (%)
Anurans	328	192	58.5	104	67	64.4	247	148	60.0
Salamanders	166	143	86.1	191	165	86.4	151	125	82.8
Caecilians	15	7	46.7	—	—	—	3	1	33.3
Subtotals	509	342	67.2	295	232	78.6	401	274	68.3
Crocodylians	3	—	—	2	1	50.0	3	—	—
Squamates	560	279	49.8	287	117	40.8	863	517	60.0
Turtles	23	2	8.3	66	48	72.7	51	20	39.2
Subtotals	586	281	48.0	355	166	46.8	917	537	58.6
Totals	1,095	623	56.9	650	398	61.2	1,318	811	61.5

salamanders, i.e., over 80%. Curiously enough, such high incidences of salamander species-level endemism are not correlated with the incidence of family-level endemism, which decreases markedly from the United States–Canada through Mexico to Central America. The 191 salamander species in the United States–Canada are organized within eight families, but with the majority allocated to the family Plethodontidae (Center for North American Herpetology website; accessed 2 January 2018). Four of these eight families, the Amphiumidae, Cryptobranchidae, Proteidae, and Rhyacotritonidae occur no farther south than the United States. Four families, the Ambystomatidae, Plethodontidae, Salamandridae, and Sirenidae are represented in Mexico, but with the greatest number of species in the Plethodontidae, as in the United States–Canada. Only a single family, the Plethodontidae, is found in Central America. In the United States–Canada, 145 of the 191 species of salamanders are in the family Plethodontidae (75.9%). Of the 151 species found in Mexico 130 (86.1%) are in the Plethodontidae (Wilson et al., 2017); the remaining 21 species are allocated to three families (see above). Finally, of the 166 species of salamanders found in Central America, all are in the Plethodontidae; obviously the percentage of occupancy is 100%.

Unlike the situation among the amphibians, the level of endemism among the crocodylians, squamates, and turtles is significantly lower in all three regions dealt with in Table 3. The level of endemism among these taxa is lowest in the United States–Canada (46.8%), next lowest in Central America (48.0%), and highest in Mexico (58.6%). Given that squamates constitute the largest group, compared to the other two, the same pattern would be expected for them as for the entire group. Thus, the level of squamate endemism is lowest in the United States–Canada (40.8%), intermediate in Central America (49.8%), and highest in Mexico (60.0%).

The differential between the percentages of endemism for amphibians versus the remainder of the herpetofauna

(Table 3) increases from that seen in Mexico (9.7%), through Central America (19.2%), to the United States–Canada (31.8%). Thus, in all three regions amphibians contribute more to the degree of endemism than the remainder of the herpetofauna (Table 3).

Physiographic Distribution of the Endemic Central American Herpetofauna

Given the considerable global significance of the diversity and endemism of the Central American herpetofauna, as documented above, it is of paramount importance to protect its elements. As an initial step to determine the distributional patterns of these organisms in Central America, we collated the available information on the occurrence of the members of the herpetofauna among the 10 physiographic regions traditionally recognized in this portion of Mesoamerica (Campbell, 1999; Wilson and Johnson, 2010; Fig. 2). Six of these regions occupy the lowlands of Central America, including the Yucatan Platform, the Caribbean lowlands of eastern Guatemala and northern Honduras, the Caribbean lowlands from Nicaragua to Panama, the Pacific lowlands from eastern Chiapas to south-central Guatemala, the Pacific lowlands from southeastern Guatemala to northwestern Costa Rica, and the Pacific lowlands from central Costa Rica through Panama (Table 4). Four regions are found in the highlands of Central America, including the western nuclear Central American highlands, the eastern nuclear Central American highlands, the Isthmian Central American highlands, and the highlands of eastern Panama (Table 4). We document the distribution of the 623 endemic members of the Central American herpetofauna among the 10 physiographic regions in Table 4, and summarize these data in Table 5 and Fig. 3.

The total number of the endemic species distributed within the 10 physiographic regions ranges from a low of six in the Yucatan Platform to 254 in the Isthmian Central American highlands (Table 5). The mean regional occupancy figure



Fig. 2. Physiographic regions of Central America, after Campbell (1999). Abbreviations are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform.

Table 4. Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
Anura (192 species)											
Bufo											
Bufo											
<i>Atelopus certus</i>			+	+							2
<i>Atelopus chiriquiensis</i>			+								1
<i>Atelopus chirripoensis</i>			+								1
<i>Atelopus limosus</i>									+		1
<i>Atelopus senex</i>			+								1
<i>Atelopus varius</i>			+						+	+	3
<i>Atelopus zeteki</i>			+							+	2

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Incilius aucoinae</i>										+	1
<i>Incilius chompipe</i>			+								1
<i>Incilius epioticus</i>			+								1
<i>Incilius fastidiosus</i>			+								1
<i>Incilius guanacaste</i>			+								1
<i>Incilius holdridgei</i>			+								1
<i>Incilius ibarrae</i>	+	+									2
<i>Incilius karenlipsae</i>			+								1
<i>Incilius leucomyos</i>		+				+					2
<i>Incilius majordomus</i>			+								1
<i>Incilius melanochlorus</i>			+								1
<i>Incilius periglenes</i>			+								1
<i>Incilius peripatetes</i>			+								1
<i>Incilius porteri</i>		+									1
<i>Incilius signifer</i>			+							+	2
<i>Rhinella centralis</i>							+			+	2
<i>Rhinella chrysophora</i>		+									1
Centrolenidae (4 species)											
<i>Cochranella granulosa</i>		+	+			+	+			+	5
<i>Hyalinobatrachium diana</i>			+				+				2
<i>Hyalinobatrachium talamancae</i>			+								1
<i>Hyalinobatrachium vireovittatum</i>			+								1
Craugastoridae (77 species)											
<i>Craugastor adamastus</i>	+										1
<i>Craugastor anciano</i>		+									1
<i>Craugastor andi</i>			+								1
<i>Craugastor angelicus</i>			+								1
<i>Craugastor aphanus</i>	+										1
<i>Craugastor aurilegulus</i>		+				+					2
<i>Craugastor azueroensis</i>										+	1
<i>Craugastor bocourti</i>	+										1
<i>Craugastor bransfordii</i>			+				+				2
<i>Craugastor campbelli</i>	+					+					2
<i>Craugastor catalinae</i>			+								1
<i>Craugastor chac</i>	+	+				+					3
<i>Craugastor charadra</i>		+				+					2
<i>Craugastor chingopetaca</i>							+				1
<i>Craugastor chrysozetetes</i>		+									1
<i>Craugastor coffeus</i>		+									1
<i>Craugastor cruzi</i>		+									1
<i>Craugastor cuaquero</i>			+								1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Craugastor cyanochthebius</i>		+									1
<i>Craugastor daryi</i>	+										1
<i>Craugastor emcelae</i>			+								1
<i>Craugastor emleni</i>		+									1
<i>Craugastor epochthidius</i>		+				+					2
<i>Craugastor escoces</i>			+								1
<i>Craugastor evanesco</i>			+				+				2
<i>Craugastor fecundus</i>		+				+					2
<i>Craugastor fleischmanni</i>			+								1
<i>Craugastor gabbi</i>			+								1
<i>Craugastor gollmeri</i>			+				+			+	3
<i>Craugastor gulosus</i>			+								1
<i>Craugastor inachus</i>	+										1
<i>Craugastor jota</i>			+								1
<i>Craugastor laevisimus</i>		+				+			+		3
<i>Craugastor lauraster</i>		+				+					2
<i>Craugastor megacephalus</i>		+	+			+	+				4
<i>Craugastor melanostictus</i>			+								1
<i>Craugastor merendonensis</i>						+					1
<i>Craugastor milesi</i>		+									1
<i>Craugastor mimus</i>		+	+			+	+				4
<i>Craugastor monnichorum</i>			+								1
<i>Craugastor myllomyllon</i>	+										1
<i>Craugastor nefrens</i>		+									1
<i>Craugastor noblei</i>		+	+			+	+			+	5
<i>Craugastor obesus</i>			+				+				2
<i>Craugastor olanchano</i>		+									1
<i>Craugastor omoaensis</i>		+									1
<i>Craugastor pechorum</i>		+				+					2
<i>Craugastor persimilis</i>			+				+				2
<i>Craugastor phasma</i>			+								1
<i>Craugastor podiciferus</i>			+								1
<i>Craugastor polyptychus</i>							+				1
<i>Craugastor punctariolus</i>			+								1
<i>Craugastor ranoides</i>			+				+		+	+	4
<i>Craugastor rayo</i>			+								1
<i>Craugastor rhyacobatrachus</i>			+								1
<i>Craugastor rivulus</i>	+										1
<i>Craugastor rostralis</i>	+	+									2
<i>Craugastor rugosus</i>			+							+	2
<i>Craugastor sabrinus</i>	+					+					2

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Craugastor saltuarius</i>		+									1
<i>Craugastor sandersoni</i>	+					+					2
<i>Craugastor stadelmani</i>		+									1
<i>Craugastor stejnegerianus</i>			+						+	+	3
<i>Craugastor tabasarae</i>			+								1
<i>Craugastor talamancae</i>								+			1
<i>Craugastor taurus</i>										+	1
<i>Craugastor trachydermus</i>	+										1
<i>Craugastor underwoodi</i>			+								1
<i>Craugastor xucanebi</i>	+										1
<i>Pristimantis adnus</i>				+							1
<i>Pristimantis altae</i>			+						+		2
<i>Pristimantis caryophyllaceus</i>			+						+		2
<i>Pristimantis cerasinus</i>			+			+	+				3
<i>Pristimantis museosus</i>			+								1
<i>Pristimantis pardalis</i>			+					+		+	3
<i>Pristimantis pirrensis</i>				+							1
<i>Strabomantis laticorpus</i>				+							1
Dendrobatidae (14 species)											
<i>Ameerega maculata?</i>											?
<i>Andinobates claudiae</i>								+			1
<i>Andinobates geminisae</i>								+			1
<i>Colostethus latinasus</i>				+							1
<i>Ectopoglossus astralogaster</i>				+							1
<i>Ectopoglossus isthminus</i>				+							1
<i>Oophaga arborea</i>			+								1
<i>Oophaga granulifera</i>								+		+	2
<i>Oophaga pumilio</i>								+			1
<i>Oophaga speciosa</i>			+								1
<i>Oophaga vicentei</i>				+			+				2
<i>Phyllobates lugubris</i>								+			1
<i>Phyllobates vittatus</i>										+	1
<i>Silverstoneia flotator</i>			+				+			+	3
Eleutherodactylidae (10 species)											
<i>Diasporus citrinobapheus</i>			+								1
<i>Diasporus darienensis</i>				+							1
<i>Diasporus diastema</i>			+			+	+			+	4
<i>Diasporus hylaeformis</i>			+								1
<i>Diasporus igneus</i>			+								1
<i>Diasporus majeensis</i>				+							1
<i>Diasporus pequeno</i>				+							1
<i>Diasporus sapo</i>				+							1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Diasporus tigrillo</i>							+				1
<i>Diasporus ventrimaculatus</i>			+								1
Hyliidae (52 species)											
<i>Atlantihyla panchoi</i>	+					+					2
<i>Atlantihyla spinipollex</i>		+				+					2
<i>Bromeliahyla melacaena</i>		+									1
<i>Dryophytes bocourti</i>	+										1
<i>Duellmanohyla legleri</i>			+								1
<i>Duellmanohyla lythrodes</i>							+				1
<i>Duellmanohyla rufiocularis</i>			+								1
<i>Duellmanohyla salvadorensis</i>	+	+									2
<i>Duellmanohyla salvavida</i>		+				+					2
<i>Duellmanohyla soralia</i>		+				+					2
<i>Duellmanohyla uranochroa</i>			+				+				2
<i>Ecnomiohyla bailarina</i>				+			+				2
<i>Ecnomiohyla fimbrimembra</i>			+				+				2
<i>Ecnomiohyla minera</i>	+										1
<i>Ecnomiohyla rabborum</i>			+								1
<i>Ecnomiohyla salvaje</i>		+									1
<i>Ecnomiohyla sukia</i>			+				+				2
<i>Ecnomiohyla thysanota</i>				+							1
<i>Ecnomiohyla veraguensis</i>			+								1
<i>Exerodonta catracha</i>		+									1
<i>Exerodonta perkinsi</i>	+										1
<i>Hyloscirtus colymba</i>			+	+							2
<i>Isthmohyla angustilineata</i>			+								1
<i>Isthmohyla calypso</i>			+								1
<i>Isthmohyla debilis</i>			+								1
<i>Isthmohyla graceae</i>			+								1
<i>Isthmohyla infucata</i>			+								1
<i>Isthmohyla insolita</i>		+									1
<i>Isthmohyla lancasteri</i>			+				+				2
<i>Isthmohyla picadoi</i>			+								1
<i>Isthmohyla pictipes</i>			+								1
<i>Isthmohyla pseudopuma</i>			+								1
<i>Isthmohyla rivularis</i>			+								1
<i>Isthmohyla tica</i>			+								1
<i>Isthmohyla xanthosticta</i>			+								1
<i>Isthmohyla zeteki</i>			+								1
<i>Plectrohyla calvata</i>		+									1
<i>Plectrohyla chrysopleura</i>		+									1

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Plectrohyla dasypus</i>		+									1
<i>Plectrohyla exquisita</i>		+									1
<i>Plectrohyla glandulosa</i>	+										1
<i>Plectrohyla pokomchi</i>	+										1
<i>Plectrohyla psiloderma</i>		+									1
<i>Plectrohyla quecchi</i>	+										1
<i>Plectrohyla tecunumani</i>	+										1
<i>Plectrohyla teuchestes</i>	+										1
<i>Ptychohyla dendrophasma</i>	+										1
<i>Ptychohyla hypomykter</i>	+	+					+				3
<i>Quilticohyla sanctaerucis</i>	+										1
<i>Scinax altae</i>										+	1
<i>Smilisca manisorum</i>									+		1
<i>Smilisca puma</i>									+		1
Leptodactylidae (1 species)											
<i>Leptodactylus silvanimbus</i>		+									1
Microhylidae (1 species)											
<i>Hypopachus pictiventris</i>									+		1
Phyllomedusidae (2 species)											
<i>Agalychnis annae</i>			+						+		2
<i>Agalychnis saltator</i>			+				+		+		3
Pipidae (1 species)											
<i>Pipa myersi</i>										+	1
Ranidae (6 species)											
<i>Lithobates juliani</i>	+										1
<i>Lithobates lenca</i>		+									1
<i>Lithobates miadis</i>									+		1
<i>Lithobates taylori</i>									+		1
<i>Lithobates vibicarius</i>			+								1
<i>Lithobates warszewitschii</i>		+	+	+		+					4
Caudata (143 species)											
Plethodontidae (143 species)											
<i>Bolitoglossa alvaradoi</i>			+						+		2
<i>Bolitoglossa anthracina</i>			+								1
<i>Bolitoglossa aurae</i>			+								1
<i>Bolitoglossa aureogularis</i>			+								1
<i>Bolitoglossa bramei</i>			+								1
<i>Bolitoglossa carri</i>		+									1
<i>Bolitoglossa cataguana</i>		+									1
<i>Bolitoglossa celaque</i>		+									1
<i>Bolitoglossa centenorum</i>	+										1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Bolitoglossa cerroensis</i>			+								1
<i>Bolitoglossa chucantiensis</i>				+							1
<i>Bolitoglossa colonnea</i>			+	+			+			+	4
<i>Bolitoglossa compacta</i>			+								1
<i>Bolitoglossa conanti</i>		+									1
<i>Bolitoglossa copia</i>			+								1
<i>Bolitoglossa cuchumatana</i>	+										1
<i>Bolitoglossa cuna</i>							+				1
<i>Bolitoglossa daryorum</i>	+										1
<i>Bolitoglossa decora</i>		+									1
<i>Bolitoglossa diaphora</i>		+									1
<i>Bolitoglossa diminuta</i>			+								1
<i>Bolitoglossa dofleini</i>	+	+			+	+					4
<i>Bolitoglossa dunni</i>		+									1
<i>Bolitoglossa epimela</i>			+								1
<i>Bolitoglossa eremia</i>	+										1
<i>Bolitoglossa gomezi</i>			+								1
<i>Bolitoglossa gracilis</i>			+								1
<i>Bolitoglossa heiroreias</i>		+									1
<i>Bolitoglossa helmrichi</i>	+										1
<i>Bolitoglossa huehuetenanguensis</i>	+										1
<i>Bolitoglossa indio</i>							+				1
<i>Bolitoglossa insularis</i>		+									1
<i>Bolitoglossa jacksoni</i>	+										1
<i>Bolitoglossa jugivagans</i>			+								1
<i>Bolitoglossa kamuk</i>			+								1
<i>Bolitoglossa kaqchikelorum</i>	+										1
<i>Bolitoglossa la</i>	+										1
<i>Bolitoglossa lignicolor</i>			+							+	2
<i>Bolitoglossa longissima</i>		+									1
<i>Bolitoglossa magnifica</i>			+								1
<i>Bolitoglossa marmorea</i>			+								1
<i>Bolitoglossa meliana</i>	+										1
<i>Bolitoglossa minutula</i>			+								1
<i>Bolitoglossa mombachoensis</i>		+									1
<i>Bolitoglossa morio</i>	+										1
<i>Bolitoglossa nigrescens</i>			+								1
<i>Bolitoglossa ninadormida</i>	+										1
<i>Bolitoglossa nussbaumi</i>	+										1
<i>Bolitoglossa nympa</i>	+	+				+					3
<i>Bolitoglossa obscura</i>			+								1

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Bolitoglossa odonnelli</i>	+					+					2
<i>Bolitoglossa omniumsanctorum</i>	+										1
<i>Bolitoglossa oresbia</i>		+									1
<i>Bolitoglossa pacaya</i>	+										1
<i>Bolitoglossa pesrubra</i>			+								1
<i>Bolitoglossa porrasorum</i>		+									1
<i>Bolitoglossa psephena</i>	+										1
<i>Bolitoglossa pygmaea</i>			+								1
<i>Bolitoglossa robinsoni</i>			+								1
<i>Bolitoglossa robusta</i>			+								1
<i>Bolitoglossa salvinii</i>	+								+		2
<i>Bolitoglossa schizodactyla</i>			+				+			+	3
<i>Bolitoglossa sombra</i>			+								1
<i>Bolitoglossa sooyorum</i>			+								1
<i>Bolitoglossa splendida</i>			+								1
<i>Bolitoglossa striatula</i>		+	+			+	+		+		5
<i>Bolitoglossa subpalmata</i>			+								1
<i>Bolitoglossa suchitanensis</i>	+										1
<i>Bolitoglossa synoria</i>		+									1
<i>Bolitoglossa taylori</i>				+							1
<i>Bolitoglossa tenebrosa</i>	+										1
<i>Bolitoglossa tica</i>			+								1
<i>Bolitoglossa tzultacaj</i>	+										1
<i>Bolitoglossa xibalba</i>	+										1
<i>Bolitoglossa zacapensis</i>	+										1
<i>Cryptotriton monzoni</i>	+										1
<i>Cryptotriton nasalis</i>	+	+									2
<i>Cryptotriton necopinus</i>		+									1
<i>Cryptotriton sierraminensis</i>	+										1
<i>Cryptotriton veraepacis</i>	+										1
<i>Cryptotriton xucaneborum</i>	+										1
<i>Dendrotriton bromeliacius</i>	+										1
<i>Dendrotriton chujorum</i>	+										1
<i>Dendrotriton cuchumatanus</i>	+										1
<i>Dendrotriton kekchiorum</i>	+										1
<i>Dendrotriton rabbi</i>	+										1
<i>Dendrotriton sanctibarbarus</i>		+									1
<i>Nototriton abscondens</i>			+								1
<i>Nototriton barbouri</i>		+									1
<i>Nototriton brodiei</i>	+										1
<i>Nototriton costaricense</i>			+								1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Nototriton gamezi</i>			+								1
<i>Nototriton guanacaste</i>			+								1
<i>Nototriton lignicola</i>		+									1
<i>Nototriton limnospectator</i>		+									1
<i>Nototriton major</i>			+								1
<i>Nototriton matama</i>			+								1
<i>Nototriton mime</i>		+									1
<i>Nototriton nelsoni</i>		+									1
<i>Nototriton oreadorum</i>		+									1
<i>Nototriton picadoi</i>			+								1
<i>Nototriton picucha</i>		+									1
<i>Nototriton richardi</i>			+								1
<i>Nototriton saslaya</i>		+									1
<i>Nototriton stuarti</i>	+										1
<i>Nototriton tapanti</i>			+								1
<i>Nototriton tomamorum</i>		+									1
<i>Oedipina alfaroi</i>			+							+	2
<i>Oedipina alleni</i>			+							+	2
<i>Oedipina altura</i>			+								1
<i>Oedipina berlini</i>			+								1
<i>Oedipina capitalina</i>		+									1
<i>Oedipina carablanca</i>										+	1
<i>Oedipina chortiorum</i>		+									1
<i>Oedipina collaris</i>										+	1
<i>Oedipina cyclocauda</i>										+	1
<i>Oedipina fortunensis</i>			+								1
<i>Oedipina geophyra</i>		+									1
<i>Oedipina gracilis</i>										+	1
<i>Oedipina grandis</i>			+								1
<i>Oedipina ignea</i>	+	+									2
<i>Oedipina kasios</i>		+									1
<i>Oedipina koehleri</i>		+									1
<i>Oedipina leptopoda</i>		+									1
<i>Oedipina maritima</i>										+	1
<i>Oedipina motaguae</i>							+				1
<i>Oedipina nica</i>		+									1
<i>Oedipina nimaso</i>			+								1
<i>Oedipina pacificensis</i>										+	1
<i>Oedipina paucidentata</i>			+								1
<i>Oedipina petiola</i>		+									1
<i>Oedipina poelzi</i>			+								1

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Oedipina pseudouniformis</i>			+				+		+		3
<i>Oedipina quadra</i>						+					1
<i>Oedipina salvadorensis</i>									+		1
<i>Oedipina savagei</i>			+								1
<i>Oedipina stenopodia</i>	+										1
<i>Oedipina stuarti</i>		+							+		2
<i>Oedipina taylora</i>		+									1
<i>Oedipina tomasi</i>		+									1
<i>Oedipina tzutujilorum</i>	+										1
<i>Oedipina uniformis</i>			+								1
<i>Pseudoeurycea exspectata</i>	+										1
Gymnophiona (7 species)											
Caeciliidae (3 species)											
<i>Caecilia volceni</i>										+	1
<i>Oscaecilia elongata</i>										+	1
<i>Oscaecilia osae</i>										+	1
Dermophiidae (4 species)											
<i>Dermophis costaricensis</i>			+								1
<i>Dermophis gracilior</i>			+								1
<i>Dermophis occidentalis</i>			+							+	2
<i>Gymnopsis multiplicata</i>		+	+			+	+			+	5
Squamata (279 Species)											
Anguidae (25 species)											
<i>Abronia anzuetoi</i>	+										1
<i>Abronia aurita</i>	+										1
<i>Abronia campbelli</i>	+										1
<i>Abronia fimbriata</i>	+										1
<i>Abronia frosti</i>	+										1
<i>Abronia gaiophasma</i>	+										1
<i>Abronia meledona</i>	+										1
<i>Abronia montecristoi</i>		+									1
<i>Abronia salvadorensis</i>		+									1
<i>Abronia vasconcelosii</i>	+										1
<i>Celestus adercus</i>			+								1
<i>Celestus atitlanensis</i>	+	+									2
<i>Celestus bivittatus</i>		+									1
<i>Celestus cyanochloris</i>			+								1
<i>Celestus hylaius</i>							+				1
<i>Celestus laf</i>			+								1
<i>Celestus montanus</i>		+									1
<i>Celestus orobius</i>			+								1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Celestus scansorius</i>		+									1
<i>Coloptychon rhombifer</i>										+	1
<i>Diploglossus bilobatus</i>			+							+	3
<i>Diploglossus montisilvestris</i>				+							1
<i>Mesaspis cuchumatanus</i>	+										1
<i>Mesaspis monticola</i>			+								1
<i>Mesaspis salvadorensis</i>		+									1
Corytophanidae (1 species)											
<i>Basiliscus plumifrons</i>			+			+	+		+	+	5
Dactyloidae (74 species)											
<i>Dactyloa brooksi</i>			+	+			+			+	4
<i>Dactyloa casildae</i>			+								1
<i>Dactyloa ginaelisae</i>			+								1
<i>Dactyloa ibanezi</i>			+							+	2
<i>Dactyloa insignis</i>			+				+			+	3
<i>Dactyloa kathydayae</i>			+								1
<i>Dactyloa kunayalae</i>			+				+				2
<i>Dactyloa maia</i>				+							1
<i>Dactyloa microtus</i>			+								1
<i>Dactyloa savagei</i>			+						+		2
<i>Norops alocomyos</i>			+								1
<i>Norops altae</i>			+								1
<i>Norops amplisquamosus</i>		+									1
<i>Norops apletophallus</i>			+	+			+			+	4
<i>Norops aquaticus</i>			+							+	2
<i>Norops benedikti</i>			+								1
<i>Norops bicaorum</i>						+					1
<i>Norops campbelli</i>	+										1
<i>Norops carpenteri</i>		+					+				2
<i>Norops charlesmyersi</i>		+								+	2
<i>Norops cobanensis</i>	+										1
<i>Norops cryptolimifrons</i>			+				+				2
<i>Norops cupreus</i>		+	+			+	+	+	+	+	6
<i>Norops cusuco</i>		+									1
<i>Norops datzorum</i>			+								1
<i>Norops elcopeensis</i>			+							+	2
<i>Norops fortunensis</i>			+								1
<i>Norops fungosus</i>			+								1
<i>Norops gruuo</i>			+								1
<i>Norops Haguei</i>	+										1
<i>Norops heterophilidotus</i>		+									1

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Norops humilis</i>			+	+			+			+	4
<i>Norops intermedius</i>			+								1
<i>Norops johnmeyeri</i>		+									1
<i>Norops kemptoni</i>			+								1
<i>Norops kreutzi</i>		+									1
<i>Norops leditzigorum</i>			+								1
<i>Norops limifrons</i>		+	+			+	+		+	+	6
<i>Norops lionotus</i>			+				+			+	3
<i>Norops loveridgei</i>		+				+					2
<i>Norops macrophallus</i>	+							+	+		3
<i>Norops magnaphallus</i>			+								1
<i>Norops marsupialis</i>			+							+	2
<i>Norops mcraniei</i>	+	+									2
<i>Norops monteverde</i>			+								1
<i>Norops morazani</i>		+									1
<i>Norops muralla</i>		+									1
<i>Norops ocelloscapularis</i>		+									1
<i>Norops osa</i>										+	1
<i>Norops oxylophus</i>		+	+							+	4
<i>Norops pachypus</i>			+								1
<i>Norops pijolensis</i>		+									1
<i>Norops polylepis</i>			+							+	2
<i>Norops pseudokemptoni</i>			+								1
<i>Norops pseudopachypus</i>			+								1
<i>Norops purpurgularis</i>		+									1
<i>Norops quaggulus</i>		+				+	+				3
<i>Norops roatanensis</i>						+					1
<i>Norops rubribarbaris</i>		+									1
<i>Norops salvini</i>			+								1
<i>Norops sminthus</i>		+									1
<i>Norops tenorioensis</i>			+								1
<i>Norops townsendi</i>										+	1
<i>Norops triumphalis</i>										+	1
<i>Norops tropidolepis</i>			+								1
<i>Norops utilis</i>						+					1
<i>Norops villai</i>							+				1
<i>Norops wampuensis</i>						+					1
<i>Norops wellbornae</i>	+	+						+	+		4
<i>Norops wermuthi</i>		+									1
<i>Norops wilsoni</i>		+				+					2
<i>Norops woodi</i>			+								1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Norops yoroensis</i>		+									1
<i>Norops zeus</i>		+				+					2
Eublepharidae (1 species)											
<i>Coleonyx mitratus</i>		+	+			+			+	+	5
Gymnophthalmidae (3 species)											
<i>Bachia blairi</i>										+	1
<i>Echinosaura panamensis</i>			+						+	+	3
<i>Echinosaura apodema</i>			+							+	2
Helodermatidae (1 species)											
<i>Heloderma charlesbogerti</i>	+							+			2
Iguanidae (7 species)											
<i>Ctenosaura bakeri</i>						+					1
<i>Ctenosaura flavidorsalis</i>		+				+			+		3
<i>Ctenosaura melanosterna</i>		+				+					2
<i>Ctenosaura oedirhina</i>						+					1
<i>Ctenosaura palearis</i>	+										1
<i>Ctenosaura praeocularis</i>		+							+		2
<i>Ctenosaura quinquecarinata</i>		+							+		2
Mabuyidae (4 species)											
<i>Marisora alliacea</i>									+		1
<i>Marisora magnacornae</i>									+		1
<i>Marisora roatanae</i>						+					1
<i>Marisora unimarginata</i>			+	+			+		+	+	5
Phrynosomatidae (2 species)											
<i>Sceloporus lunaei</i>	+					+					2
<i>Sceloporus malachiticus</i>		+	+			+					3
Phyllodactylidae (3 species)											
<i>Phyllodactylus insularis</i>						+					1
<i>Phyllodactylus palmeus</i>						+					1
<i>Phyllodactylus paralepis</i>						+					1
Scincidae (1 species)											
<i>Mesoscincus managuae</i>		+							+		2
Sphaerodactylidae (13 species)											
<i>Lepidoblepharis emberawoundule</i>				+							1
<i>Lepidoblepharis rufigularis</i>				+							1
<i>Lepidoblepharis victormartinezi</i>			+				+				2
<i>Sphaerodactylus alphus</i>						+					1
<i>Sphaerodactylus dunni</i>						+					1
<i>Sphaerodactylus graptolaemus</i>										+	1
<i>Sphaerodactylus guanaje</i>						+					1
<i>Sphaerodactylus homolepis</i>							+				1

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Sphaerodactylus leonardovaldesi</i>						+					1
<i>Sphaerodactylus millepunctatus</i>	+	+			+	+	+		+		6
<i>Sphaerodactylus pacificus</i>										+	1
<i>Sphaerodactylus poindexteri</i>						+					1
<i>Sphaerodactylus rosaurae</i>						+					1
Sphenomorphidae (1 species)											
<i>Scincella rara</i>								+			1
Teiidae (5 species)											
<i>Cnemidophorus duellmani</i>										+	1
<i>Cnemidophorus ruatanus</i>						+	+				2
<i>Holcosus leptophrys</i>			+	+			+			+	4
<i>Holcosus miadis</i>							+				1
<i>Holcosus quadrilineatus</i>			+	+			+			+	4
Xantusiidae (2 species)											
<i>Lepidophyma mayae</i>	+				+						2
<i>Lepidophyma reticulatum</i>			+							+	2
Anomalepididae (1 species)											
<i>Helminthophis frontalis</i>			+				+				2
Colubridae (30 species)											
<i>Dendrophidion apharocybe</i>		+	+			+	+				4
<i>Dendrophidion crybelum</i>			+								1
<i>Dendrophidion paucicarinatum</i>			+								1
<i>Dendrophidion rufiterminorum</i>	+	+	+			+	+				5
<i>Drymobius melanotropis</i>			+			+	+				3
<i>Leptodrymus pulcherrimus</i>		+				+		+	+		4
<i>Leptophis nebulosus</i>			+			+	+			+	4
<i>Mastigodryas alternatus</i>		+	+			+	+		+	+	6
<i>Mastigodryas dorsalis</i>	+	+									2
<i>Oxybelis wilsoni</i>						+					1
<i>Scolecophis atrocinctus</i>		+	+						+		3
<i>Tantilla albiceps</i>							+				1
<i>Tantilla armillata</i>	+	+	+						+		4
<i>Tantilla bairdi</i>	+										1
<i>Tantilla berguidoii</i>				+							1
<i>Tantilla brevicauda</i>								+	+		2
<i>Tantilla excelsa</i>		+				+					2
<i>Tantilla gottei</i>		+									1
<i>Tantilla hendersoni</i>	+										1
<i>Tantilla jani</i>	+							+			2
<i>Tantilla lempira</i>		+									1
<i>Tantilla olympia</i>		+									1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Tantilla psittaca</i>						+					1
<i>Tantilla ruficeps</i>			+				+			+	3
<i>Tantilla stenigrammi</i>		+									1
<i>Tantilla taeniata</i>	+								+		2
<i>Tantilla tecta</i>					+						1
<i>Tantilla tritaeniata</i>						+					1
<i>Tantilla vermiformis</i>									+		1
<i>Trimorphodon quadruplex</i>	+	+				+	+		+		5
Dipsadidae (77 species)											
<i>Adelphicos daryi</i>	+										1
<i>Adelphicos ibarrorum</i>	+										1
<i>Adelphicos veraepacis</i>	+										1
<i>Atractus darienensis</i>										+	1
<i>Atractus depressiocellus</i>										+	1
<i>Atractus hostiltractus</i>							+				1
<i>Atractus imperfectus</i>				+							1
<i>Chapinophis xanthocheilus</i>	+										1
<i>Coniophanes joanae</i>				+							1
<i>Crisantophis nevermanni</i>		+	+					+	+		4
<i>Cubophis brooksi</i>						+					1
<i>Dipsas articulata</i>			+				+		+		3
<i>Dipsas bicolor</i>		+				+	+				3
<i>Dipsas nicholsi</i>										+	1
<i>Dipsas tenuissima</i>										+	1
<i>Enulius bifoveatus</i>						+					1
<i>Enulius roatanensis</i>						+					1
<i>Geophis bellus</i>				+							1
<i>Geophis brachycephalus</i>			+				+			+	3
<i>Geophis championi</i>			+								1
<i>Geophis damiani</i>		+									1
<i>Geophis downsi</i>			+								1
<i>Geophis dunni</i>		+									1
<i>Geophis fulvoguttatus</i>		+									1
<i>Geophis godmani</i>			+								1
<i>Geophis hoffmanni</i>		+	+			+	+		+	+	6
<i>Geophis nephodrymus</i>		+									1
<i>Geophis ruthveni</i>			+				+				2
<i>Geophis talamancae</i>			+								1
<i>Geophis tectus</i>			+				+				2
<i>Geophis zeledoni</i>			+								1
<i>Hydromorphus concolor</i>	+	+	+			+	+		+	+	7

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Hydromorphus dunni</i>			+								1
<i>Imantodes phantasma</i>				+							1
<i>Leptodeira rhombifera</i>	+	+	+			+	+	+	+	+	8
<i>Leptodeira rubricata</i>										+	1
<i>Ninia celata</i>			+								1
<i>Ninia espinali</i>		+									1
<i>Ninia maculata</i>		+	+			+	+			+	5
<i>Ninia pavimentata</i>	+	+									2
<i>Ninia psephota</i>			+								1
<i>Omoadiphas aurula</i>		+									1
<i>Omoadiphas cannula</i>		+									1
<i>Omoadiphas texiguatensis</i>		+									1
<i>Rhadinaea calligaster</i>			+								1
<i>Rhadinaea pulveriventris</i>			+								1
<i>Rhadinaea sargenti</i>				+			+				2
<i>Rhadinaea stadelmani</i>	+										1
<i>Rhadinaea vermiculaticeps</i>			+				+			+	3
<i>Rhadinella anachoreta</i>	+	+			+	+					4
<i>Rhadinella hempsteadae</i>	+										1
<i>Rhadinella lisyae</i>		+									1
<i>Rhadinella montecristi</i>	+	+									2
<i>Rhadinella pegosalyta</i>		+									1
<i>Rhadinella pilonaorum</i>	+	+									2
<i>Rhadinella rogerromani</i>		+									1
<i>Rhadinella serperaster</i>			+								1
<i>Rhadinella tolpanorum</i>		+									1
<i>Sibon anthracops</i>	+	+							+		3
<i>Sibon argus</i>			+	+			+			+	4
<i>Sibon carri</i>	+	+							+		3
<i>Sibon lamari</i>							+				1
<i>Sibon longifrenis</i>		+	+			+	+				4
<i>Sibon manzanaresi</i>						+					1
<i>Sibon merendonensis</i>		+									1
<i>Sibon miskitus</i>						+					1
<i>Sibon noalamina</i>			+								1
<i>Sibon perissostichon</i>			+								1
<i>Trimetopon barbouri</i>			+				+			+	3
<i>Trimetopon gracile</i>			+								1
<i>Trimetopon pliolepis</i>			+				+				2
<i>Trimetopon simile</i>			+				+				2
<i>Trimetopon slevini</i>			+								1

The endemic herpetofauna of Central America

Table 4 (continued). Distribution of the 623 endemic herpetofaunal species in Central America among the 10 physiographic regions. Abbreviations for the regions are as follows: CG = western nuclear Central American highlands; CGU = Pacific lowlands from eastern Chiapas to south-central Guatemala; CP = Pacific lowlands from central Costa Rica through Panama (area includes associated Pacific islands); CRP = Isthmian Central American highlands; EP = highlands of eastern Panama; GCR = Pacific lowlands from southeastern Guatemala to northwestern Costa Rica; GH = Caribbean lowlands of eastern Guatemala and northern Honduras (area includes associated Caribbean islands); HN = eastern nuclear Central American highlands; NP = Caribbean lowlands from Nicaragua to Panama (area includes associated Caribbean islands); and YP = Yucatan Platform. ? = species known from indeterminate locality (see text).

	Physiographic Regions of Central America										Totals
	CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP	
<i>Trimetopon viquezi</i>							+				1
<i>Urotheca guentheri</i>		+	+			+	+			+	5
<i>Urotheca myersi</i>			+								1
<i>Urotheca pachyura</i>			+							+	2
Elapidae (6 species)											
<i>Micrurus alleni</i>			+			+	+			+	4
<i>Micrurus hippocrepis</i>	+					+					2
<i>Micrurus mosquitensis</i>							+				1
<i>Micrurus ruatanus</i>						+					1
<i>Micrurus stewarti</i>				+			+			+	3
<i>Micrurus stuarti</i>	+										1
Leptotyphlopidae (3 species)											
<i>Epictia ater</i>	+	+	+			+			+		5
<i>Epictia martinezi</i>		+									1
<i>Epictia pauldwyeri</i>										+	1
Typhlopidae (3 species)											
<i>Amerotyphlops costaricensis</i>		+	+			+	+				4
<i>Amerotyphlops stadelmani</i>		+				+					2
<i>Typhlops tycherus</i>		+									1
Viperidae (16 species)											
<i>Agkistrodon howardgloydi</i>	+				+			+	+		4
<i>Atropoides indomitus</i>		+									1
<i>Atropoides picadoi</i>			+							+	3
<i>Bothriechis guifarroi</i>		+									1
<i>Bothriechis lateralis</i>			+								1
<i>Bothriechis marchi</i>		+									1
<i>Bothriechis nigroviridis</i>			+								1
<i>Bothriechis nubestris</i>			+								1
<i>Bothriechis supraciliaris</i>			+							+	2
<i>Bothriechis thalassinus</i>		+									1
<i>Cerrophidion sasai</i>			+								1
<i>Cerrophidion wilsoni</i>		+									1
<i>Lachesis melanocephala</i>			+							+	2
<i>Lachesis stenophrys</i>							+			+	2
<i>Porthidium porrasi</i>								+			1
<i>Porthidium volcanicum</i>										+	1
Testudines (2 species)											
Geoemydidae (1 species)											
<i>Rhinoclemmys funerea</i>						+	+				2
Kinosternidae (1 species)											
<i>Kinosternon angustipons</i>							+				1




Fig. 3. Graph indicating the number and percentage of Central American endemic species in each of the 10 physiographic regions recognized.

Table 5. Distributional summary of herpetofaunal families containing priority level one species in Central America, among the 10 physiographic regions. See Table 4 for explanation of abbreviations.

Families	Number of Species	Physiographic Regions									
		CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP
Bufonidae	12	—	1	9	—	—	—	1	—	—	1
Centrolenidae	2	—	—	2	—	—	—	—	—	—	—
Craugastoridae	50	9	12	20	3	—	1	3	—	—	2
Dendrobatidae	10	—	—	2	3	—	—	4	—	—	1
Eleutherodactylidae	9	—	—	4	4	—	—	1	—	—	—
Hylidae	31	7	8	11	1	—	—	3	—	—	1
Leptodactylidae	1	—	1	—	—	—	—	—	—	—	—
Microhylidae	1	—	—	—	—	—	—	1	—	—	—
Pipidae	1	—	—	—	—	—	—	—	—	—	1
Ranidae	3	—	1	1	—	—	—	1	—	—	—
Subtotals	120	16	23	49	11	—	1	14	—	—	6
Plethodontidae	127	34	35	45	2	—	2	7	—	1	1
Subtotals	127	34	35	45	2	—	2	7	—	1	1
Caecilidae	3	—	—	—	—	—	—	—	—	—	3
Dermophiidae	2	—	—	2	—	—	—	—	—	—	—
Subtotals	5	—	—	2	—	—	—	—	—	—	3
Totals	252	50	58	96	13	—	3	21	—	1	10
Anguidae	23	9	6	5	1	—	—	1	—	—	1
Dactyloidae	46	2	14	22	—	—	4	1	—	—	3
Gymnophthalmidae	1	—	—	—	—	—	—	—	—	—	1

The endemic herpetofauna of Central America

Table 5 (continued). Distributional summary of herpetofaunal families containing priority level one species in Central America, among the 10 physiographic regions. See Table 4 for explanation of abbreviations.

Families	Number of Species	Physiographic Regions									
		CG	HN	CRP	EP	YP	GH	NP	CGU	GCR	CP
Iguanidae	3	1	—	—	—	—	2	—	—	—	—
Mabuyidae	3	—	—	—	—	—	1	2	—	—	—
Phyllodactylidae	3	—	—	—	—	—	3	—	—	—	—
Sphaerodactylidae	11	—	—	—	2	—	6	1	—	—	2
Sphenomorphidae	1	—	—	—	—	—	—	1	—	—	—
Teiidae	2	—	—	—	—	—	—	1	—	—	1
Colubridae	15	2	4	2	1	1	3	1	—	1	—
Dipsadidae	48	4	13	14	4	—	5	3	—	—	5
Elapidae	3	1	—	—	—	—	1	1	—	—	—
Leptotyphlopidae	2	—	1	—	—	—	—	—	—	—	1
Typhlopidae	1	—	1	—	—	—	—	—	—	—	—
Viperidae	11	—	5	4	—	—	—	—	—	1	1
Subtotals	173	19	44	47	8	1	25	12	—	2	15
Kinosternidae	1	—	—	—	—	—	—	1	—	—	—
Subtotals	1	—	—	—	—	—	—	1	—	—	—
Totals	174	19	44	47	8	1	25	13	—	2	15
Sum Totals	426	69	102	143	21	1	28	34	—	3	25

is 94.3. Four of the regional values lie above or close to the mean figure, as follows: western nuclear Central American highlands (CG; 117), eastern nuclear Central American highlands (HN; 178), Isthmian Central American highlands (CRP; 254), Caribbean lowlands of eastern Guatemala and northern Honduras (GH; 94), and Caribbean lowlands from Nicaragua to Panama (NP; 123). Given these species numbers, the five regions with values above or close to the mean are the most significant for conservation remediation.

The other five of the regional values lie below the mean figure, as follows: highlands of eastern Panama (EP; 37), Yucatan Platform (YP; 6), Pacific lowlands from eastern Chiapas to south-central Guatemala (CGU; 9), Pacific lowlands from southeastern Guatemala to northwestern Costa Rica (GCR; 39), and Pacific lowlands from Central Costa Rica through Panama (CP; 86). Even though these values are relatively low, collectively they amount to 177 species, 28.4% of the total of 623 endemic species; thus, they also are of considerable importance.

The five regions containing the highest numbers of endemic species include three in highland and two in lowland areas. The five regions with the lowest numbers include one in highland and four in lowland areas. The numbers in the four highland regions range from 37 to 254, and in the six lowland regions from six to 86.

Obviously, the 623 Central American endemic species are distributed unevenly throughout the 10 physiographic regions we recognize. In order to examine their distribution, we constructed a table indicating the total number of regions inhabited by the component species (Table 6). The regions, listed in order of their total number of constituent species, range from six in the Yucatan Platform to 254 in the Isthmian

Central American highlands. The number of physiographic regions occupied by these species ranges from one to eight, and their corresponding number of species also decreases markedly (Table 6). Thus, 450 species occupy a single region, with the numbers ranging from one in the Yucatan Platform to 154 in the Isthmian Central American highlands; no single-region species are present in the Pacific lowlands from eastern Chiapas to south-central Guatemala. At the opposite extreme, a single species (*Leptodeira rhombifera*) occupies eight regions, and one species (*Hydromorphus concolor*) inhabits seven regions. The single-region species comprise the most speciose categories for seven of the 10 physiographic regions (Table 6). The three exceptions are subhumid regions on the Atlantic (Yucatan Platform) and Pacific versants (Pacific lowlands from eastern Chiapas to south-central Guatemala and Pacific lowlands from southeastern Guatemala to northwestern Costa Rica).

The 450 single-region species comprise 72.2% of the 623 Central American endemic species. The 95 two-region species contribute 15.2% of the total number. Together, the single-region and two-region species constitute 545 taxa, 87.5% of the total. Thus, only 78 of the remaining species occupy from three to eight regions. This feature is of tremendous conservation significance for Central America, and we review this matter in greater detail below.

Conservation Status of the Endemic Central American Herpetofauna

In a previous paper on the Mexican endemic herpetofauna (Johnson et al. 2017), we utilized the Environmental Vulnerability Score (EVS) system of conservation



Plate 10. *Diasporus ventrimaculatus* Chaves, García-Rodríguez, Mora, and Leal, 2009. This frog is a priority one species “known only from the Valle del Silencio on the Caribbean versant of the Cordillera de Talamanca, Limon Province, Costa Rica” (Frost 2018). This individual was observed in Valle del Silencio, Parque Internacional La Amistad, in the province of Limón, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 11. *Duellmanohyla rufioculis* (Taylor, 1952). This treefrog is a priority one species with an EVS of 14, which ranges on both “the Caribbean and Pacific slopes of the mountains of Costa Rica” (Frost 2018). This individual came from Centro Soltis, San Isidro de Peñas Blancas, in the province of Alajuela, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 12. *Isthmohyla lancasteri* (Barbour, 1928). Lancaster’s Treefrog is a priority two species with an EVS of 14, which occurs in “the Cordillera de Talamanca of Costa Rica and western Panama” (Frost 2018). This individual was found in Guayacán, in the province of Limón, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 13. *Plectrohyla pokomchi* Duellman and Campbell, 1984. The Rio Sanaja Spikethumb Frog is a priority eight species with an EVS of 13, which is distributed in “the Sierra de las Minas and the contiguous Sierra de Xucaneb in central and eastern Guatemala” (Frost 2018). This individual came from Purullhá, in the department of Baja Verapaz, Guatemala. *Photo by Andres Novales.*



Plate 14. *Ptychohyla legleri* (Taylor, 1958). Legler’s Stream Frog is a priority one species with an EVS of 14, which is found on the “Pacific slopes of the Sierra de Talamanca [of] eastern Costa Rica and western Panama” (Frost 2018). This individual was located in Alfombra de Pérez Zeledón, in the province of San José, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 15. *Smilisca puma* (Cope, 1885). The Tawny Smilisca is a priority one species with an EVS of 14, distributed on the “Caribbean lowlands of Costa Rica and adjacent Nicaragua” (Frost 2018). This individual was encountered in La Selva Biological Station, in the province of Heredia, Costa Rica. *Photo by Víctor Acosta-Chaves.*

The endemic herpetofauna of Central America

Table 6. Number of endemic species in each of the 10 physiographic regions inhabited in Central America. See Table 4 for explanation of abbreviations.

Physiographic Regions	Number of endemic species in each physiographic region								Totals
	1	2	3	4	5	6	7	8	
YP	1	1	—	3	—	1	—	—	6
CGU	—	3	1	4	—	—	—	1	9
EP	23	4	1	6	1	—	1	—	36
GCR	4	8	10	6	6	5	2	1	42
CP	28	20	14	9	8	4	2	1	86
GH	28	25	12	11	11	5	2	1	95
CG	77	23	6	5	3	1	1	1	117
NP	36	35	22	13	11	5	2	1	125
HN	102	32	12	11	9	5	2	1	174
CRP	153	40	21	16	11	4	2	1	248
Totals	452	191	99	84	60	30	14	8	—

assessment. Along with various other authors, we have been involved with a series of papers published on the Mesoamerican herpetofauna since 2013 (see Johnson et al. 2017, for a listing) including a recent paper on the herpetofauna of the Mexican state of Puebla (Woolrich-Piña et al. 2017). Herein, we use the same system to evaluate the conservation status of the 623 species comprising the Central American herpetofauna. In calculating the EVS for these species, we used the scores included in Johnson et al. (2015), supplemented by the scores we determined for the 43 species described since this paper was published. We placed these scores in Table 7, incorporated them into those for the entire Central American endemic herpetofauna in Table 8, and provide a graph of the data in Fig. 4.

To illustrate the pattern of distribution of the EVS, we organized these scores by family in Table 9. The data in this table indicate that the scores range from 10 to 20, out of a total theoretical range of 3 to 20. Thus, the scores occupy the entire range of medium vulnerability (10–13) and high vulnerability (14–20) in the EVS scale. None of the scores for these endemic species extend into the low vulnerability range (3–9).

The highest score of 20 is found only among the anurans and, in particular, within the family Hylidae. This score is shared by six hylid species, including four species of *Enomiophyla*, one of *Bromeliophyla*, and one of *Ptychohyla* (Table 8). The lowest score of 10 is seen in a broader range of herpetofaunal families (Table 9), including the Hylidae (one species), Ranidae (one), Dactyloidae (one), Phrynosomatidae (one), and Leptotyphlopidae (one). The greatest number of species, i.e., 146, were assessed an EVS of 16, with species numbers decreasing more or less gradually on either side of this apex to both extremes, i.e., 10 and 20.

Of the 623 total scores, 63 (10.1%) lie within the medium range and the remaining 560 (89.9%) in the high range (Table 9). This large representation of high vulnerability species among the endemic species is of tremendous conservation significance, and figures prominently in the system of prioritization we present below.

Priority Listing for Central American Endemic Herpetofaunal Species

In prioritizing the conservation significance of the endemic herpetofaunal species in Central America, we used the same simple system developed by Johnson et al. (2017). This system involves combining the data on physiographic distribution (Table 4) and the Environmental Vulnerability Scores (Table 8) for the 623 endemic species. This procedure resulted in the recognition of 14 priority levels, of which six are high vulnerability and eight are medium vulnerability groupings (Table 10).

We organized the high vulnerability species into six groups based on the number of physiographic regions they occupy, ranging from one to six (Table 10, Fig. 5). The numbers of species in these seven groups decrease markedly and consistently, as follows: Priority Level One (429 species); Priority Level Two (73); Priority Level Three (27); Priority Level Four (21); Priority Level Five (nine); and Priority Level Six (three). The most significant conclusion of this study is that 562 (90.2%) of the endemic species in Central America are allocated to the six high vulnerability groups. This proportion is 10 percentage points higher than the comparable figure (80.2%) for the Mexican endemic species (Johnson et al., 2017). Furthermore, we believe that the difficulty of protecting these high vulnerability species increases with the fewer physiographic regions they occupy. Thus, the most critically vulnerable species are in the Priority One grouping, the 429 species that constitute 68.9% of the total number of Central American endemics. The challenge of protecting the high vulnerability species increases commensurately with the decrease in the priority level number.

We arranged the medium vulnerability species into eight groups, also on the basis of the number of physiographic regions inhabited (Table 10, Fig. 5). Fewer species are included in these eight groups compared to the high vulnerability ones, as follows: Priority Level Seven (23); Priority Level Eight (21 species); Priority Level Nine (5); Priority Level Ten (four); Priority Level Eleven (four);

Table 7. Environmental Vulnerability Scores (EVS) for 43 endemic members of the Central American herpetofauna not included in Johnson et al. (2015) or requiring recalculation. Question marks indicate decisions made about reproductive mode based on phylogenetic relationships.

Taxa	Environmental Vulnerability Score (EVS)			
	Geographic Distribution	Ecological Distribution	Reproductive Mode/ Degree of Persecution	Total Score
<i>Incilius majordomus</i>	6	8	1?	15
<i>Hyalinobatrachium diana</i>	5	7	3	15
<i>Craugastor gabbi</i>	5	8	4?	17
<i>Diasporus dariensis</i>	5	8	4?	17
<i>Diasporus majeensis</i>	6	8	4?	18
<i>Diasporus pequeno</i>	6	8	4?	18
<i>Diasporus sapo</i>	6	8	4?	18
<i>Plectrohyla calvata</i>	5	8	1	14
<i>Smilisca manisorum</i>	5	8	1?	14
<i>Lithobates lenca</i>	5	8	1	14
<i>Bolitoglossa aurae</i>	6	8	4?	18
<i>Bolitoglossa chucutaniensis</i>	6	8	4?	18
<i>Cryptotriton xucaneborum</i>	6	8	4?	18
<i>Nototriton costaricense</i>	6	8	4?	18
<i>Nototriton nelson</i>	6	7	4?	17
<i>Nototriton oreadorum</i>	6	8	4?	18
<i>Oedipina berlina</i>	5	8	4?	17
<i>Oedipina capitalina</i>	6	8	4?	18
<i>Oedipina salvadorensis</i>	5	8	4?	17
<i>Celestus laf</i>	6	8	3	17
<i>Mesaspis cuchumatana</i>	5	7	3	15
<i>Mesaspis salvadorensis</i>	5	7	3	15
<i>Dactyloa brooksi</i>	5	7	3	15
<i>Dactyloa kathydayae</i>	6	8	3	17
<i>Dactyloa maia</i>	5	7	3	15
<i>Dactyloa savage</i>	5	7	3	15
<i>Norops elcopeensis</i>	5	7	3	15
<i>Norops mccraniei</i>	5	2	3	10
<i>Norops oxylophus</i>	5	6	3	14
<i>Norops wilsoni</i>	5	7	3	15
<i>Lepidoblepharis emberawoundule</i>	5	7	3	15
<i>Lepidoblepharis rufigularis</i>	6	8	3	17
<i>Lepidoblepharis victormartinezi</i>	5	7	3	15
<i>Ameiva fuliginosa</i>	5	8	3	16
<i>Holcosus miadis</i>	6	8	3	17
<i>Tantilla berguido</i>	6	8	2	16
<i>Tantilla excelsa</i>	5	6	2	13
<i>Tantilla gottei</i>	5	7	2	14
<i>Tantilla stenigrammi</i>	5	8	2	15
<i>Rhadinella lisyae</i>	6	7	2	15
<i>Epictia martinezi</i>	6	8	1	15
<i>Epictia pauldwyeri</i>	5	8	1	14
<i>Bothriechis nubestris</i>	5	7	5	17



Plate 16. *Agalychnis annae* (Duellman, 1963). The Orange-eyed Treefrog is a priority two species with an EVS of 15, with a distribution in the “Northern Cordillera de Talamanca, Cordillera de Tilarán and Cordillera Central of Costa Rica” (Frost 2018). This individual was found in Heredia, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 17. *Agalychnis saltator* Taylor, 1955. This leaf frog is a priority three species with an EVS of 14, which ranges along the “Caribbean lowlands of northeastern Honduras, Nicaragua, to east-central Costa Rica” (Frost 2018). This individual was located in Centro Soltis, San Isidro de Peñas Blancas, in the province of Alajuela, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 18. *Lithobates taylori* (Smith, 1959). The Peralta Frog is a priority eight species with an EVS of 12, distributed “at scattered localities on the humid Atlantic lowlands from eastern Nicaragua to southeastern Costa Rica and in the humid premontane and lower montane areas of upland Costa Rica, including the Meseta Oriental and Meseta Occidental and probably the Cordillera Central” (Savage 2002: 402). This individual was found in a pond at Llano Tugrú, in the Serranía de Tabasará. *Photo by Abel Batista.*

Table 8. Environmental Vulnerability Scores (EVS) for the endemic members of the herpetofauna of Central America.

Taxa	EVS	Taxa	EVS
<i>Atelopus certus</i>	14	<i>Oedipina ignea</i>	15
<i>Atelopus chiriquiensis</i>	14	<i>Oedipina kasios</i>	16
<i>Atelopus chirripoensis</i>	15	<i>Oedipina koehleri</i>	16
<i>Atelopus limosus</i>	14	<i>Oedipina leptopoda</i>	17
<i>Atelopus senex</i>	13	<i>Oedipina maritima</i>	18
<i>Atelopus varius</i>	11	<i>Oedipina motaguae</i>	18
<i>Atelopus zeteki</i>	13	<i>Oedipina nica</i>	17
<i>Incilius aucoinae</i>	14	<i>Oedipina nimaso</i>	18
<i>Incilius chompipe</i>	13	<i>Oedipina pacificensis</i>	16
<i>Incilius epioticus</i>	16	<i>Oedipina paucidentata</i>	18
<i>Incilius fastidiosus</i>	13	<i>Oedipina petiola</i>	18
<i>Incilius guanacaste</i>	17	<i>Oedipina poelzi</i>	16
<i>Incilius holdridgei</i>	14	<i>Oedipina pseudouniformis</i>	16
<i>Incilius ibarra</i>	13	<i>Oedipina quadra</i>	17
<i>Incilius karenlipsae</i>	15	<i>Oedipina salvadorensis</i>	17
<i>Incilius leucomyos</i>	12	<i>Oedipina savagei</i>	18
<i>Incilius majordomus</i>	15	<i>Oedipina stenopodia</i>	17
<i>Incilius melanochlorus</i>	12	<i>Oedipina stuarti</i>	15
<i>Incilius periglenes</i>	15	<i>Oedipina taylori</i>	14
<i>Incilius peripatetes</i>	14	<i>Oedipina tomasi</i>	18
<i>Incilius porter</i>	14	<i>Oedipina tzutujilorum</i>	18
<i>Incilius signifier</i>	14	<i>Oedipina uniformis</i>	15
<i>Rhinella centralis</i>	14	<i>Pseudoerycea exspectata</i>	18
<i>Rhinella chrysophora</i>	13	<i>Caecilia volceni</i>	17
<i>Cochranella granulosa</i>	15	<i>Oscaecilia elongata</i>	19
<i>Hyalinobatrachium diana</i>	15	<i>Oscaecilia osae</i>	19
<i>Hyalinobatrachium talamancae</i>	16	<i>Dermophis costaricensis</i>	18
<i>Hyalinobatrachium vireovittatum</i>	16	<i>Dermophis gracilior</i>	18
<i>Craugastor adamastus</i>	18	<i>Dermophis occidentalis</i>	17
<i>Craugastor anciano</i>	16	<i>Gymnopsis multiplicata</i>	14
<i>Craugastor andi</i>	17	<i>Abronia anzueto</i>	18
<i>Craugastor angelicus</i>	15	<i>Abronia aurita</i>	16
<i>Craugastor aphanus</i>	17	<i>Abronia campbelli</i>	18
<i>Craugastor aurilegulus</i>	15	<i>Abronia fimbriata</i>	16
<i>Craugastor azueroensis</i>	16	<i>Abronia frosti</i>	18
<i>Craugastor bocourti</i>	16	<i>Abronia gaiophantasma</i>	16
<i>Craugastor bransfordii</i>	13	<i>Abronia meledona</i>	18
<i>Craugastor campbelli</i>	16	<i>Abronia montecristoi</i>	17
<i>Craugastor catalinae</i>	17	<i>Abronia salvadorensis</i>	17
<i>Craugastor chac</i>	16	<i>Abronia vasconcelosii</i>	16
<i>Craugastor charadra</i>	15	<i>Celestus adercus</i>	17
<i>Craugastor chingopetaca</i>	18	<i>Celestus atitlanensis</i>	15
<i>Craugastor chrysozetetes</i>	18	<i>Celestus bivittatus</i>	15
<i>Craugastor coffeus</i>	18	<i>Celestus cyanochloris</i>	14
<i>Craugastor cruzi</i>	18	<i>Celestus hylaius</i>	16

The endemic herpetofauna of Central America

Table 8 (continued). Environmental Vulnerability Scores (EVS) for the endemic members of the herpetofauna of Central America.

Taxa	EVS	Taxa	EVS
<i>Craugastor cuaquero</i>	18	<i>Celestus laf</i>	17
<i>Craugastor cyanochthebius</i>	18	<i>Celestus montanus</i>	15
<i>Craugastor daryi</i>	17	<i>Celestus orobius</i>	16
<i>Craugastor emcelae</i>	17	<i>Celestus scansorius</i>	15
<i>Craugastor emleni</i>	15	<i>Coloptychon rhombifer</i>	16
<i>Craugastor epochthidius</i>	16	<i>Diploglossus bilobatus</i>	16
<i>Craugastor escoces</i>	15	<i>Diploglossus montisilvestris</i>	18
<i>Craugastor evanesce</i>	17	<i>Mesaspis cuchumatanus</i>	15
<i>Craugastor fecundus</i>	16	<i>Mesaspis monticola</i>	14
<i>Craugastor fleischmanni</i>	16	<i>Mesaspis salvadorensis</i>	15
<i>Craugastor gabbi</i>	17	<i>Basiliscus plumifrons</i>	15
<i>Craugastor gollmeri</i>	16	<i>Dactyloa brooksi</i>	15
<i>Craugastor gulosus</i>	17	<i>Dactyloa casildae</i>	16
<i>Craugastor inachus</i>	17	<i>Dactyloa ginaelisae</i>	12
<i>Craugastor jota</i>	18	<i>Dactyloa ibanezi</i>	15
<i>Craugastor laevisimus</i>	12	<i>Dactyloa insignis</i>	14
<i>Craugastor lauraster</i>	16	<i>Dactyloa kathydayae</i>	17
<i>Craugastor megacephalus</i>	16	<i>Dactyloa kunayalae</i>	15
<i>Craugastor melanostictus</i>	16	<i>Dactyloa maia</i>	15
<i>Craugastor merendonensis</i>	18	<i>Dactyloa microtus</i>	15
<i>Craugastor milesi</i>	16	<i>Dactyloa savagei</i>	15
<i>Craugastor mimus</i>	16	<i>Norops alocomyos</i>	16
<i>Craugastor monnichorum</i>	16	<i>Norops altae</i>	15
<i>Craugastor myllomyllon</i>	18	<i>Norops amplisquamosus</i>	17
<i>Craugastor nefrens</i>	18	<i>Norops apletophallus</i>	15
<i>Craugastor noblei</i>	16	<i>Norops aquaticus</i>	15
<i>Craugastor obesus</i>	17	<i>Norops benedikti</i>	16
<i>Craugastor olanchano</i>	18	<i>Norops bicaorum</i>	17
<i>Craugastor omoaensis</i>	18	<i>Norops campbelli</i>	17
<i>Craugastor pechorum</i>	16	<i>Norops carpenteri</i>	16
<i>Craugastor persimilis</i>	16	<i>Norops charlesmyersi</i>	16
<i>Craugastor phasma</i>	18	<i>Norops cobanensis</i>	13
<i>Craugastor podiciferus</i>	15	<i>Norops cryptolimifrons</i>	16
<i>Craugastor polyptychus</i>	17	<i>Norops cupreus</i>	13
<i>Craugastor punctariolus</i>	16	<i>Norops cusuco</i>	17
<i>Craugastor ranoides</i>	15	<i>Norops datzorum</i>	15
<i>Craugastor rayo</i>	16	<i>Norops elcopeensis</i>	15
<i>Craugastor rhyacobatrachus</i>	16	<i>Norops fortunensis</i>	17
<i>Craugastor rivulus</i>	17	<i>Norops fungosus</i>	15
<i>Craugastor rostralis</i>	16	<i>Norops gruuo</i>	17
<i>Craugastor rugosus</i>	16	<i>Norops haguei</i>	17
<i>Craugastor sabrinus</i>	16	<i>Norops heteropholidotus</i>	16
<i>Craugastor saltuarius</i>	18	<i>Norops humilis</i>	14
<i>Craugastor sandersoni</i>	18	<i>Norops intermedius</i>	14
<i>Craugastor stadelmani</i>	16	<i>Norops johnmeyeri</i>	16

Table 8 (continued). Environmental Vulnerability Scores (EVS) for the endemic members of the herpetofauna of Central America.

Taxa	EVS	Taxa	EVS
<i>Craugastor stejnegerianus</i>	14	<i>Norops kemptoni</i>	15
<i>Craugastor tabasarae</i>	17	<i>Norops kreutzi</i>	17
<i>Craugastor talamancae</i>	17	<i>Norops leditzigorum</i>	15
<i>Craugastor taurus</i>	17	<i>Norops limifrons</i>	15
<i>Craugastor trachydermus</i>	18	<i>Norops lionotus</i>	14
<i>Craugastor underwoodi</i>	16	<i>Norops loveridgei</i>	14
<i>Craugastor xucanebi</i>	16	<i>Norops macrophallus</i>	15
<i>Pristimantis adnus</i>	18	<i>Norops magnaphallus</i>	17
<i>Pristimantis altae</i>	16	<i>Norops marsupialis</i>	16
<i>Pristimantis caryophyllaceus</i>	15	<i>Norops mccraniei</i>	10
<i>Pristimantis cerasinus</i>	16	<i>Norops monteverde</i>	17
<i>Pristimantis museosus</i>	17	<i>Norops morazani</i>	17
<i>Pristimantis pardalis</i>	17	<i>Norops muralla</i>	17
<i>Pristimantis pirrensis</i>	18	<i>Norops ocelloscapularis</i>	15
<i>Strabomantis laticorpus</i>	17	<i>Norops osa</i>	16
<i>Ameerega maculate</i>	18	<i>Norops oxylophus</i>	14
<i>Andinobates claudiae</i>	18	<i>Norops pachypus</i>	15
<i>Andinobates geminisae</i>	18	<i>Norops pijolensis</i>	16
<i>Colostethus latinasus</i>	15	<i>Norops polylepis</i>	15
<i>Ectopoglossus astralogaster</i>	18	<i>Norops pseudokemptoni</i>	17
<i>Ectopoglossus isthminus</i>	16	<i>Norops pseudopachypus</i>	17
<i>Oophaga arborea</i>	16	<i>Norops purpurgularis</i>	16
<i>Oophaga granulifera</i>	17	<i>Norops quaggulus</i>	15
<i>Oophaga pumilio</i>	16	<i>Norops roatanensis</i>	17
<i>Oophaga speciosa</i>	16	<i>Norops rubribarbaris</i>	17
<i>Oophaga vicentei</i>	16	<i>Norops salvini</i>	15
<i>Phyllobates lugubris</i>	17	<i>Norops sminthus</i>	15
<i>Phyllobates vittatus</i>	17	<i>Norops tenorioensis</i>	17
<i>Silverstoneia flotator</i>	16	<i>Norops townsendi</i>	17
<i>Diasporus citrinobapheus</i>	17	<i>Norops triumphalis</i>	17
<i>Diasporus darienensis</i>	17	<i>Norops tropidolepis</i>	15
<i>Diasporus diastema</i>	15	<i>Norops utilensis</i>	17
<i>Diasporus hylaeformis</i>	17	<i>Norops villai</i>	17
<i>Diasporus igneus</i>	18	<i>Norops wampuensis</i>	17
<i>Diasporus majeensis</i>	18	<i>Norops wellbornae</i>	15
<i>Diasporus pequeno</i>	18	<i>Norops wermuthi</i>	16
<i>Diasporus sapo</i>	18	<i>Norops wilsoni</i>	15
<i>Diasporus tigrillo</i>	18	<i>Norops woodi</i>	14
<i>Diasporus ventrimaculatus</i>	18	<i>Norops yoroensis</i>	15
<i>Atlantihyla panchoi</i>	13	<i>Norops zeus</i>	15
<i>Atlantihyla spinipollex</i>	12	<i>Coleonyx mitratus</i>	14
<i>Bromelioshyla melacaena</i>	20	<i>Bachia blairi</i>	15
<i>Dryophytes bocourti</i>	14	<i>Echinosaura panamensis</i>	14
<i>Duellmanohyla legleri</i>	14	<i>Echinosaura apodema</i>	15
<i>Duellmanohyla lythroides</i>	14	<i>Heloderma charlesbogerti</i>	18
<i>Duellmanohyla rufioculis</i>	14	<i>Ctenosaura bakeri</i>	19

The endemic herpetofauna of Central America

Table 8 (continued). Environmental Vulnerability Scores (EVS) for the endemic members of the herpetofauna of Central America.

Taxa	EVS	Taxa	EVS
<i>Duellmanohyla salvadorensis</i>	12	<i>Ctenosaura flavidorsalis</i>	18
<i>Duellmanohyla salvavida</i>	13	<i>Ctenosaura melanosterna</i>	18
<i>Duellmanohyla soralia</i>	12	<i>Ctenosaura oedirhina</i>	19
<i>Duellmanohyla uranochroa</i>	12	<i>Ctenosaura palearis</i>	19
<i>Ecnomiohyla bailarina</i>	20	<i>Ctenosura praeocularis</i>	18
<i>Ecnomiohyla fimbrimembra</i>	19	<i>Ctenosaura quinquecarinata</i>	19
<i>Ecnomiohyla minera</i>	18	<i>Marisora alliacea</i>	15
<i>Ecnomiohyla rabborum</i>	20	<i>Marisora magnacornae</i>	17
<i>Ecnomiohyla salvaje</i>	19	<i>Marisora roatanae</i>	16
<i>Ecnomiohyla sukia</i>	18	<i>Marisora unimarginata</i>	15
<i>Ecnomiohyla thysanota</i>	20	<i>Sceloporus lunaei</i>	15
<i>Ecnomiohyla veraguensis</i>	20	<i>Sceloporus malachiticus</i>	10
<i>Exerodonta catracha</i>	14	<i>Phyllodactylus insularis</i>	17
<i>Exerodonta perkinsi</i>	15	<i>Phyllodactylus palmeus</i>	16
<i>Hyloscirtus colymba</i>	13	<i>Phyllodactylus paralepis</i>	17
<i>Isthmohyla angustilineata</i>	13	<i>Mesoscincus managuae</i>	14
<i>Isthmohyla calypso</i>	16	<i>Lepidoblepharis emberawoundule</i>	15
<i>Isthmohyla debilis</i>	14	<i>Lepidoblepharis rufingularis</i>	17
<i>Isthmohyla graceae</i>	13	<i>Lepidoblepharis victormartinezi</i>	15
<i>Isthmohyla infucata</i>	14	<i>Sphaerodactylus alphas</i>	17
<i>Isthmohyla insolita</i>	17	<i>Sphaerodactylus dunni</i>	15
<i>Isthmohyla lancasteri</i>	14	<i>Sphaerodactylus graptolaemus</i>	16
<i>Isthmohyla picadoi</i>	19	<i>Sphaerodactylus guanaje</i>	17
<i>Isthmohyla pictipes</i>	14	<i>Sphaerodactylus homolepis</i>	16
<i>Isthmohyla pseudopuma</i>	13	<i>Sphaerodactylus leonardovaldesi</i>	16
<i>Isthmohyla rivularis</i>	13	<i>Sphaerodactylus millepunctatus</i>	15
<i>Isthmohyla tica</i>	13	<i>Sphaerodactylus pacificus</i>	17
<i>Isthmohyla xanthosticta</i>	15	<i>Sphaerodactylus poindexteri</i>	17
<i>Isthmohyla zeteki</i>	18	<i>Sphaerodactylus rosaurae</i>	16
<i>Plectrohyla calvata</i>	14	<i>Scincella rara</i>	17
<i>Plectrohyla chrysopleura</i>	13	<i>Cnemidophorus duellmani</i>	16
<i>Plectrohyla dasypus</i>	14	<i>Cnemidophorus ruatanus</i>	15
<i>Plectrohyla exquisita</i>	15	<i>Holcosus leptophrys</i>	16
<i>Plectrohyla glandulosa</i>	12	<i>Holcosus miadis</i>	17
<i>Plectrohyla pokomchi</i>	13	<i>Holcosus quadrilineatus</i>	16
<i>Plectrohyla psiloderma</i>	14	<i>Lepidophyma mayae</i>	13
<i>Plectrohyla quecchi</i>	13	<i>Lepidophyma reticulatum</i>	13
<i>Plectrohyla tecunumani</i>	14	<i>Helminthophis frontalis</i>	12
<i>Plectrohyla teuchestes</i>	15	<i>Dendrophidion apharocybe</i>	16
<i>Ptychohyla dendrophasma</i>	20	<i>Dendrophidion crybelum</i>	17
<i>Ptychohyla hypomykter</i>	10	<i>Dendrophidion paucicarinatum</i>	16
<i>Quilticohyla sanctaecrucis</i>	14	<i>Dendrophidion rufiterminorum</i>	16
<i>Scinax altae</i>	14	<i>Drymobius melanotropis</i>	16
<i>Smilisca manisorum</i>	14	<i>Leptodrymus pulcherrimus</i>	13
<i>Smilisca puma</i>	14	<i>Leptophis nebulosus</i>	14
<i>Leptodactylus silvanimbus</i>	14	<i>Mastigodryas alternatus</i>	12

Table 8 (continued). Environmental Vulnerability Scores (EVS) for the endemic members of the herpetofauna of Central America.

Taxa	EVS	Taxa	EVS
<i>Hypopachus pictiventris</i>	14	<i>Mastigodryas dorsalis</i>	14
<i>Agalychnis annae</i>	15	<i>Oxybelis wilsoni</i>	17
<i>Agalychnis saltatory</i>	14	<i>Scolecophis atrocinctus</i>	13
<i>Pipa myersi</i>	17	<i>Tantilla albiceps</i>	16
<i>Lithobates juliani</i>	12	<i>Tantilla armillata</i>	11
<i>Lithobates lenca</i>	14	<i>Tantilla bairdi</i>	16
<i>Lithobates miadis</i>	15	<i>Tantilla berguidoi</i>	16
<i>Lithobates taylori</i>	12	<i>Tantilla brevicauda</i>	13
<i>Lithobates vibicarius</i>	14	<i>Tantilla excelsa</i>	13
<i>Lithobates warszewitschii</i>	10	<i>Tantilla gottei</i>	14
<i>Bolitoglossa alvaradoi</i>	16	<i>Tantilla hendersoni</i>	16
<i>Bolitoglossa anthracina</i>	18	<i>Tantilla jani</i>	14
<i>Bolitoglossa aurae</i>	18	<i>Tantilla lempira</i>	14
<i>Bolitoglossa aureogularis</i>	18	<i>Tantilla olympia</i>	16
<i>Bolitoglossa bramei</i>	17	<i>Tantilla psittaca</i>	15
<i>Bolitoglossa carri</i>	18	<i>Tantilla ruficeps</i>	12
<i>Bolitoglossa cataguana</i>	18	<i>Tantilla stenigrammi</i>	15
<i>Bolitoglossa celaque</i>	17	<i>Tantilla taeniata</i>	14
<i>Bolitoglossa centenorum</i>	18	<i>Tantilla tecta</i>	16
<i>Bolitoglossa cerroensis</i>	16	<i>Tantilla tritaeniata</i>	16
<i>Bolitoglossa chucantiensis</i>	18	<i>Tantilla vermiformis</i>	14
<i>Bolitoglossa colonnea</i>	16	<i>Trimorphodon quadruplex</i>	14
<i>Bolitoglossa compacta</i>	17	<i>Adelphicos daryi</i>	16
<i>Bolitoglossa conanti</i>	16	<i>Adelphicos ibarrorum</i>	15
<i>Bolitoglossa copia</i>	18	<i>Adelphicos veraepacis</i>	14
<i>Bolitoglossa cuchumatana</i>	14	<i>Atractus darienensis</i>	16
<i>Bolitoglossa cuna</i>	17	<i>Atractus depressiocellus</i>	15
<i>Bolitoglossa daryorum</i>	17	<i>Atractus hostilitractus</i>	16
<i>Bolitoglossa decora</i>	18	<i>Atractus imperfectus</i>	16
<i>Bolitoglossa diaphora</i>	18	<i>Chapinophis xanthocheilus</i>	16
<i>Bolitoglossa diminuta</i>	18	<i>Coniophanes joanae</i>	15
<i>Bolitoglossa dofleini</i>	15	<i>Crisantophis nevermanni</i>	16
<i>Bolitoglossa dunni</i>	16	<i>Cubophis brooksi</i>	14
<i>Bolitoglossa epimela</i>	17	<i>Dipsas articulata</i>	15
<i>Bolitoglossa eremia</i>	18	<i>Dipsas bicolor</i>	17
<i>Bolitoglossa gomezi</i>	16	<i>Dipsas nicholsi</i>	15
<i>Bolitoglossa gracilis</i>	18	<i>Dipsas tenuissima</i>	14
<i>Bolitoglossa heiroreias</i>	17	<i>Enulius bifoveatus</i>	16
<i>Bolitoglossa helmrichi</i>	16	<i>Enulius roatanensis</i>	16
<i>Bolitoglossa huehuetenanguensis</i>	18	<i>Geophis bellus</i>	16
<i>Bolitoglossa indio</i>	17	<i>Geophis brachycephalus</i>	11
<i>Bolitoglossa insularis</i>	18	<i>Geophis championi</i>	16
<i>Bolitoglossa jacksoni</i>	18	<i>Geophis damiani</i>	16
<i>Bolitoglossa jugivagans</i>	18	<i>Geophis downsi</i>	16
<i>Bolitoglossa kamuk</i>	18	<i>Geophis dunni</i>	16

The endemic herpetofauna of Central America

Table 8 (continued). Environmental Vulnerability Scores (EVS) for the endemic members of the herpetofauna of Central America.

Taxa	EVS	Taxa	EVS
<i>Bolitoglossa kaqchikelorum</i>	17	<i>Geophis fulvoguttatus</i>	14
<i>Bolitoglossa la</i>	17	<i>Geophis godmani</i>	14
<i>Bolitoglossa lignicolor</i>	16	<i>Geophis hoffmanni</i>	12
<i>Bolitoglossa longissima</i>	18	<i>Geophis nephodrymus</i>	16
<i>Bolitoglossa magnifica</i>	16	<i>Geophis ruthveni</i>	14
<i>Bolitoglossa marmorea</i>	17	<i>Geophis talamancae</i>	15
<i>Bolitoglossa meliana</i>	16	<i>Geophis tectus</i>	13
<i>Bolitoglossa minutula</i>	17	<i>Geophis zeledoni</i>	15
<i>Bolitoglossa mombachoensis</i>	17	<i>Hydromorphus concolor</i>	12
<i>Bolitoglossa morio</i>	13	<i>Hydromorphus dunnii</i>	16
<i>Bolitoglossa nigrescens</i>	16	<i>Imantodes phantasma</i>	16
<i>Bolitoglossa ninadormida</i>	18	<i>Leptodeira rhombifera</i>	12
<i>Bolitoglossa nussbaumi</i>	18	<i>Leptodeira rubricata</i>	17
<i>Bolitoglossa nympa</i>	16	<i>Ninia celata</i>	15
<i>Bolitoglossa obscura</i>	18	<i>Ninia espinali</i>	14
<i>Bolitoglossa odonnelli</i>	16	<i>Ninia maculata</i>	12
<i>Bolitoglossa omniumsanctorum</i>	16	<i>Ninia pavimentata</i>	15
<i>Bolitoglossa oresbia</i>	17	<i>Ninia psephota</i>	13
<i>Bolitoglossa pacaya</i>	17	<i>Omoadiphas aurula</i>	16
<i>Bolitoglossa pesrubra</i>	15	<i>Omoadiphas cannula</i>	16
<i>Bolitoglossa porrasorum</i>	16	<i>Omoadiphas texiguatensis</i>	16
<i>Bolitoglossa psephena</i>	18	<i>Rhadinaea calligaster</i>	14
<i>Bolitoglossa pygmaea</i>	17	<i>Rhadinaea pulveriventris</i>	14
<i>Bolitoglossa robinsoni</i>	16	<i>Rhadinaea sargenti</i>	14
<i>Bolitoglossa robusta</i>	16	<i>Rhadinaea stadelmani</i>	13
<i>Bolitoglossa salvinii</i>	16	<i>Rhadinaea vermiculaticeps</i>	15
<i>Bolitoglossa schizodactyla</i>	15	<i>Rhadinella anachoreta</i>	14
<i>Bolitoglossa sombra</i>	16	<i>Rhadinella hempsteadae</i>	13
<i>Bolitoglossa sooyorum</i>	16	<i>Rhadinella lisyae</i>	15
<i>Bolitoglossa splendida</i>	18	<i>Rhadinella montecristi</i>	14
<i>Bolitoglossa striatula</i>	16	<i>Rhadinella pegosalyla</i>	16
<i>Bolitoglossa subpalmata</i>	15	<i>Rhadinella pilonaorum</i>	15
<i>Bolitoglossa suchitanensis</i>	18	<i>Rhadinella rogerromani</i>	16
<i>Bolitoglossa synoria</i>	17	<i>Rhadinella serperaster</i>	13
<i>Bolitoglossa taylori</i>	17	<i>Rhadinella tolpanorum</i>	16
<i>Bolitoglossa tenebrosa</i>	17	<i>Sibon anthracops</i>	15
<i>Bolitoglossa tica</i>	17	<i>Sibon argus</i>	16
<i>Bolitoglossa tzultacaj</i>	18	<i>Sibon carri</i>	14
<i>Bolitoglossa xibalba</i>	17	<i>Sibon lamari</i>	16
<i>Bolitoglossa zacapensis</i>	18	<i>Sibon longifrenis</i>	14
<i>Cryptotriton monzoni</i>	18	<i>Sibon manzanaresi</i>	15
<i>Cryptotriton nasalis</i>	18	<i>Sibon merendonensis</i>	16
<i>Cryptotriton necopinus</i>	18	<i>Sibon miskitus</i>	15
<i>Cryptotriton sierraminensis</i>	17	<i>Sibon noalamina</i>	15
<i>Cryptotriton veraepacis</i>	17	<i>Sibon perissostichon</i>	16

Table 8 (continued). Environmental Vulnerability Scores (EVS) for the endemic members of the herpetofauna of Central America.

Taxa	EVS	Taxa	EVS
<i>Cryptotriton xucaneborum</i>	18	<i>Trimetopon barbouri</i>	15
<i>Dendrotriton bromeliacus</i>	17	<i>Trimetopon gracile</i>	14
<i>Dendrotriton chujorum</i>	18	<i>Trimetopon pliolepis</i>	12
<i>Dendrotriton cuchumatanus</i>	18	<i>Trimetopon simile</i>	13
<i>Dendrotriton kekchiorum</i>	18	<i>Trimetopon slevini</i>	14
<i>Dendrotriton rabbi</i>	17	<i>Trimetopon viquezi</i>	15
<i>Dendrotriton sanctibarbarus</i>	18	<i>Urotheca guentheri</i>	12
<i>Nototriton abscondens</i>	16	<i>Urotheca myersi</i>	15
<i>Nototriton barbouri</i>	16	<i>Urotheca pachyura</i>	14
<i>Nototriton brodiei</i>	17	<i>Micrurus alleni</i>	16
<i>Nototriton costaricense</i>	18	<i>Micrurus hippocrepis</i>	18
<i>Nototriton gamezi</i>	18	<i>Micrurus mosquitensis</i>	17
<i>Nototriton guanacaste</i>	17	<i>Micrurus ruatanus</i>	18
<i>Nototriton lignicola</i>	18	<i>Micrurus stewarti</i>	17
<i>Nototriton limnospectator</i>	17	<i>Micrurus stuarti</i>	17
<i>Nototriton major</i>	18	<i>Epictia ater</i>	10
<i>Nototriton matama</i>	18	<i>Epictia martinezi</i>	15
<i>Nototriton mime</i>	18	<i>Epictia pauldwyeri</i>	14
<i>Nototriton nelsoni</i>	17	<i>Amerotyphlops costaricensis</i>	11
<i>Nototriton oreadorum</i>	18	<i>Amerotyphlops stadelmani</i>	12
<i>Nototriton picadoi</i>	16	<i>Typhlops tycherus</i>	14
<i>Nototriton picucha</i>	18	<i>Agkistrodon howardgloydi</i>	17
<i>Nototriton richardi</i>	16	<i>Atropoides indomitus</i>	18
<i>Nototriton saslaya</i>	18	<i>Atropoides picadoi</i>	16
<i>Nototriton stuarti</i>	18	<i>Bothriechis guifarroi</i>	19
<i>Nototriton tapanti</i>	18	<i>Bothriechis lateralis</i>	16
<i>Nototriton tomamorum</i>	18	<i>Bothriechis marchi</i>	16
<i>Oedipina alfaroi</i>	16	<i>Bothriechis nigroviridis</i>	17
<i>Oedipina alleni</i>	16	<i>Bothriechis nubestris</i>	17
<i>Oedipina altura</i>	18	<i>Bothriechis supraciliaris</i>	17
<i>Oedipina berlini</i>	17	<i>Bothriechis thalassinus</i>	17
<i>Oedipina capitalina</i>	18	<i>Cerrophidion sasai</i>	16
<i>Oedipina carablanca</i>	18	<i>Cerrophidion wilsoni</i>	15
<i>Oedipina chortiorum</i>	18	<i>Lachesis melanocephala</i>	17
<i>Oedipina collaris</i>	17	<i>Lachesis stenophrys</i>	17
<i>Oedipina cyclocauda</i>	15	<i>Porthidium porrasi</i>	18
<i>Oedipina fortunensis</i>	18	<i>Porthidium volcanicum</i>	18
<i>Oedipina gephyra</i>	17	<i>Rhinoclemmys funerea</i>	16
<i>Oedipina gracilis</i>	16	<i>Kinosternon angustipons</i>	16
<i>Oedipina grandis</i>	17		

The endemic herpetofauna of Central America




Fig. 4. Graph showing Central American endemic species and their corresponding Environmental Vulnerability Scores (EVS) ranging from 10 to 20.




Fig. 5. Graph of Central American endemic species allocated to the 14 conservation priority groups.

Priority Level Twelve (two); Priority Level Thirteen (one); and Priority Level Fourteen (one). Even so, the next most important conclusion of this study is that these 61 species make up 9.8% of the total compendium of endemic species in Central America. The number of species in these eight groups also decreases sharply, as 69.8% of the 61 species fall into the first two priority levels, i.e., Seven and Eight.

When we examined the 623 endemic species relative to the number of physiographic regions inhabited, the results are as follows: one region ($429+23 = 452$); two regions ($73+21 = 94$); three regions ($27+5 = 32$); four regions ($21+4 = 25$); five regions ($9+4 = 13$); six regions ($3+2 = 5$); seven regions (1); and eight regions (1). Perusal of these data supports another conclusion, i.e., that 72.6% of the total number of species occupy a single physiographic region. Based on the assumptions of this study, these 452 species can be expected to offer the major challenge in efforts to protect the endemic component of the Central American herpetofauna. The next most challenging group contains the 94 species occupying two regions. Together, the single-group and double-group species comprise 546 (87.6%) of the total of 623 Central American endemic species.

Our analysis in this paper indicates that most of the 623 endemic Central American herpetofaunal species are judged as high vulnerability based on the EVS methodology, and are demonstrated to occupy relatively few physiographic regions (one or two). The endemic component of the Central American herpetofauna, just as with the Mexican endemic component (Johnson et al. 2017), is of global significance and constitutes the most significant challenge to conservation professionals working within this segment of the Mesoamerican herpetofauna. Johnson et al. (2017) arrived at the same conclusion in their work on the Mexican endemic herpetofauna. Considered as a whole, the Mesoamerican endemic herpetofauna comprises the 789 Mexican endemic species dealt with by Johnson et al. (2017) and the 623 Central American endemic species dealt with here, as well as the 225 species restricted in distribution to Mexico and Central America (i.e., Mesoamerica; Wilson et al., 2017) for a total of 1,637 species. This figure represents more than three quarters of the entire Mesoamerican herpetofauna (Wilson et al., 2017). We examine the parameters of the challenge facing conservation biologists working in Central America in the following section.

Prognosis for the Endemic Central American Herpetofauna

The same environmental issues impacting the Mexican endemic herpetofauna, as discussed by Johnson et al. (2017), also impinge upon the Central American endemic herpetofauna. In light of this situation, we emphasize that the survival of the 623 endemic species inhabiting Central America ultimately depends on addressing the underlying issues that lead to all environmental problems, including biodiversity decline, that in turn stand in the way of designing a sustainable existence for humanity's tenure on

Earth. Johnson et al. (2017: 609) explained what we face as follows: "Fundamentally, humans have created and maintain these environmental problems because of their capacity for rational thought, i.e., their ability to connect cause to effect through the passing of time, and adopting an anthropocentric worldview that stresses the exploitation of the world's resources to support the burgeoning human population. Such a worldview contrasts markedly with that of environmentalists, who have adopted 'a worldview that helps us make sense of how the environment works, our place in the environment, and right and wrong environmental behaviors' (Raven and Berg 2004: G-6). Obviously, the present anthropocentric worldview held by most people represents the fundamental reason why these environmental problems exist, and continued human population growth allows them to worsen over time."

The anthropocentric worldview, also known as the Western worldview, "includes human superiority and dominance over nature, the unrestricted use of natural resources, increased economic growth to manage an expanding industrial base, the inherent rights of individuals, and accumulation of wealth and unlimited consumption of goods and services to provide material comforts" (Raven and Berg 2004: 17). This worldview not only creates the entire spectrum of environmental problems, but also the entire panoply of human societal issues we see played out every day in various media outlets. Ultimately, they arise from a commitment to discriminate among groups of people, i.e., on the basis of racial background, gender, religion, economic wealth, political persuasion, and so forth. Thus, not only is humanity poised against the rest of the living world, but also varying groups of humans are in conflict with one another. As the focus of humanity decreases from larger to increasingly smaller realms of interest, it can be argued that mental stability gives way to instability, and eventually gives rise to the increased incidence of the narcissistic personality disorder (NPD). This disorder is highly variable in presentation and can manifest across a broad spectrum of severity, but is generally characterized by pervasive grandiosity, an excessive need for admiration, and a lack of empathy (Caligor et al. 2015). Envisioning NPD as an extreme end-point of the intensification of anthropocentrism might explain why the potential causes of this disorder remain unknown and that clinical guidelines have yet to emerge (Caligor et al. 2015). Given that none of the authors of this paper possesses credentials in psychology or psychiatry, our idea about the connection between the anthropocentric worldview and the narcissistic personality disorder can be best understood as a hypothesis remaining to be tested, hopefully by a cross-discipline team of environmental scientists, deep ecology philosophers, and biocentric psychologists/psychiatrists. Studying such a connection could lie within the realm of environmental psychology, defined as an interdisciplinary field that focuses on the interplay between environments and human cognition and behavior; considering the term "environment" broadly, including both natural and human-

The endemic herpetofauna of Central America

Table 9. Summary of EVS values for Central American endemic species, arranged by family. Shaded area encompasses high vulnerability scores.

Families	Number of Species	Environmental Vulnerability Scores										
		10	11	12	13	14	15	16	17	18	19	20
Bufonidae	24	—	1	2	6	9	4	1	1	—	—	—
Centrolenidae	4	—	—	—	—	—	2	2	—	—	—	—
Craugastoridae	77	—	—	1	1	1	8	29	18	19	—	—
Dendrobatidae	14	—	—	—	—	—	1	6	3	4	—	—
Eleutherodactyludae	10	—	—	—	—	—	1	—	3	6	—	—
Hylidae	52	1	—	5	11	17	4	1	1	3	3	6
Leptodactylidae	1	—	—	—	—	1	—	—	—	—	—	—
Microhylidae	1	—	—	—	—	1	—	—	—	—	—	—
Phyllomedusidae	2	—	—	—	—	1	1	—	—	—	—	—
Pipidae	1	—	—	—	—	—	—	—	1	—	—	—
Ranidae	6	1	—	2	—	2	1	—	—	—	—	—
Subtotals	192	2	1	10	18	32	22	39	27	32	3	6
Plethodontidae	143	—	—	—	1	2	8	33	38	61	—	—
Subtotals	143	—	—	—	1	2	8	33	38	61	—	—
Caeciliidae	3	—	—	—	—	—	—	—	1	—	2	—
Dermophiidae	4	—	—	—	—	1	—	—	1	2	—	—
Subtotals	7	—	—	—	—	1	—	—	2	2	2	—
Totals	342	2	1	10	19	35	30	72	67	95	5	6
Anguidae	25	—	—	—	—	2	6	8	4	5	—	—
Corytophanidae	1	—	—	—	—	—	1	—	—	—	—	—
Dactyloidae	74	1	—	1	2	7	27	13	23	—	—	—
Eublepharidae	1	—	—	—	—	1	—	—	—	—	—	—
Gymnophthalmidae	3	—	—	—	—	1	2	—	—	—	—	—
Helodermatidae	1	—	—	—	—	—	—	—	—	1	—	—
Iguanidae	7	—	—	—	—	—	—	—	—	3	4	—
Mabuyidae	4	—	—	—	—	—	2	1	1	—	—	—
Phrynosomatidae	2	1	—	—	—	—	1	—	—	—	—	—
Phyllodactylidae	3	—	—	—	—	—	—	1	2	—	—	—
Scincidae	1	—	—	—	—	1	—	—	—	—	—	—
Sphaerodactylidae	13	—	—	—	—	—	4	4	5	—	—	—
Sphenomorphidae	1	—	—	—	—	—	—	—	1	—	—	—
Teiidae	5	—	—	—	—	—	1	3	1	—	—	—
Xantusiidae	2	—	—	—	2	—	—	—	—	—	—	—
Subtotals	143	2	—	1	4	12	44	30	37	9	4	—
Anomalepididae	1	—	—	1	—	—	—	—	—	—	—	—
Colubridae	30	—	1	2	4	8	2	11	2	—	—	—
Dipsadidae	77	—	1	6	6	17	19	26	2	—	—	—
Elapidae	6	—	—	—	—	—	—	1	3	2	—	—
Leptotyphlopidae	3	1	—	—	—	1	1	—	—	—	—	—
Typhlopidae	3	—	1	1	—	1	—	—	—	—	—	—
Viperidae	16	—	—	—	—	—	1	4	7	3	1	—
Subtotals	136	1	3	10	10	27	23	42	14	5	1	—
Geoemydidae	1	—	—	—	—	—	—	1	—	—	—	—

Table 9 (continued). Summary of EVS values for Central American endemic species, arranged by family. Shaded area encompasses high vulnerability scores.

Families	Number of Species	Environmental Vulnerability Scores										
		10	11	12	13	14	15	16	17	18	19	20
Kinosternidae	1	—	—	—	—	—	—	1	—	—	—	—
Subtotals	2	—	—	—	—	—	—	2	—	—	—	—
Totals	281	3	3	11	14	39	67	74	51	14	5	—
Sum Totals	623	5	4	21	33	74	97	146	118	109	10	6
Category Totals	623	63					560					



Plate 19. *Lithobates warszewitschii* (Schmidt, 1857). Warszewitsch’s Frog is a priority eleven species with an EVS of 10, found on “the Atlantic versant from northeastern Honduras to central Panama, both slopes of the cordilleras of Costa Rica and western Panama, lowlands of southwestern Costa Rica, and eastern Panama and gallery forests in nonpeninsular northwestern Costa Rica” (Savage 2002: 405). This individual came from Nectandra Reserve, in the province of Alajuela, Costa Rica. *Photo by Sean Michael Rovito.*



Plate 20. *Bolitoglossa alvaradoi* Taylor, 1954. The Moravia de Chirripó Salamander is a priority two species with an EVS of 16, distributed on “the Atlantic versant of Costa Rica” (Frost, 2018). This individual was found in Veragua Rainforest, in the province of Limón, Costa Rica. *Photo by Victor Acosta-Chaves.*



Plate 21. *Bolitoglossa centenorum* Campbell, Smith, Streicher, Acevedo, and Brodie, 2010. This salamander is a priority one species with an EVS of 18, which is known “only from the type locality in the Sierra Cuchumatanes, Huehuetenango, Guatemala” (Frost 2018). This individual was encountered at the type locality, near San Mateo Ixtatán. *Photo by Todd Pierson.*

made environments (De Young 2013). Since its conception, research in environmental psychology has often targeted human attitudes towards the natural environment, and current trends are now shifting to a focus on sustainable living in the context of environmental issues (De Young 2013).

Finding lasting solutions to environmental problems must be based on a realistic, fact-based approach that evaluates the symptoms of these problems until their causes are identified (Wilson and McCranie 2004). Often, the search for an ultimate cause stops when exposed to the anthropocentric worldview. A worldview, however, is a collection of basic values that “help us to make sense of the world, understand our place in it, and determine right and wrong behaviors” (Raven and Berg 2004: G-17). How the values that characterize the anthropocentric worldview have arisen through the evolution of human behavior to become predominant, however, generally has not been explored. Our working assumption, i.e., our hypothesis, is that the ultimate causes are deeply engrained in the origins of human behavior and have become so pervasive as to underlie our efforts to understand our world, and our place in it. Even a discipline called environmental psychology might not expose the steps in behavioral evolution that would allow present-day humans to address the malady known as the anthropocentric worldview. In particular, this viewpoint is evident when considering that environmental psychology adopts a broad array of theories, methods, and interpretations from other disciplines as needed, and this mosaic approach can make it difficult to understand the field as a whole and the role it might play in these societal issues moving forward. An encouraging sign is the recent emergence of even more specific sub-fields, such as conservation psychology and ecopsychology, which aim to provide solutions or interventions for problems specifically related to conservation of the natural world (Steg and Vlek 2009; De Young 2013).

There is clearly a critical need to develop novel approaches for studying animal behavior and human psychology that emphasize reasons why the anthropocentric worldview has become so predominant, and what needs to be done to replace it with the environmental worldview. If, as we hypothesized, there is a psychological connection among centrist forms of thinking at larger scales (i.e., the anthropocentric worldview) and those at smaller scales (i.e., narcissism), then we are faced with an even greater challenge than commonly is envisioned.

Searching for Ultimate Solutions

Johnson et al. (2017: 613) offered some ideas about searching for ultimate solutions to the problem of biodiversity decline, based on opinions promulgated by Wilson and Townsend (2010), Wilson (2016), and Kopnina (2016), and concluded as follows: “Our opinion is that humans have the rational capacity to design a sustainable world through cooperative action, but our species’ attitudes and actions will have to

change. Our preparedness will have to improve as well. Such change will have to be based on realistic, fact-based appraisals of where we are now and where we want to be in the future. Biologists will have to commit to helping the rest of us understand why the protection of biodiversity is critical to enjoying a sustainable world. Cultural anthropologists also will have to assist humanity at large to understand why the maintenance of cultural diversity also is essential to living sustainably. Educational reform will have to be central to such efforts, to help people learn how to think and act critically and base decisions on the way things really are, and not how we might wish them to be by denying reality. The devotion humans have for structuring beliefs on the basis of little or no evidence, essentially reversing the benefit of rationality, will have to surrender to critical-thinking education established by top-to-bottom educational reform.”

Critical-thinking educational reform, however, is much easier to conceive than to bring into reality. A fundamental question is why such reform has not been undertaken. This question is not easy to answer, but perhaps the most fundamental reason is that the educational systems currently in existence are products of the anthropocentric worldview and reflect its mindsets. These educational systems also have developed within the current economic systems responsible for the huge disparities between the rich and poor, and act to reinforce these disparities.

Ultimate solutions will emerge only from a clear understanding of the evolution of human psychology, as confronted with the problems we face. If not, then the endemic herpetofauna of Central America, as well as the remainder of life on Earth, will become casualties of the biodiversity crisis that eventually will envelop all humanity.

Conclusions and Recommendations

Conclusions

- A. As concluded by Johnson et al. (2017), life on Earth exists as a result of the interplay among the planet’s three abiotic spheres, the atmosphere, hydrosphere, and lithosphere.
- B. Environmental problems and the biodiversity crisis exist because of the impact of humans on all of the planetary spheres, including the biosphere, and extend along their existing energy and materials pathways.
- C. The biodiversity crisis impacts all life across the globe, and all levels of its organization.
- D. The endemic component of the Central American herpetofauna is of global significance, and its importance increases with the addition of new information. Forty-three species have been added to this component within the last two years, bringing the total to 623 species.
- E. The percentage of endemism of the Central American herpetofauna is 56.9, compared to 61.1 in Mexico, the other major portion of Mesoamerica.



Plate 22. *Bolitoglossa conanti* McCranie and Wilson, 1993. Conant’s Mushroomtongue Salamander is a priority one species with an EVS of 16, with a distribution “on the Atlantic and Pacific versants in western Honduras and eastern Guatemala” (Townsend and Wilson 2008). This individual came from Aldea San Joaquin, in the department of Copán, Honduras. *Photo by Sean Michael Rovito.*



Plate 23. *Bolitoglossa cuchumatana* (Stuart, 1943). The Oak Forest Salamander is a priority one species with an EVS of 14, which is found in the “departments of El Quiché and Huehuetenango in the Sierra de Cuchumatanes, Guatemala” (Frost, 2018). This individual came from near Laguna Maxbal, in the department of Huehuetenango, Guatemala. *Photo by Todd Pierson.*



Plate 24. *Bolitoglossa diaphora* McCranie and Wilson, 1995. The Cusuco Salamander is a priority one species with an EVS of 18, which ranges on “the Atlantic versant in northwestern Honduras” (Townsend and Wilson 2008). This individual was found in Parque Nacional Cusuco, in the department of Cortés, Honduras. *Photo by Sean Michael Rovito.*



Plate 25. *Bolitoglossa subpalmata* (Boulenger, 1896). This salamander is a priority one species with an EVS of 15, which occurs “on both slopes of the Cordillera de Guanacaste, Cordillera de Tilarán, Cordillera Central, and their outliers in central to northern Costa Rica” (Frost 2018). This individual was encountered in Volcán Barva, Parque Nacional Braulio Carrillo, in the province of, Heredia, Costa Rica. *Photo by Víctor Acosta-Chaves.*



Plate 26. *Oedipina koehleri* Sunyer, Townsend, Wake, Travers, Gonzalez, Obando, and Quintana, 2011. This worm salamander is a priority one species with an EVS of 16, restricted to “three highland areas in northern Nicaragua” (Frost 2018). This individual came from Reserva Natural Cerro Musún, in the department of Matagalpa, Nicaragua. *Photo by Javier Sunyer.*



Plate 27. *Coloptychon rhombifer* (Peters, 1876). The Isthmian Alligator Lizard is a priority one species with an EVS of 16, distributed from “southwestern Costa Rica and adjacent western Panama (Savage, 2002: 533). This individual was found in San Juan de Rincón, Península de Osa, in the province of Puntarenas, Costa Rica. *Photo by César Barrio-Amorós.*

The endemic herpetofauna of Central America

Table 10. Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

Priority One: High Vulnerability Species in Single Physiographic Region (429 species)	
<i>Atelopus chiriquiensis</i>	<i>Nototriton tapanti</i>
<i>Atelopus chirripoensis</i>	<i>Nototriton tomamorum</i>
<i>Atelopus limosus</i>	<i>Oedipina altura</i>
<i>Incilius aucoinae</i>	<i>Oedipina berlini</i>
<i>Incilius epioticus</i>	<i>Oedipina capitalina</i>
<i>Incilius guanacaste</i>	<i>Oedipina carablanca</i>
<i>Incilius holdridgei</i>	<i>Oedipina chortiorum</i>
<i>Incilius karenlipsae</i>	<i>Oedipina collaris</i>
<i>Incilius majordomus</i>	<i>Oedipina cyclocauda</i>
<i>Incilius periglenes</i>	<i>Oedipina fortunensis</i>
<i>Incilius peripatetes</i>	<i>Oedipina gephyra</i>
<i>Incilius porteri</i>	<i>Oedipina gracilis</i>
<i>Hyalinobatrachium talamancae</i>	<i>Oedipina grandis</i>
<i>Hyalinobatrachium vireovittatum</i>	<i>Oedipina kasios</i>
<i>Craugastor adamastus</i>	<i>Oedipina koehleri</i>
<i>Craugastor anciano</i>	<i>Oedipina leptopoda</i>
<i>Craugastor andi</i>	<i>Oedipina maritima</i>
<i>Craugastor angelicus</i>	<i>Oedipina motaguae</i>
<i>Craugastor aphanus</i>	<i>Oedipina nica</i>
<i>Craugastor azueroensis</i>	<i>Oedipina nimaso</i>
<i>Craugastor bocourti</i>	<i>Oedipina pacificensis</i>
<i>Craugastor catalinae</i>	<i>Oedipina paucidentata</i>
<i>Craugastor chingopetaca</i>	<i>Oedipina petiola</i>
<i>Craugastor chrysozetetes</i>	<i>Oedipina poelzi</i>
<i>Craugastor coffeus</i>	<i>Oedipina quadra</i>
<i>Craugastor cruzi</i>	<i>Oedipina salvadorensis</i>
<i>Craugastor cuaquero</i>	<i>Oedipina savagei</i>
<i>Craugastor cyanochthebius</i>	<i>Oedipina stenopodia</i>
<i>Craugastor daryi</i>	<i>Oedipina taylori</i>
<i>Craugastor emcelae</i>	<i>Oedipina tomasi</i>
<i>Craugastor emleni</i>	<i>Oedipina tzutujilorum</i>
<i>Craugastor escoces</i>	<i>Oedipina uniformis</i>
<i>Craugastor fleischmanni</i>	<i>Pseudoeurycea expectata</i>
<i>Craugastor gabbi</i>	<i>Caecilia volcani</i>
<i>Craugastor gulosus</i>	<i>Oscaecilia elongata</i>
<i>Craugastor inachus</i>	<i>Oscaecilia osae</i>
<i>Craugastor jota</i>	<i>Dermophis costaricensis</i>
<i>Craugastor melanostictus</i>	<i>Dermophis gracilior</i>
<i>Craugastor merendonensis</i>	<i>Abronia anzuetoii</i>
<i>Craugastor milesi</i>	<i>Abronia aurita</i>
<i>Craugastor monnichorum</i>	<i>Abronia campbelli</i>
<i>Craugastor myllomyllon</i>	<i>Abronia fimbriata</i>
<i>Craugastor nefrens</i>	<i>Abronia frosti</i>
<i>Craugastor olanchano</i>	<i>Abronia gaiophantasma</i>
<i>Craugastor omoaensis</i>	<i>Abronia meledona</i>

Table 10 (continued). Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

<i>Craugastor phasma</i>	<i>Abronia montecristoi</i>
<i>Craugastor podiciferus</i>	<i>Abronia salvadorensis</i>
<i>Craugastor polyptychus</i>	<i>Abronia vasconcelosii</i>
<i>Craugastor punctariolus</i>	<i>Celestus adercus</i>
<i>Craugastor rayo</i>	<i>Celestus bivittatus</i>
<i>Craugastor rhyacobatrachus</i>	<i>Celestus cyanochloris</i>
<i>Craugastor rivulus</i>	<i>Celestus hylaius</i>
<i>Craugastor saltuarius</i>	<i>Celestus laf</i>
<i>Craugastor stadelmani</i>	<i>Celestus montanus</i>
<i>Craugastor tabasarae</i>	<i>Celestus orobius</i>
<i>Craugastor talamancae</i>	<i>Celestus scansorius</i>
<i>Craugastor taurus</i>	<i>Coloptychon rhombifer</i>
<i>Craugastor trachydermus</i>	<i>Diploglossus montisilvestris</i>
<i>Craugastor underwoodi</i>	<i>Mesaspis cuchumatanus</i>
<i>Craugastor xucanebi</i>	<i>Mesaspis monticola</i>
<i>Pristimantis adnus</i>	<i>Mesaspis salvadorensis</i>
<i>Pristimantis museosus</i>	<i>Dactyloa casildae</i>
<i>Pristimantis pirrensis</i>	<i>Dactyloa kathydayae</i>
<i>Strabomantis laticorpus</i>	<i>Dactyloa microtus</i>
<i>Ameerega maculata</i>	<i>Norops alocomyos</i>
<i>Andinobates claudiae</i>	<i>Norops altae</i>
<i>Andinobates geminisae</i>	<i>Norops amplisquamosus</i>
<i>Colostethus latinasus</i>	<i>Norops benedikti</i>
<i>Ectopoglossus astralogaster</i>	<i>Norops bicaorum</i>
<i>Ectopoglossus isthminus</i>	<i>Norops campbelli</i>
<i>Oophaga arborea</i>	<i>Norops cusuco</i>
<i>Oophaga pumilio</i>	<i>Norops datzorum</i>
<i>Oophaga speciosa</i>	<i>Norops fortunensis</i>
<i>Phyllobates lugubris</i>	<i>Norops fungosus</i>
<i>Phyllobates vittatus</i>	<i>Norops gruuo</i>
<i>Diasporus citrinobapheus</i>	<i>Norops haguei</i>
<i>Diasporus darienensis</i>	<i>Norops heteropholidotus</i>
<i>Diasporus hylaeformis</i>	<i>Norops intermedius</i>
<i>Diasporus igneus</i>	<i>Norops johnmeyeri</i>
<i>Diasporus majeensis</i>	<i>Norops kemptoni</i>
<i>Diasporus pequeno</i>	<i>Norops kreutzi</i>
<i>Diasporus sapo</i>	<i>Norops leditzigorum</i>
<i>Diasporus tigrillo</i>	<i>Norops magnaphallus</i>
<i>Diasporus ventrimaculatus</i>	<i>Norops marsupialis</i>
<i>Bromeliohyla melacaena</i>	<i>Norops monteverde</i>
<i>Dryophytes bocourti</i>	<i>Norops morazani</i>
<i>Duellmanohyla legleri</i>	<i>Norops muralla</i>
<i>Duellmanohyla lythrodus</i>	<i>Norops ocelloscapularis</i>
<i>Duellmanohyla rufioculis</i>	<i>Norops osa</i>
<i>Ecnomiohyla minera</i>	<i>Norops pachypus</i>
<i>Ecnomiohyla rabborum</i>	<i>Norops pijolensis</i>

The endemic herpetofauna of Central America

Table 10 (continued). Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

<i>Ecnomiohyla salvaje</i>	<i>Norops pseudokemptoni</i>
<i>Ecnomiohyla thysanota</i>	<i>Norops pseudopachypus</i>
<i>Ecnomiohyla veraguensis</i>	<i>Norops purpurgularis</i>
<i>Exerodonta catracha</i>	<i>Norops roatanensis</i>
<i>Exerodonta perkinsi</i>	<i>Norops rubribarbaris</i>
<i>Isthmohyla calypsa</i>	<i>Norops salvini</i>
<i>Isthmohyla debilis</i>	<i>Norops sminthus</i>
<i>Isthmohyla infucata</i>	<i>Norops tenorioensis</i>
<i>Isthmohyla insolita</i>	<i>Norops townsendi</i>
<i>Isthmohyla picadoi</i>	<i>Norops triumphalis</i>
<i>Isthmohyla pictipes</i>	<i>Norops tropidolepis</i>
<i>Isthmohyla xanthosticta</i>	<i>Norops utilensis</i>
<i>Isthmohyla zeteki</i>	<i>Norops villai</i>
<i>Plectrohyla calvata</i>	<i>Norops wampuensis</i>
<i>Plectrohyla dasypus</i>	<i>Norops wermuthi</i>
<i>Plectrohyla exquisita</i>	<i>Norops woodi</i>
<i>Plectrohyla psiloderma</i>	<i>Norops yoroensis</i>
<i>Plectrohyla tecunumani</i>	<i>Bachia blairi</i>
<i>Plectrohyla teuchestes</i>	<i>Potamites apodemus</i>
<i>Ptychohyla dendrophasma</i>	<i>Ctenosaura bakeri</i>
<i>Quilticohyla sanctaerucis</i>	<i>Ctenosaura oedirhina</i>
<i>Scinax alata</i>	<i>Ctenosaura palearis</i>
<i>Smilisca manisorum</i>	<i>Marisora alliacea</i>
<i>Smilisca puma</i>	<i>Marisora magnacornae</i>
<i>Leptodactylus silvanimbus</i>	<i>Marisora roatanae</i>
<i>Hypopachus pictiventris</i>	<i>Phyllodactylus insularis</i>
<i>Pipa myersi</i>	<i>Phyllodactylus palmeus</i>
<i>Lithobates lenca</i>	<i>Phyllodactylus paralepis</i>
<i>Lithobates miadis</i>	<i>Lepidoblepharis rufigularis</i>
<i>Lithobates vibicarius</i>	<i>Lepidoblepharis victormartinezi</i>
<i>Bolitoglossa anthracina</i>	<i>Sphaerodactylus alphas</i>
<i>Bolitoglossa aurae</i>	<i>Sphaerodactylus dunni</i>
<i>Bolitoglossa aureogularis</i>	<i>Sphaerodactylus graptolaemus</i>
<i>Bolitoglossa bramei</i>	<i>Sphaerodactylus guanaje</i>
<i>Bolitoglossa carri</i>	<i>Sphaerodactylus homolepis</i>
<i>Bolitoglossa cataguana</i>	<i>Sphaerodactylus leonardovaldesi</i>
<i>Bolitoglossa celaque</i>	<i>Sphaerodactylus pacificus</i>
<i>Bolitoglossa centenorum</i>	<i>Sphaerodactylus poindexteri</i>
<i>Bolitoglossa cerroensis</i>	<i>Sphaerodactylus rosaurae</i>
<i>Bolitoglossa chucantiensis</i>	<i>Scincella rara</i>
<i>Bolitoglossa compacta</i>	<i>Cnemidophorus duellmani</i>
<i>Bolitoglossa conanti</i>	<i>Holcosus miadis</i>
<i>Bolitoglossa copia</i>	<i>Dendrophidion paucicarinatum</i>
<i>Bolitoglossa cuchumatana</i>	<i>Oxybelis wilsoni</i>
<i>Bolitoglossa cuna</i>	<i>Tantilla albiceps</i>
<i>Bolitoglossa daryorum</i>	<i>Tantilla bairdi</i>

Table 10 (continued). Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

<i>Bolitoglossa decora</i>	<i>Tantilla berguidoi</i>
<i>Bolitoglossa diaphora</i>	<i>Tantilla gottei</i>
<i>Bolitoglossa diminuta</i>	<i>Tantilla hendersoni</i>
<i>Bolitoglossa dunni</i>	<i>Tantilla lempira</i>
<i>Bolitoglossa epimela</i>	<i>Tantilla olympia</i>
<i>Bolitoglossa eremia</i>	<i>Tantilla psittaca</i>
<i>Bolitoglossa gomezi</i>	<i>Tantilla stenigrammi</i>
<i>Bolitoglossa gracilis</i>	<i>Tantilla tecta</i>
<i>Bolitoglossa heiroreias</i>	<i>Tantilla tritaeniata</i>
<i>Bolitoglossa helmrichi</i>	<i>Tantilla vermiformis</i>
<i>Bolitoglossa huehuetenanguensis</i>	<i>Adelphicos daryi</i>
<i>Bolitoglossa indio</i>	<i>Adelphicos ibarrorum</i>
<i>Bolitoglossa insularis</i>	<i>Adelphis veraepacis</i>
<i>Bolitoglossa jacksoni</i>	<i>Atractus dariensis</i>
<i>Bolitoglossa jugivagans</i>	<i>Atractus depressiocellus</i>
<i>Bolitoglossa kamuk</i>	<i>Atractus hostilitractus</i>
<i>Bolitoglossa kaqchikelorum</i>	<i>Atractus imperfectus</i>
<i>Bolitoglossa la</i>	<i>Chapinophis xanthocheilus</i>
<i>Bolitoglossa longissima</i>	<i>Coniophanes joanae</i>
<i>Bolitoglossa magnifica</i>	<i>Cubophis brooksi</i>
<i>Bolitoglossa marmorea</i>	<i>Dipsas nicholsi</i>
<i>Bolitoglossa meliana</i>	<i>Dipsas tenuissima</i>
<i>Bolitoglossa minutula</i>	<i>Enulius bifoveatus</i>
<i>Bolitoglossa mombachoensis</i>	<i>Enulius roatanensis</i>
<i>Bolitoglossa nigrescens</i>	<i>Geophis bellus</i>
<i>Bolitoglossa ninadormida</i>	<i>Geophis championi</i>
<i>Bolitoglossa nussbaumi</i>	<i>Geophis damiani</i>
<i>Bolitoglossa obscura</i>	<i>Geophis downsi</i>
<i>Bolitoglossa omniumsanctorum</i>	<i>Geophis dunni</i>
<i>Bolitoglossa oresbia</i>	<i>Geophis fulvoguttatus</i>
<i>Bolitoglossa pacaya</i>	<i>Geophis godmani</i>
<i>Bolitoglossa pesrubra</i>	<i>Geophis nephodrymus</i>
<i>Bolitoglossa porrasorum</i>	<i>Geophis talamancae</i>
<i>Bolitoglossa psephena</i>	<i>Geophis zeledoni</i>
<i>Bolitoglossa pygmaea</i>	<i>Hydromorphus dunni</i>
<i>Bolitoglossa robinsoni</i>	<i>Imantodes phantasma</i>
<i>Bolitoglossa robusta</i>	<i>Leptodeira rubricata</i>
<i>Bolitoglossa sombra</i>	<i>Ninia celata</i>
<i>Bolitoglossa sooyorum</i>	<i>Ninia espinali</i>
<i>Bolitoglossa splendida</i>	<i>Omoadiphas aurula</i>
<i>Bolitoglossa subpalmata</i>	<i>Omoadiphas cannula</i>
<i>Bolitoglossa suchitanensis</i>	<i>Omoadiphas texiguatensis</i>
<i>Bolitoglossa synoria</i>	<i>Rhadinaea calligaster</i>
<i>Bolitoglossa taylori</i>	<i>Rhadinaea pulveriventris</i>
<i>Bolitoglossa tenebrosa</i>	<i>Rhadinella lisyae</i>
<i>Bolitoglossa tica</i>	<i>Rhadinella pegosalyta</i>

The endemic herpetofauna of Central America

Table 10 (continued). Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

<i>Bolitoglossa tzultacaj</i>	<i>Rhadinella pilonaorum</i>
<i>Bolitoglossa xibalba</i>	<i>Rhadinella rogerromani</i>
<i>Bolitoglossa zacapensis</i>	<i>Rhadinella tolpanorum</i>
<i>Cryptotriton monzoni</i>	<i>Sibon lamari</i>
<i>Cryptotriton necopinus</i>	<i>Sibon manzanaresi</i>
<i>Cryptotriton sierraminensis</i>	<i>Sibon merendonensis</i>
<i>Cryptotriton veraepacis</i>	<i>Sibon miskitus</i>
<i>Cryptotriton xucaneborum</i>	<i>Sibon noalamina</i>
<i>Dendrotriton bromeliacius</i>	<i>Sibon perissostichon</i>
<i>Dendrotriton chujorum</i>	<i>Trimetopon gracile</i>
<i>Dendrotriton cuchumatanus</i>	<i>Trimetopon slevini</i>
<i>Dendrotriton kekchiorum</i>	<i>Trimetopon viquezi</i>
<i>Dendrotriton rabbi</i>	<i>Urotheca myersi</i>
<i>Dendrotriton sanctibarbarus</i>	<i>Micrurus mosquitensis</i>
<i>Nototriton abscondens</i>	<i>Micrurus ruatanus</i>
<i>Nototriton barbouri</i>	<i>Micrurus stuarti</i>
<i>Nototriton brodiei</i>	<i>Epictia martinezi</i>
<i>Nototriton costaricense</i>	<i>Epictia pauldwyeri</i>
<i>Nototriton gamezi</i>	<i>Typhlops tycherus</i>
<i>Nototriton guanacaste</i>	<i>Atropoides indomitus</i>
<i>Nototriton lignicola</i>	<i>Bothriechis guifarroi</i>
<i>Nototriton limnospectator</i>	<i>Bothriechis lateralis</i>
<i>Nototriton major</i>	<i>Bothriechis marchi</i>
<i>Nototriton matama</i>	<i>Bothriechis nigroviridis</i>
<i>Nototriton mime</i>	<i>Bothriechis nubestris</i>
<i>Nototriton nelsoni</i>	<i>Bothriechis thalassinus</i>
<i>Nototriton oreadorum</i>	<i>Cerrophidion sasai</i>
<i>Nototriton picadoi</i>	<i>Cerrophidion wilsoni</i>
<i>Nototriton picucha</i>	<i>Porthidium porrasi</i>
<i>Nototriton richardi</i>	<i>Porthidium volcanicum</i>
<i>Nototriton saslaya</i>	<i>Kinosternon angustipons</i>
<i>Nototriton stuarti</i>	
Priority Two: High Vulnerability Species in Two Physiographic Regions (73)	
<i>Atelopus certus</i>	<i>Celestus atitlanensis</i>
<i>Incilius signifer</i>	<i>Dactyloa ibanezi</i>
<i>Rhinella centralis</i>	<i>Dactyloa kunayalae</i>
<i>Hyalinobatrachium diana</i>	<i>Dactyloa maia</i>
<i>Craugastor aurilegulus</i>	<i>Dactyloa savagei</i>
<i>Craugastor campbelli</i>	<i>Norops aquaticus</i>
<i>Craugastor charadra</i>	<i>Norops carpenteri</i>
<i>Craugastor epochthidius</i>	<i>Norops charlesmyersi</i>
<i>Craugastor evanesco</i>	<i>Norops cryptolimifrons</i>
<i>Craugastor fecundus</i>	<i>Norops loveridgei</i>
<i>Craugastor lauraster</i>	<i>Norops polylepis</i>
<i>Craugastor obesus</i>	<i>Norops wilsoni</i>
<i>Craugastor pechorum</i>	<i>Norops zeus</i>

Table 10 (continued). Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

<i>Craugastor persimilis</i>	<i>Heloderma charlesbogerti</i>
<i>Craugastor rostralis</i>	<i>Ctenosaura melanosterna</i>
<i>Craugastor rugosus</i>	<i>Ctenosaura praeocularis</i>
<i>Craugastor sabrinus</i>	<i>Ctenosaura quinquecarinata</i>
<i>Craugastor sandersoni</i>	<i>Sceloporus lunaei</i>
<i>Pristimantis altae</i>	<i>Mesoscincus managuae</i>
<i>Pristimantis caryophyllaceus</i>	<i>Lepidoblepharis emberawoundule</i>
<i>Oophaga granulifera</i>	<i>Cnemidophorus ruatanus</i>
<i>Oophaga vicentei</i>	<i>Dendrophidion crybelum</i>
<i>Ecnomiohyla bailarina</i>	<i>Mastigodryas dorsalis</i>
<i>Ecnomiohyla fimbrimembra</i>	<i>Tantilla jani</i>
<i>Ecnomiohyla sukia</i>	<i>Tantilla taeniata</i>
<i>Isthmohyla lancasteri</i>	<i>Geophis ruthveni</i>
<i>Agalychnis annae</i>	<i>Ninia pavimentata</i>
<i>Bolitoglossa alvaradoi</i>	<i>Rhadinaea sargenti</i>
<i>Bolitoglossa lignicolor</i>	<i>Rhadinella montecristi</i>
<i>Bolitoglossa nympa</i>	<i>Urotheca pachyura</i>
<i>Bolitoglossa odonnelli</i>	<i>Micrurus hippocrepis</i>
<i>Bolitoglossa salvinii</i>	<i>Bothriechis supraciliaris</i>
<i>Cryptotriton nasalis</i>	<i>Lachesis melanocephala</i>
<i>Oedipina alfaroi</i>	<i>Lachesis stenophrys</i>
<i>Oedipina alleni</i>	<i>Rhinoclemmys funerea</i>
<i>Oedipina ignea</i>	
<i>Oedipina stuarti</i>	
<i>Dermophis occidentalis</i>	
Priority Three: High Vulnerability Species in Three Physiographic Regions (27)	
<i>Craugastor chac</i>	<i>Echinosaura panamensis</i>
<i>Craugastor gollmeri</i>	<i>Ctenosaura flavidorsalis</i>
<i>Craugastor stejnegerianus</i>	<i>Sceloporus malachiticus</i>
<i>Pristimantis cerasinus</i>	<i>Drymobius melanotropis</i>
<i>Pristimantis pardalis</i>	<i>Scolecophis atrocinctus</i>
<i>Silverstoneia flotator</i>	<i>Dipsas articulata</i>
<i>Agalychnis saltator</i>	<i>Dipsas bicolor</i>
<i>Bolitoglossa schizodactyla</i>	<i>Rhadinaea vermiculaticeps</i>
<i>Oedipina pseudouniformis</i>	<i>Sibon anthracops</i>
<i>Diploglossus bilobatus</i>	<i>Sibon carri</i>
<i>Dactyloa insignis</i>	<i>Trimetopon barbouri</i>
<i>Norops lionotus</i>	<i>Micrurus stewarti</i>
<i>Norops macrophallus</i>	<i>Atropoides picadoi</i>
<i>Norops quaggulus</i>	
Priority Four: High Vulnerability Species in Four Physiographic Regions (21)	
<i>Craugastor megacephalus</i>	<i>Norops wellbornae</i>
<i>Craugastor mimus</i>	<i>Holcosus leptophrys</i>
<i>Craugastor ranoides</i>	<i>Holcosus quadrilineatus</i>
<i>Diasporus diastema</i>	<i>Leptophis nebulosus</i>
<i>Bolitoglossa colonnea</i>	<i>Crisantophis nevermanni</i>

The endemic herpetofauna of Central America

Table 10 (continued). Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

<i>Bolitoglossa dofleini</i>	<i>Rhadinella anachoreta</i>
<i>Dactyloa brooksi</i>	<i>Sibon argus</i>
<i>Norops apletophallus</i>	<i>Sibon longifrenis</i>
<i>Norops elcopeensis</i>	<i>Micrurus alleni</i>
<i>Norops humilis</i>	<i>Agkistrodon howardgloydi</i>
<i>Norops oxylophus</i>	
Priority Five: High Vulnerability Species in Five Physiographic Regions (9)	
<i>Cochranella granulosa</i>	<i>Coleonyx mitratus</i>
<i>Craugastor noblei</i>	<i>Marisora unimarginata</i>
<i>Bolitoglossa striatula</i>	<i>Dendrophidion rufiterminorum</i>
<i>Gymnopsis multiplicata</i>	<i>Trimorphodon quadruplex</i>
<i>Basiliscus plumifrons</i>	
Priority Six: High Vulnerability Species in Six Physiographic Regions (3)	
<i>Norops limifrons</i>	<i>Dendrophidion apharocybe</i>
<i>Sphaerodactylus millepunctatus</i>	
Priority Seven: Medium Vulnerability Species in Single Physiographic Region (23)	
<i>Atelopus senex</i>	<i>Plectrohyla pokomchi</i>
<i>Incilius chompipe</i>	<i>Plectrohyla quechi</i>
<i>Incilius fastidiosus</i>	<i>Lithobates juliani</i>
<i>Incilius melanochlorus</i>	<i>Lithobates taylori</i>
<i>Rhinella chrysophora</i>	<i>Bolitoglossa morio</i>
<i>Isthmohyla angustilineata</i>	<i>Dactyloa ginaelisae</i>
<i>Isthmohyla graceae</i>	<i>Norops cobanensis</i>
<i>Isthmohyla pseudopuma</i>	<i>Ninia psephota</i>
<i>Isthmohyla rivularis</i>	<i>Rhadinaea stadelmani</i>
<i>Isthmohyla tica</i>	<i>Rhadinella hempsteadae</i>
<i>Plectrohyla chrysopleura</i>	<i>Rhadinella serperaster</i>
<i>Plectrohyla glandulosa</i>	
Priority Eight: Medium Vulnerability Species in Two Physiographic Regions (21)	
<i>Atelopus zeteki</i>	<i>Norops mcraniei</i>
<i>Incilius ibarraii</i>	<i>Lepidophyma mayae</i>
<i>Incilius leucomyos</i>	<i>Lepidophyma reticulatum</i>
<i>Craugastor bransfordii</i>	<i>Helminthophis frontalis</i>
<i>Atlantihyla panchoi</i>	<i>Tantilla brevicauda</i>
<i>Atlantihyla spinipollex</i>	<i>Tantilla excelsa</i>
<i>Duellmanohyla salvadorensis</i>	<i>Geophis tectus</i>
<i>Duellmanohyla salvavida</i>	<i>Trimetopon plioplepis</i>
<i>Duellmanohyla soralia</i>	<i>Trimetopon simile</i>
<i>Duellmanohyla uranochroa</i>	<i>Amerotyphlops stadelmani</i>
<i>Hyloscirtus colymba</i>	
Priority Nine: Medium Vulnerability Species in Three Physiographic Regions (5)	
<i>Atelopus varius</i>	<i>Tantilla ruficeps</i>
<i>Craugastor laevisimus</i>	<i>Geophis brachycephalus</i>
<i>Ptychohyla hypomykter</i>	
Priority Ten: Medium Vulnerability Species in Four Physiographic Regions (4)	
<i>Lithobates warszewitschii</i>	<i>Tantilla armillata</i>

Table 10 (continued). Conservation priority listing of the endemic herpetofaunal species in Central America based on the EVS categorization and the range of physiographic occurrence.

<i>Leptodrymus pulcherrimus</i>	<i>Amerotyphlops costaricensis</i>
Priority Eleven: Medium Vulnerability Species in Five Physiographic Regions (4)	
<i>Mastigodryas alternatus</i>	<i>Urotheca guentheri</i>
<i>Ninia maculate</i>	<i>Epictia ater</i>
Priority Twelve: Medium Vulnerability Species in Six Physiographic Regions (2)	
<i>Norops cupreus</i>	<i>Geophis hoffmanni</i>
Priority Thirteen: Medium Vulnerability Species in Seven Physiographic Regions (1)	
<i>Hydromorphus concolor</i>	
Priority Fourteen: Medium Vulnerability Species in Eight Physiographic Regions (1)	
<i>Leptodeira rhombifera</i>	



Plate 28. *Dactyloa insignis* (Cope, 1871). The Decorated Anole is a priority four species with an EVS of 14, with a distribution restricted to “the Cordillera Tilarán and Cordillera Central of Costa Rica” (Poe and Ryan 2017: 6). This individual was encountered in Estación Pocosol, Bosque Eterno de los Niños, in the province of Alajuela, Costa Rica. *Photo by Victor Acosta-Chaves.*



Plate 29. *Dactyloa kunayalae* (Huleback, Poe, Ibáñez, and Williams, 2007). This anole is a priority two species with an EVS of 15, restricted in distribution to “central Panama” (Köhler 2008: 100). This individual came from Parque Nacional General de División Omar Torrijos Herrera, in the province of Coclé, Panama. *Photo by Abel Batista.*



Plate 30. *Norops carpenteri* (Echelle, Echelle, and Fitch, 1971). Carpenter’s Anole is a priority two species with an EVS of 16, distributed “on the Atlantic versant from northeastern Honduras to northwestern Panama” (McCranie and Köhler 2015: 45). This individual was found in Reserva El Copal, Jiménez, Costa Rica. *Photo by Victor Acosta-Chaves.*



Plate 31. *Norops kemptoni* (Dunn, 1940). Kempton’s Anole is a priority one species with an EVS of 15, found in the “highlands of Chiriquí, Panama” (Köhler 2008: 106). This individual is from Alto Chiquero, Parque Nacional Volcán Barú, in the province of Chiriquí, Panama. *Photo by Javier Sunyer.*



Plate 32. *Coleonyx mitratus* (Peters, 1863). The Central American Banded Gecko is a priority six species with an EVS of 14, which ranges along the “Atlantic lowlands of northeastern Guatemala and northwestern Honduras [and the] Pacific lowlands from Guatemala to southwestern Costa Rica” (Savage 2002: 482). This individual was located in Sector Santa Rosa, Parque Nacional Santa Rosa, in the province of, Guanacaste, Costa Rica. *Photo by Victor Acosta-Chaves.*



Plate 34. *Sceloporus malachiticus* Cope, 1864. The Green Spiny Lizard is a priority three species with an EVS of 10, distributed from “El Salvador and Honduras across Nicaragua and Costa Rica to Panama” (Köhler 2008: 152). This individual was encountered in Cerro de la Muerte, in the province of San José, Costa Rica. *Photo by Victor Acosta-Chaves.*



Plate 36. *Rhadinaea calligaster* (Cope, 1876). The Thick Graceful Brownsnake is a priority one species with an EVS of 14, distributed in “the Cordillera de Tilarán, Cordillera Central, and Cordillera de Talamanca of Costa Rica and on Volcán Tenorio in the Cordillera de Guanacaste...to extreme western Panama” (Savage 2002: 623). This individual came from San Gerardo de Dota, in the province of San José, Costa Rica. *Photo by Victor Acosta-Chaves.*



Plate 33. *Heloderma charlesbogerti* Campbell and Vanini, 1988. The Guatemalan Beaded Lizard is a priority two species that inhabits “the Río Motagua Valley, in the Atlantic versant of eastern Guatemala” (Reiserer et al. 2013: 81). This individual was found at Cabañas, in the department of Zacapa, Guatemala. *Photo by Andres Novales.*



Plate 35. *Scolocophis atrocinctus* (Schlegel, 1837). priority three species with an EVS of 13, which ranges along the “Pacific versant from southeastern Guatemala to northwestern Costa Rica; the species also is found on the Atlantic versant in southwestern Honduras, western Nicaragua, and northwestern Costa Rica” (Wilson and Mata-Silva 2015: 422). This individual was found in Tilarán, in the province of Guanacaste, Costa Rica. *Photo by Victor Acosta-Chaves.*



Plate 37. *Bothriechis lateralis* Peters, 1862. The Coffee Palmviper is a priority one species with an EVS of 16. distributed in “the cordilleras of Costa Rica and western Panama (Savage, 2002: 724). This individual came from La Nevera, Serranía de Tabasará, Panama. *Photo by Abel Batista.*

F. The Central American endemic herpetofauna is distributed unevenly among the 10 physiographic regions we recognize in this portion of Mesoamerica. The number of endemic species in these regions ranges from six in the Yucatan Platform to 254 in the Isthmian Central American highlands. Most of the 623 endemic species are limited to a single physiographic region, i.e., 450 (72.2%). The next largest number of 95 (15.2%) is for those occupying two regions. Thus, 545 (87.5%) of the species occur in only one or two of the 10 regions.

G. An implementation of the EVS system of conservation assessment demonstrates that all of the scores for the 623 endemic species in Central America lie within the medium vulnerability (63) or high vulnerability (560) categories.

H. We used the same means as Johnson et al. (2017) for the Mexican endemic species to prioritize the conservation significance of the Central American endemic species, i.e., by combining the data on physiographic distribution with that on EVS. This procedure allowed us to identify 14 priority levels, of which six are high vulnerability and eight are medium vulnerability groupings.

I. The number of species occupying the six high vulnerability levels decrease markedly and consistently from 429 in Priority Level One (high vulnerability species occupying a single physiographic region) to one in Priority Level Six (high vulnerability species occurring in six physiographic regions). The total number of species allocated to the seven priority levels amounts to 562, 90.2% of the 623 Central American endemic species.

J. The number of species occurring in the eight medium vulnerability levels also decreases consistently, but not as markedly, from 23 in Priority Level Seven (medium vulnerability species occupying single physiographic regions) to one in Priority Level Fourteen (medium vulnerability species distributed in eight physiographic regions). The total number of species placed in these eight priority levels is 61, 9.8% of the total number of Central American endemics.

K. Our analysis demonstrates that most Central American endemic species are assessed as high vulnerability species that occupy only one or two physiographic regions. A significant number of species (44) are of medium vulnerability, and they also inhabit one or two physiographic regions.

L. Protecting the endemic component of the globally significant Central American herpetofauna represents the most significant challenge for conservation professionals working in this portion of Mesoamerica.

M. One of the conclusions of the conservation analyses we have conducted in recent years, including in this paper, is that perpetual protection of the Mesoamerican herpetofauna presently is a goal far from realization. Our prognosis is that this goal will not be attained until humans are willing to address the widespread problems created by the anthropocentric worldview, and the impediment

it represents for allowing the environment “to function indefinitely without going into a decline from the stresses imposed by human society on natural systems such as fertile soil, water, and air” (Raven and Berg 2004: G-15).

N. If there is any merit to our hypothesis that anthropocentrism is part of a cascade of psychological ailments, which extend through ethnocentrism and culminate in the narcissistic personality disorder, it might predict that the critical-thinking educational reform called for by Johnson et al. (2017) will have to be recognized as requiring species-wide psychotherapy to treat a species-wide mental disease. If so, addressing this disease will be the largest problem undertaken by humanity during its existence on planet Earth.

Recommendations

A. The recommendations presented in the Johnson et al. (2017: 616) study on the Mexican endemic herpetofauna dealt with the establishment of a global coalition “to document all of Earth’s inhabitants within the 21st century, and to provide for their perpetual protection.” This recommendation applies here as well.

B. Johnson et al. (2017) also recommended that the system of prioritization they developed could help determine how funding for a countrywide system of sustainable reserves could be best utilized to protect the Mexican endemic herpetofauna. The same can be said for the Central American endemic herpetofauna, with respect to the seven nations comprising this region of the world.

C. We propose that until such steps are taken, the Central American endemic herpetofauna, as well as the entire planetary biota, will become a casualty of anthropocentrism. Nonetheless, these steps only will constitute stopgap measures, as the underlying anthropocentric worldview held by most humans will continue to accelerate the degradation of the planetary life-support systems, and eventually will render the planet unsuitable to support life.

“... Each loss of a species is a loss of a companion on the long journey of evolution ...”

—David W. Orr (2016).

Acknowledgments.—We thank the people who graciously allowed us to use their beautiful images of various Central American endemic species to illustrate this paper, including Victor Acosta-Chaves, César Barrio-Amorós, Abel Batista, Andres Novales, Todd Pierson, Sean Michael Rovito, and Javier Sunyer. We also are appreciative of the assistance of Alan Resetar of the Field Museum of Natural History in providing us with photographs of FMNH 74376, the specimen tentatively identified as *Dipsas viguieri* by Cadle (2005). Abel Batista, Sebastian Lotzkat, and Louis W. Porras significantly improved the quality of the manuscript with their outstanding reviews. We are indebted hugely to these colleagues for sharing their expertise with us.



Plate 38. *Bothriechis nigroviridis* Peters, 1859. The Black-speckled Palm-pitviper is a priority one species with an EVS of 17, which ranges along “the cordilleras of Costa Rica and western Panama” (Savage, 2002: 725). This individual was found at Jurutungo, in the province of Chiriquí, Panama. *Photo by Javier Sunyer.*



Plate 39. *Lachesis stenophrys* Cope, 1875. The Central American Bushmaster is a priority two species with an EVS of 17, distributed from “southeastern Nicaragua to central Panama” (Köhler 2008: 330). This individual came from Parque Nacional Braulio Carrillo, in the province of Heredia, Costa Rica. *Photo by César Barrio-Amorós.*



Plate 40. *Porthidium porrasi* Lamar, 2003. The White-tailed Hog-nosed Pitviper is a priority one species with an EVS of 18, with a distribution restricted to the Península de Osa and adjacent areas of southwestern Costa Rica (Solórzano 2004). Pictured here is an individual from this region, in the province of Puntarenas, Costa Rica. *Photo by César Barrio-Amorós, courtesy of Roel de Plecker.*

Literature Cited

- Ariano-Sánchez D, Campbell JA. 2018. A new species of the *Rhadinella* (Serpentes: Dipsadidae) from the dry forest of Motagua Valley, Guatemala. *Zootaxa* 4442: 338–344.
- Arias E, Chaves G, Crawford AJ, Parra-Olea G. 2016. A new species of the *Craugastor podiciferus* species group (Anura: Craugastoridae) from the premontane forest of southwestern Costa Rica. *Zootaxa* 4132: 347–363.
- Arias E, Kubicki B. 2018. A new moss salamander, genus *Nototriton* (Caudata: Plethodontidae), from the Cordillera de Talamanca, in the Costa Rica-Panama border region. *Zootaxa* 4369: 487–500.
- Batista A, Köhler G, Mebert K, Vesely M. 2014. A new species of *Bolitoglossa* (Amphibia: Plethodontidae) from eastern Panama, with comments on other members of the *adpersa* species group from eastern Panama. *Mesoamerican Herpetology* 1: 96–121.
- Batista A, Köhler G, Mebert K, Hertz A, Vesely M. 2016a. An integrative approach to reveal speciation and species richness in the genus (Amphibia: Anura: Eleutherodactylidae) in eastern Panama. *Zoological Journal of the Linnean Society* 178: 267–311.
- Batista A, Mebert K, Lotzkat S, Wilson LD. 2016b. A new species of centipede snake of the genus *Tantilla* (Squamata: Colubridae) from an isolated premontane forest in eastern Panama. *Mesoamerican Herpetology* 3: 949–960.
- Batista A, Ponce M, Vesely M, Mebert K, Hertz A, Köhler G, Carrizo A, Lotzkat S. 2015a. Revision of the genus *Lepidoblepharis* (Reptilia: Squamata: Sphaerodactylidae) in Central America, with the description of three new species. *Zootaxa* 3994: 187–221.
- Batista A, Vesely M, Mebert K, Lotzkat S, Köhler G. 2015b. A new species of *Dactyloa* from eastern Panama, with comments on other *Dactyloa* species present in the region. *Zootaxa* 4039: 57–84.
- Brodie ED Jr, Acevedo ME, Campbell JA. 2012. New salamanders of the genus *Oedipina* (Caudata: Plethodontidae) from Guatemala. *Journal of Herpetology* 46: 233–240.
- Cadle JE. 2005. Systematics of snakes of the *Dipsas oreas* complex (Colubridae: Dipsadinae) in western Ecuador and Peru, with revalidation of *D. elegans* (Boulenger) and *D. ellipsifera* (Boulenger). *Bulletin of the Museum of Comparative Zoology* 158: 67–136.
- Caligor E, Levy KN, Yeomans FE. 2015. Narcissistic personality disorder: diagnostic and clinical challenges. *The American Journal of Psychiatry* 172: 415–422.
- Campbell JA. 1999. Distribution patterns of amphibians in Middle America. Pp. 111–210 In: *Patterns of Distribution of Amphibians: A Global Perspective*. Editor, Duellman WE. The Johns Hopkins University Press, Baltimore, Maryland, USA. 633 p.
- Carr A. 1953. *High Jungles and Low*. University of Florida Press, Gainesville, Florida, USA. 226 p.
- Ceballos G, Ehrlich PR, Dirzo R. 2017. Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. *Proceedings of the National Academy of Sciences* 114(30): E6089–E6096.
- De Young R. 2013. Environmental psychology overview. Pp. 17–33 In: *Green Organizations: Driving Change with IO Psychology*. Editors, Huffman AH, Klein SR. Routledge Academic, New York, New York, USA. 440 p.
- Doan TM, Mason AJ, Castoe TA, Sasa M, Parkinson CL. 2016. A cryptic palm-pitviper species (Squamata: Viperidae: *Bothriechis*) from the Costa Rican highlands, with notes on the variation within *B. nigroviridis*. *Zootaxa* 4138: 271–290.
- Frost DR. 2018. Amphibian Species of the World: An Online Reference. Version 6.0. Available: www.research.amnh.org/herpetology/amphibia/index.html [Accessed 07 March 2018]. American Museum of Natural History, New York, New York, USA.
- García-Vázquez UO, Nieto-Montes de Oca A, Bryson RW Jr, Schmidt-Ballardo W, Pavón-Vázquez CJ. 2018. Molecular systematics and historical biogeography of the genus *Gerrhonotus* (Squamata: Anguillidae). *Journal of Biogeography* 45(7): 1,640–1,652.
- Harvey MB. 2008. New and poorly known *Dipsas* (Serpentes: Colubridae) from northern South America. *Herpetologica* 64: 422–451.
- Hofmann EP, Townsend JH. 2018. A cryptic new species of anole (Squamata: Dactyloidae) from the Lenca Highlands of Honduras, previously referred to as *Norops crassulus* (Cope, 1864). *Annals of Carnegie Museum* 85: 91–111.
- Jadin RC, Townsend JH, Castoe TA, Campbell JA. 2012. Cryptic diversity in disjunct populations of Middle American Montane Pitvipers: a systematic reassessment of *Cerrophidion godmani*. *Zoologica Scripta* 41: 455–470.
- Jaramillo C, Wilson LD, Ibáñez R, Jaramillo F. 2010. The herpetofauna of Panama: distribution and conservation status. Pp. 604–673 In: *Conservation of Mesoamerican Amphibians and Reptiles*. Editors, Wilson LD, Townsend JH, Johnson JD. Eagle Mountain Publishing, LC, Eagle Mountain, Utah, USA. 812 p.
- Johnson JD, Wilson LD, Townsend JH. 2010. Summary of recent changes in higher taxonomic categories of Mesoamerican amphibians and reptiles. Pp. 18–27 In: *Conservation of Mesoamerican Amphibians and Reptiles*. Editors, Wilson LD, Townsend JH, Johnson JD. Eagle Mountain Publishing, LC, Eagle Mountain, Utah USA. 812 p.
- Johnson JD, Mata-Silva V, Wilson LD. 2015. A conservation reassessment of the Central American herpetofauna based on the EVS measure. *Amphibian & Reptile Conservation* 9: 1–94 (e100).
- Johnson JD, Wilson LD, Mata-Silva V, García-Padilla

- E, DeSantis DL. 2017. The endemic herpetofauna of Mexico: organisms of global significance in severe peril. *Mesoamerican Herpetology* 4: 543–620.
- Köhler G. 2001. *Reptilien und Amphibien Mittelamerikas*. Band 2: Schlangen. Herpeton Verlag, Offenbach, Germany. 160 p.
- Köhler G. 2003. *Reptiles of Central America*. Herpeton Verlag, Offenbach, Germany. 367 p.
- Köhler G. 2008. *Reptiles of Central America*. 2nd edition. Herpeton, Offenbach, Germany. 400 p.
- Köhler G, Townsend JH, Petersen CBP. 2016. Taxonomic revision of the *Norops tropidonotus* complex (Squamata, Dactyloidae), with the resurrection of *N. spilorhipis* (Alvarez del Toro and Smith, 1956) and the description of two new species. *Mesoamerican Herpetology* 3: 8–41.
- Köhler JJ, Poe S, Ryan MJ, Köhler G. 2015. *Anolis marsupialis* Taylor 1956, a valid species from southern Pacific Costa Rica (Reptilia, Squamata, Dactyloidae). *Zootaxa* 3915: 111–122.
- Kopnina H. 2016. Half the Earth for people (or more?) Addressing ethical questions in conservation. *Biological Conservation* 203: 176–185.
- Kubicki B. 2016. A new species of salamander (Caudata: Plethodontidae: *Oedipina*) from the central Caribbean foothills of Costa Rica. *Mesoamerican Herpetology* 3: 819–840.
- Kubicki B, Arias E. 2016. A beautiful new yellow salamander, genus *Bolitoglossa* (Caudata: Plethodontidae), from the northwestern slopes of the Cordillera de Talamanca, Costa Rica. *Zootaxa* 4184: 329–346.
- Kubicki B, Salazar S, Puschendorf R. 2015. A new species of glassfrog, genus *Hyalinobatrachium* (Anura: Centrolenidae), from the Caribbean foothills of Costa Rica. *Zootaxa* 3920: 69–84.
- Lotzkat S, Hertz A, Köhler G. 2016. A new species of *Celestus* (Squamata: Anguillidae) from western Panama. *Mesoamerican Herpetology* 3: 962–975.
- Luque-Montes I, Austin JD, Weinfurter KD, Wilson LD, Hofmann EP, Townsend JH. 2018. An integrative assessment of the taxonomic status of putative hybrid leopard frogs (Anura: Ranidae) from the Chortís Highlands of Central America, with description of a new species. *Systematics and Biodiversity* 2018: 1–17.
- McCranie JR. 2017a. Specific status of the Montaña de Celaque Honduran frogs previously referred to as *Plectrohyla guatemalensis* (Anura: Hylidae: Hylinae). *Mesoamerican Herpetology* 4: 389–401.
- McCranie JR. 2017b. A new species of *Rhadinella* (Serpentes: Dipsadidae) from the Sierra de Agalta, Honduras. *Mesoamerican Herpetology* 4: 243–253.
- McCranie JR. 2017c. Morphological and systematic comments on the Caribbean lowland population of *Smilisca baudinii* (Anura: Hylidae: Hylinae) in northeastern Honduras, with the resurrection of *Hyla manisorum* Taylor. *Mesoamerican Herpetology* 4: 512–526.
- McCranie JR. 2018. A discussion of the phenetic-based *Craugastor laticeps* species group (Anura: Brachycephaloidea: Craugastoridae) from north-central Honduras, with the description of two new species. *Herpetologica* 72: 169–180.
- McCranie JR, Gotte SW. 2014. An investigation into the Swan Island Honduras collecting event of *Tiaporus fuliginosus* Cope (Reptilia: Teiidae) and its systematic status. *Proceedings of the Biological Society of Washington* 127: 543–556.
- McCranie JR, Köhler G. 2015. The Anoles of Honduras (Reptilia: Squamata: Dactyloidae: *Anolis*: *Norops*) of Honduras. Systematics, distribution, and conservation. *Bulletin of the Museum of Comparative Zoology* SPS(1): 1–292.
- McCranie JR, Smith ER. 2017. A review of the *Tantilla taeniata* species group (Reptilia: Squamata: Colubridae: Colubrinae) in Honduras, with the description of three new species. *Herpetologica* 73: 338–348.
- McCranie JR, Townsend JH, Wilson LD. 2006. *The Amphibians and Reptiles of the Honduran Mosquitia*. Krieger Publishing Company, Malabar, Florida, USA. 291 p.
- Meza-Lázaro RN, Nieto-Montes de Oca A. 2015. Long forsaken species diversity in the Middle American lizard *Holcosus undulatus* (Teiidae). *Zoological Journal of the Linnean Society* 175(1): 189–210.
- Nicholson KE, Köhler G. 2014. A new species of the genus *Norops* from Darién, Panama with comments on *N. sulcifrons* (Cope 1899) (Reptilia, Squamata, Dactyloidae). *Zootaxa* 3895: 225–237.
- Orr DW. 2016. *Dangerous Years: Climate Change, the Long Emergency, and the Way Forward*. Yale University Press, New Haven, Connecticut, USA. 320 p.
- Pérez Santos, C, Moreno, AG. 1988. *Ofidios de Colombia*. Monografía VI. Museo Regionale di Scienze Naturali Torino, Torino, Italy. 517 p.
- Pérez Santos, C. 1999. *Serpientes de Panamá / Snakes of Panama*. BIOSFERA, Publicaciones del Comité Español del Programa MaB y de la Red IberoMaB de la UNESCO, Madrid, Spain. 312 p.
- Peters JA. 1960. The snakes of the subfamily Dipsadinae. *Miscellaneous Publications of the Museum of Zoology, University of Michigan* 114: 1–224.
- Poe S, Ryan MJ. 2017. Description of two new species similar to *Anolis insignis* (Squamata: Iguanidae) and resurrection of *Anolis (Diaphoranolis) brooksi*. *Amphibian & Reptile Conservation* 11: 1–16 (e141).
- Poe S, Scarpetta S, Schaad EW. 2015. A new species of *Anolis* (Squamata: Iguanidae) from Panama. *Amphibian & Reptile Conservation* 9: 1–13 (e194).
- Pyron RA. 2017. We don't need to save endangered species. Extinction is part of evolution. *The Washington Post* (22 November 2017).
- Raven PH, Berg LR. 2004. *Environment*. 4th edition. John Wiley & Sons, Inc., Hoboken, New Jersey, USA. 688 p.

- Ray J. 2017. *Snakes of Panama: A Field Guide to All Species*. Team Snake Panama, CreateSpace Independent Publishing Platform, Lexington, Kentucky, USA. 213 p.
- Rovito SM, Vásquez-Almazán CR, Papenfuss TJ, Parra-Olea G, Wake DB. 2015. Biogeography and evolution of Central American cloud forest salamanders (Caudata: Plethodontidae: *Cryptotriton*), with the description of a new species. *Zoological Journal of the Linnean Society* 175: 150–166.
- Savage JM, Ugarte CA, Donnelly MA. 2013. A new species of earless toad (Bufonidae: *Incilius*) from western Panama. *Copeia* 2013: 8–12.
- Solano-Zavaleta I, Nieto-Montes de Oca A, Campbell JA. 2016. A new species of *Mesaspis* (Squamata: Anguillidae) from the high Cuchumatanes of Guatemala. *Journal of Herpetology* 50: 327–335.
- Solano-Zavaleta I, Nieto-Montes de Oca A. 2018. Species limits in the Morelet's Alligator Lizard (Anguillidae: Gerrhonotinae). *Molecular Phylogenetics and Evolution* 120: 16–27.
- Solís JM, Espinal MR, Valle RE, O'Reilly CM, Itgen MS, Townsend JH. 2016. On the taxonomy of *Oedipina stuarti* (Caudata: Plethodontidae), with description of a new species from suburban Tegucigalpa, Honduras. *Salamandra* 42: 125–133.
- Steg L, Vlek C. 2009. Encouraging pro-environmental behaviour: an integrative review and research agenda. *Journal of Environmental Psychology* 29: 309–317.
- Townsend JH. 2016. Taxonomic revision of the moss salamander *Nototriton barbouri* (Schmidt) (Caudata: Plethodontidae), with description of two new species from the Cordillera Nombre de Dios, Honduras. *Zootaxa* 4196: 511–528.
- Townsend JH. 2014. Characterizing the Chortís Block biogeographic province: geological, physiographic, and ecological association and herpetofaunal diversity. *Mesoamerican Herpetology* 1: 203–252.
- Townsend JH, Wilson LD. 2008. *Guide to the Amphibians and Reptiles of Cusuco National Park, Honduras/Guía de los Anfíbios y Reptiles del Parque Nacional El Cusuco, Honduras*. Bibliomania!, Salt Lake City, Utah, USA. 322 p.
- Townsend JH, Wilson LD. 2010a. Conservation of the Honduran herpetofauna: issues and imperatives. Pp. 460–487 In: *Conservation of Mesoamerican Amphibians and Reptiles*. Editors, Wilson LD, Townsend JH, Johnson JD. Eagle Mountain Publishing, LC, Eagle Mountain, Utah, USA. 812 p.
- Townsend JH, Wilson LD. 2010b. Biogeography and conservation of the Honduran subhumid forest herpetofauna. Pp. 686–705 In: *Conservation of Mesoamerican Amphibians and Reptiles*. Editors, Wilson LD, Townsend JH, Johnson JD. Eagle Mountain Publishing, LC, Eagle Mountain, Utah, USA. 812 p.
- Townsend JH, Wilson LD. 2016. Amphibians of the Cordillera Nombre de Dios, Honduras: COI barcoding suggests underestimated taxonomic richness in a threatened endemic fauna. *Mesoamerican Herpetology* 3: 909–928.
- Wallach V. 2016. Morphological review and taxonomic status of the *Epictia phenops* species group of Mesoamerica, with description of six new species and discussion of South American *Epictia albifrons*, *E. goudotii*, and *E. tenella* (Serpentes: Leptotyphlopidae: Epictinae). *Mesoamerican Herpetology* 3: 216–374.
- Wallach V, Williams KL, Boundy J. 2014. *Snakes of the World: A Catalogue of Living and Extinct Species*. CRC Press, Taylor & Francis Group, Boca Raton, Florida, USA. 1,237 p.
- Wilson EO. 2016. *Half-Earth: Our Planet's Fight for Life*. Liveright Publishing Corporation, New York, New York, USA. 272 p.
- Wilson LD, McCranie JR. 2004. The conservation status of the herpetofauna of Honduras. *Amphibian & Reptile Conservation* 3: 6–33 (e12).
- Wilson LD, Johnson JD. 2010. Distributional patterns of the herpetofauna of Mesoamerica, a biodiversity hotspot. Pp. 30–235 In: *Conservation of Mesoamerican Amphibians and Reptiles*. Editors, Wilson LD, Townsend JH, Johnson JD. Eagle Mountain Publishing, LC, Eagle Mountain, Utah, USA. 812 p.
- Wilson LD, Johnson JD, Porras LW, Mata-Silva V, García-Padilla E. 2017. A system for categorizing the distribution of the Mesoamerican herpetofauna. *Mesoamerican Herpetology* 4: 901–913.
- Wilson LD, Mata-Silva V. 2015. A checklist and key to the snakes of the *Tantilla* clade (Squamata: Colubridae), with comments on taxonomy, distribution, and conservation. *Mesoamerican Herpetology* 2: 418–498.
- Wilson LD, Townsend JH, Johnson JD. 2010. Preface. Pp. xiv–xvii In: *Conservation of Mesoamerican Amphibians and Reptiles*. Editors, Wilson LD, Townsend JH, Johnson JD. Eagle Mountain Publishing, LC, Eagle Mountain, Utah USA. 812 p.
- Wilson LD, Townsend JH. 2010. The herpetofauna of Mesoamerica: biodiversity significance, conservation status, and future challenges. Pp. 760–812 In: *Conservation of Mesoamerican Amphibians and Reptiles*. Editors, Wilson LD, Townsend JH, Johnson JD. Eagle Mountain Publishing, LC, Eagle Mountain, Utah, USA. 812 p.
- Woolrich-Piña GA, García-Padilla E, DeSantis DL, Johnson JD, Mata-Silva V, Wilson LD. 2017. The herpetofauna of Puebla, Mexico: composition, distribution, and conservation status. *Mesoamerican Herpetology* 4: 790–884.

Addendum (changes past conclusion of analyses)

We chose a cut-off date of 30 April 2018 to discontinue revising the huge number of calculations dealing with the 623 endemic herpetofaunal species documented in Central America. Past this date, we added pertinent taxa and publications to this addendum until the correction of the proof, as follows:

(1) *Craugastor aenigmaticus*. Arias et al. (2018) described this new species of rainfrog from the Cordillera de Talamanca in southern Costa Rica. This frog is known from several localities in montane rainforest, and its EVS can be calculated as $5+8+4=17$, placing it in the middle portion of the high vulnerability category. This species is limited to a single physiographic region, the Isthmian Central American highlands, and can be placed in conservation priority level one.

(2) *Craugastor castanedai*. McCranie (2018) described this new species of rainfrog from the Parque Nacional Pico Bonito in north-central Honduras. This frog is known from only two localities on either side of the Quebrada de Oro, and its EVS can be calculated as $6+8+4=18$, placing it in the upper portion of the high vulnerability category. This species is limited to a single physiographic region, the eastern nuclear Central American highlands, and can be placed in conservation priority level one.

(3) *Craugastor gutschei*. McCranie (2018) also described this new species of rainfrog from the western portion of the Cordillera Nombre de Dios in north-central Honduras. Its EVS can be calculated as $5+7+4=16$, placing it in the middle portion of the high vulnerability category. This species is restricted to the eastern nuclear Central American highlands, and can be allocated to conservation priority level one.

(4) *Hemiphractus elioti*. Hill et al. (2018) described this new species of horned frog from the Cordillera de Talamanca in western Panama. Its EVS can be determined as $5+7+5=17$, placing it in the middle portion of the high vulnerability category. This species is restricted to the Isthmian Central American highlands, and can be allocated to conservation priority level one.

(5) *Hemiphractus kaylockae*. Hill et al. (2018) named this new species of horned frog from the highlands of eastern

Panama. Its EVS can be estimated as $6+8+5=19$, placing it in the upper portion of the high vulnerability category. This species is restricted to the eastern Panamanian highlands, and can be placed in conservation priority level one.

(6) *Hemiphractus panamensis*. Hill et al. (2018) revalidated this taxon of horned frogs from the eastern highlands of Panama. Its EVS can be determined as $5+8+5=18$, placing it in the upper portion of the high vulnerability category. This species is restricted to the western portion of the eastern Panamanian highlands in the central portion of the country, and can be placed in conservation priority level one.

(7) *Hemiphractus fasciatus*. Hill et al. (2018) determined that this species, formerly considered to be the single representative of this peculiar hylid genus in Central America, should be considered to be a South American taxon not resident in Panama.

(8) *Norops caceresae*. Hofmann and Townsend (2018) described this anole from the Lenca highlands of southwestern Honduras. Its EVS can be estimated as $5+7+3=15$, placing in the lower portion of the high vulnerability category. This species is limited to the eastern nuclear Central American highlands, and can be placed in conservation priority level one.

(9) *Rhadinella xerophila*. Ariano-Sánchez and Campbell (2018) described this dipsadid snake from dry forest and thorn scrub of the Valle del Motagua, Guatemala. Its EVS can be calculated as $6+8+2=16$, placing it in the middle portion of the high vulnerability category. This species is restricted to a single physiographic region, the Caribbean lowlands of eastern Guatemala and northern Honduras, and can be placed in conservation priority level one.

(10) The genera *Coloptychon* and *Gerrhonotus*. García-Vázquez et al. (2018), in a paper on the molecular systematics and historical biogeography of the anguid genus *Gerrhonotus*, left open a number of taxonomic questions about the alligator lizards, but at least seemed to conclude that the monotypic genus *Coloptychon* (and its species *rhombifer*) should be returned to the genus *Gerrhonotus*. The genus *Coloptychon* has had a history of wobbling between recognition as distinct or not from *Gerrhonotus* over the course of its 142-year history. In this addendum, we follow the decision of García-Vázquez et al. (2018).



Vicente Mata-Silva is a herpetologist originally from Río Grande, Oaxaca, Mexico. His interests include ecology, conservation, natural history, and biogeography of the herpetofaunas of Mexico, Central America, and the southwestern United States. He received his B.S. degree from the Universidad Nacional Autónoma de México (UNAM), and his M.S. and Ph.D. degrees from the University of Texas at El Paso (UTEP). Vicente is an Assistant Professor of Biological Sciences at UTEP in the Ecology and Evolutionary Biology Program, and Assistant Director of UTEP's 40,000 acre Indio Mountains Research Station, located in the Chihuahuan Desert of Trans-Pecos, Texas. To date, Vicente has authored or co-authored over 100 peer-reviewed scientific publications. He also was the Distribution Notes Section Editor for the journal *Mesoamerican Herpetology*.



Dominic L. DeSantis is currently a Ph.D. candidate and National Science Foundation-Graduate Research Fellow at the University of Texas at El Paso. He received his Bachelor's degree at Texas State University where he also completed multiple research projects on the antipredator behavior of the critically endangered Barton Springs Salamander (*Eurycea sosorum*). His ongoing dissertation research integrates multiple field monitoring technologies to study snake movement and behavioral ecology. Dominic accompanied Vicente Mata-Silva, Eli García-Padilla, and Larry David Wilson on survey and collecting trips to Oaxaca in 2015, 2016, and 2017 and is a co-author on numerous natural

history publications produced from those visits, including an invited book chapter entitled: "Conservation of Herpetofauna in Disturbed Habitats: Perspectives from Short-term Surveys in the Sierra Madre del Sur, Oaxaca, Mexico." Overall, Dominic has co-authored over 50 peer-reviewed scientific publications.



Eli García-Padilla is a herpetologist primarily focused on the study of the ecology and natural history of the Mexican herpetofauna. His research efforts have centered on the Mexican states of Baja California, Tamaulipas, Chiapas, and Oaxaca. His first experience in the field was researching the ecology of the insular endemic populations of the rattlesnakes *Crotalus catalinensis*, *C. muertensis* (now allocated to *C. pyrrhus*) and *C. tortugensis* (now allocated to *C. atrox*) in the Gulf of California. For his Bachelor's degree he presented a thesis on the ecology of *C. muertensis* (now allocated to *C. pyrrhus*) on Isla El Muerto, Baja California, Mexico. To date, he has authored or co-authored over 75 peer-reviewed scientific publications.

Currently, he is employed as a formal Curator of Amphibians and Reptiles from Mexico in the electronic platform "Naturalista" of the Comisión Nacional para el Uso y Conocimiento de la Biodiversidad (CONABIO; www.naturalista.mx). One of his main passions is environmental education, and for several years he has been working on a variety of projects that include the use of audiovisual media as a powerful tool to reach large audiences and to promote the importance of the knowledge, protection, and conservation of biodiversity in Mexico. Eli's interests include wildlife and conservation photography, and his art has been published in several recognized scientific, artistic, and educational books, magazines, and websites. Presently, he is collaborating on a research project evaluating the Jaguar (*Panthera onca*) as an umbrella species for the conservation of the herpetofauna of Nuclear Central America.



Jerry D. Johnson is Professor of Biological Sciences at The University of Texas at El Paso, and has extensive experience studying the herpetofauna of Mesoamerica, especially that of southern Mexico. Jerry is the Director of the 40,000-acre "Indio Mountains Research Station," was a co-editor on *Conservation of Mesoamerican Amphibians and Reptiles* and co-author of four of its chapters, co-editor of *Mesoamerican Herpetology: Systematics, Zoogeography, and Conservation*, and co-author of *Middle American Herpetology: A Bibliographic Checklist*. He also is the senior author of the recent paper "A conservation reassessment of the Central American herpetofauna based on the EVS measure" and is Mesoamerica/Caribbean editor for Geographic Distribution section of *Herpetological Review*. Johnson has authored or co-authored over 120 peer-reviewed papers, including other studies on the conservation status of the Mesoamerican herpetofauna. One species, *Tantilla johnsoni*, has been named in his honor. For several years, he was an Associate Editor and Co-chair of the Taxonomic Board for the journal *Mesoamerican Herpetology*.



Larry David Wilson is a herpetologist with lengthy experience in Mesoamerica. He was born in Taylorville, Illinois, United States, and received his university education at the University of Illinois at Champaign-Urbana (B.S. degree) and at Louisiana State University in Baton Rouge (M.S. and Ph.D. degrees). He has authored or co-authored over 400 peer-reviewed papers and books on herpetology, including numerous papers on the conservation status of Mesoamerica and its constituent parts. Larry is the senior editor of *Conservation of Mesoamerican Amphibians and Reptiles* and the co-author of seven of its 21 chapters. His other books include *The Snakes of Honduras*, *Middle American Herpetology*, *The Amphibians of Honduras*, *Amphibians & Reptiles of the Bay Islands and Cayos Cochinos, Honduras*, *The Amphibians and Reptiles of the Honduran Mosquitia*, and *Guide to the Amphibians & Reptiles of Cusuco National Park, Honduras*.

To date, he has authored or co-authored the descriptions of 71 currently recognized herpetofaunal species, and seven species have been named in his honor, including the anuran *Craugastor lauraster*, the lizard *Norops wilsoni*, and the snakes *Oxybelis wilsoni*, *Myriopholis wilsoni*, and *Cerrophidion wilsoni*. For several years, Larry was an Associate Editor and Co-chair of the Taxonomic Board for the journal *Mesoamerican Herpetology*.