



# Native anuran species as prey of invasive American Bullfrog, *Lithobates catesbeianus*, in Brazil: a review with new predation records

<sup>1,2,\*</sup>Fabrizio H. Oda, <sup>3</sup>Vinicius Guerra, <sup>4</sup>Eduardo Grou, <sup>5</sup>Lucas D. de Lima, <sup>6</sup>Helen C. Proença, <sup>6</sup>Priscilla G. Gambale, <sup>4,5</sup>Ricardo M. Takemoto, <sup>7</sup>Cauê P. Teixeira, <sup>7</sup>Karla M. Campião, and <sup>8</sup>Jean Carlo G. Ortega

<sup>1</sup>Departamento de Química Biológica, Programa de Pós-graduação em Bioprospecção Molecular, Universidade Regional do Cariri, Campus Pimenta, 63105-000, Crato, Ceará, BRAZIL <sup>2</sup>Departamento de Química Biológica, Laboratório de Zoologia, Universidade Regional do Cariri, Campus Pimenta, Crato, Ceará, BRAZIL <sup>3</sup>Departamento de Ecologia, Laboratório de Herpetologia e Comportamento Animal, Instituto de Ciências Biológicas, Universidade Federal de Goiás, Campus Samambaia, Goiânia, Goiás, BRAZIL <sup>4</sup>Centro de Ciências Biológicas, Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura, Laboratório de Ictioparasitologia, Universidade Estadual de Maringá, Maringá, Paraná, BRAZIL <sup>5</sup>Centro de Ciências Biológicas, Programa de Pós-graduação em Biologia Comparada, Universidade Estadual de Maringá, Paraná, BRAZIL <sup>6</sup>Universidade Estadual de Mato Grosso do Sul, Dourados, Mato Grosso do Sul, BRAZIL <sup>7</sup>Departamento de Zoologia, Laboratório de Ecologia de Interações Antagonistas, Universidade Federal do Paraná, Centro Politécnico, Curitiba, Paraná, BRAZIL <sup>8</sup>Departamento de Ecologia, Programa de Pós-graduação em Ecologia e Evolução, Instituto de Ciências Biológicas, Universidade Federal de Goiás, Campus Samambaia, Goiânia, Goiás, BRAZIL

**Abstract.**—The American Bullfrog (*Lithobates catesbeianus*) is widely distributed throughout the world as an invasive species, and causes negative impacts on the fauna resulting from its voracious predatory activity. This study documents two new predation reports and reviews the previous predation reports of the American Bullfrog on native Brazilian anurans. Twenty-one species of native anurans were recorded as American Bullfrog prey in Brazil. A positive correlation was found between the number of native anurans preyed on by American Bullfrog and the respective family or number of species per genus. Most of the prey species are small or medium-sized, and the results suggest that the generalist diet and intraguild predation may have favored the widespread establishment of the American Bullfrog.

**Keywords.** Amphibia, Atlantic Forest, biological invasion, conservation, global change, intraguild predation, exotic species

**Citation:** Oda FH, Guerra V, Grou E, de Lima LD, Proença HC, Gambale PG, Takemoto RM, Teixeira CP, Campião KM, Ortega JCG. 2019. Native anuran species as prey of invasive American Bullfrog, *Lithobates catesbeianus*, in Brazil: a review with new predation records. *Amphibian & Reptile Conservation* 13(2) [General Section]: 217–226 (e207).

**Copyright:** © 2019 Oda et al. This is an open access article distributed under the terms of the Creative Commons Attribution License [Attribution 4.0 International (CC BY 4.0): <https://creativecommons.org/licenses/by/4.0/>], which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website: [amphibian-reptile-conservation.org](http://amphibian-reptile-conservation.org).

**Received:** 8 February 2017; **Accepted:** 23 June 2019; **Published:** 20 December 2019

## Introduction

Biological invasions represent a major threat to natural ecosystems and their respective biodiversity, human health, and food security (IUCN 2012). In this context, the American Bullfrog, *Lithobates catesbeianus* (Shaw, 1802), is a globally widespread introduced species (Lowe et al. 2004). It is native in North America, occurring from eastern Canada and the central and eastern United States to northeastern Mexico (Quiroga et al. 2015). The introduction of *L. catesbeianus* in non-native

environments has direct (e.g., predation and competition) and indirect (e.g., parasites, disease introduction, and biotic homogenization) impacts on biodiversity (Batista 2002; Batista et al. 2015; Kiesecker and Blaustein 1998; Kraus 2009).

*Lithobates catesbeianus* is a voracious predator whose diet includes a wide variety of prey (Boelter and Cechin 2007; Boelter et al. 2012; Silva et al. 2009). Juveniles feed mainly on insects (Silva et al. 2009), whereas adults prey upon invertebrates and small vertebrates, such as fish, reptiles, birds, and mammals (Quiroga et al. 2015).

**Correspondence.** [fabrizio\\_oda@hotmail.com](mailto:fabrizio_oda@hotmail.com) (FHO), [vinicius.guerrabatista@gmail.com](mailto:vinicius.guerrabatista@gmail.com) (VG), [eduardogrou@hotmail.com](mailto:eduardogrou@hotmail.com) (EG), [lucasduartelima@hotmail.com](mailto:lucasduartelima@hotmail.com) (LDL), [helencassia23@hotmail.com](mailto:helencassia23@hotmail.com) (HCP), [priscillagambale@gmail.com](mailto:priscillagambale@gmail.com) (PGG), [takemotorm@nupelia.uem.br](mailto:takemotorm@nupelia.uem.br) (RMT), [caue.cpt@gmail.com](mailto:caue.cpt@gmail.com) (CPT), [karla\\_mcamp@yahoo.com.br](mailto:karla_mcamp@yahoo.com.br) (KMC), [ortegajejan@gmail.com](mailto:ortegajejan@gmail.com) (JCGO)

American Bullfrogs are considered opportunistic feeders, also preying on amphibians, including conspecifics and other species (Silva et al. 2011; Toledo et al. 2007).

The American Bullfrog is now established in nearly 40 countries around the world (Frost 2019; Kraus 2009). In Brazil, the first specimens were introduced in 1935 for commercial exploitation at the municipality of Itaguaí, Rio de Janeiro state (Vizotto 1984). The introduction of the American Bullfrog for commercial frog farming was due to its fast reproduction and greater development in captivity compared to native species. It occurs mainly in the southern and southeastern Brazilian states because of its easy adaptation to the climatic conditions (Vizotto 1984). Approximately 2,000 commercial frog farms were active in the early 1990s in Brazil, but many closed their activities because of low profitability (Lima and Agostinho 1988), which led to American Bullfrog specimens being abandoned or released into the natural environments, and consequently several accidental invasions have occurred in Brazil (Both et al. 2011).

Populations of *Lithobates catesbeianus* are now known to be present in 155 Brazilian municipalities (Both et al. 2011; Instituto Horus 2016), a context in which many studies have revealed the localized impacts of its predatory activity on native anuran fauna (Batista et al. 2015; Boelter and Cechin 2007; Boelter et al. 2012; Leivas et al. 2012; Silva et al. 2011). In addition, global-scale studies have demonstrated trophic niche-width shifts in bullfrog populations from both native and invaded areas (Bissattini and Vignoli 2017), as well as the effects of the interactions between bullfrogs and crayfish on native amphibians (Bissattini et al. 2018, 2019; Liu et al. 2018). However, studies summarizing data on the predation of native anurans by American Bullfrogs have yet to be presented; therefore, knowledge on the impact and the native anuran species preyed upon by such an invasive frog may benefit our understanding of their predator-prey relationships.

Herein, the predation of *Boana raniceps* and *Phyllomedusa distincta* by males of *Lithobates catesbeianus* are reported, and the available literature on the predation of native anurans by the invasive frog *L. catesbeianus* in Brazil is reviewed. An overview on the number and identities of native species reported as prey and the potential impact of the American Bullfrog on native anurans are provided.

## Material and Methods

### Bibliographic Review

An extensive literature review was conducted to find scientific articles, natural history notes, and theses which contain reports on the predation of native anurans by the

invasive American Bullfrog *Lithobates catesbeianus* in Brazil. The sources included articles or natural history notes published in *Herpetological Review* (1967–2018), *Herpetological Bulletin* (2008–2018), *Herpetology Notes* (2008–2018), and *South American Journal of Herpetology* (2006–2018). Searches were also conducted in Web of Science using the following query: (“*Rana catesbeiana*” OR “*Lithobates catesbeianus*”) AND (“diet” OR “feeding biology” OR “predation”), applied in the field “topic” on 30 December 2018, without applying any filters for year or other parameters. Considering that predation attempts would not necessarily result in a predation event (Toledo et al. 2007), reports of predation attempts in the field, laboratory experiments, or captivity were not included. Masters and doctoral papers in digital format were obtained from the library databases of Brazilian universities (especially Universidade Estadual Paulista and Universidade Regional de Blumenau) by using the search terms mentioned above in the Google search engine.

The Web of Science query resulted in 159 studies, three of which met the criteria and were included in the study. Eight additional predation records were selected for inclusion in the study by searching the selected journals (six studies) and the library databases of Brazilian universities (two studies). Information was extracted from each diet analysis (i.e., the diet was described through the analyses of stomach contents or predation records), study location, anuran prey species, geographic range, and body size. The geographic range follows the list of anuran species for each Brazilian federal state and the biomes proposed in Toledo and Batista (2012). The body sizes of anuran species follow the size values available in Uetanabaro et al. (2008) and Haddad et al. (2013). The spatial distribution map of *Lithobates catesbeianus* invasive populations and predation reports were generated with 155 occurrence points for American Bullfrog in Brazil, obtained from Both et al. (2011) and Instituto Horus (2016).

### Data Analysis

The relation between the number of native anuran species preyed upon by the American Bullfrog and the number of native anurans per family or genus was tested with a Pearson correlation analysis. The numbers of native anuran prey species per family and genus were compiled following the Frost (2019) database. Toledo et al. (2007) stated that a positive correlation between the number of predation events and taxonomic richness may be a proxy for search representativeness, by reasoning that taxa with more species would be more frequently preyed by chance (i.e., a sampling effect). Such a correlation could indicate the possible mechanisms of *L. catesbeianus* impacts on native biota apart from search representativeness.



**Fig. 1.** Adult *Lithobates catesbeianus* swallowing an adult *Boana raniceps* in an artificial permanent pond within pasture area in southern Brazil.

## Results

### Field Observations and New Predation Records

An adult *Lithobates catesbeianus* swallowing an adult *Boana raniceps* (Fig. 1) was recorded on 11 October 2014 at 2100 h, in an artificial permanent pond inside a pasture area (23°20'38"S, 51°52'07"W), in the northern region of Paraná state, southern Brazil. Although the specimens escaped, voucher specimens of the native anuran species and *L. catesbeianus* had been previously collected by Affonso et al. (2014) and stored at the Amphibian Collection from the Zoology and Botany Department, Bioscience Institute, Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil.

A second predation event recorded a male adult specimen of Bullfrog swallowing a treefrog (Fig. 2A). The specimen was collected during an *L. catesbeianus* survey on 22 January 2019, at 2200 h, in an artificial permanent pond in a rural property at Iporanga, southern São Paulo state, southeastern Brazil (24°35'01.2"S, 48°36'00.4"W). The *L. catesbeianus* specimen was taken to the laboratory where the anuran prey was removed and identified as an adult *Phyllomedusa distincta* (Fig. 2B).



**Fig. 2.** (A) Predation of an adult *Phyllomedusa distincta* by *Lithobates catesbeianus*, (B) Adult *P. distincta* partially digested, removed from the oral cavity of *L. catesbeianus*.

### Exploratory Analysis


Overall, 11 publications reported predation events, corresponding to 41 records of native anurans as prey of *L. catesbeianus* (Table 1). Nine of the publications discussed the diet in a broader sense, and two were natural history notes reporting predation events. Most of the records occurred in Minas Gerais state (39%), followed by Rio Grande do Sul (~32%), Paraná (~12%), São Paulo (~12%), and Santa Catarina (~5%), at sites inside the Atlantic Forest, in addition to another site in a transition zone between Cerrado and Atlantic Forest (Fig. 3, Table 1).

This survey accounted for 21 anuran species as prey of *L. catesbeianus*, all widely distributed and possibly coexisting with American Bullfrogs in their breeding sites. The anuran family Hylidae had the highest number of species (11 species), followed by Leptodactylidae (four species), Bufonidae and Microhylidae with two species each, and Odontophrynidae and Phyllomedusidae with one species each. *Lithobates catesbeianus* often preyed on medium-sized species, but small-sized species were also preyed upon (Table 1).

A positive correlation was found between the number of native anuran species preyed on by American Bullfrog and genus richness ( $r = 0.71$ ,  $P = 0.01$ ), whereas at the family level no relationship was found ( $r = 0.53$ ,  $P = 0.22$ ). Thus, considering the studies analyzed, genera with higher numbers of species presented more potential prey for American Bullfrogs in Brazil.

### Discussion

Most of the predation records found in this review came from a few studies which assessed the overall dietary composition of *L. catesbeianus*, and revealed that the diet of these invasive frog populations is represented by a wide variety of native anuran species (Boelter and Cechin 2007; Silva et al. 2009, 2010, 2011). Only two predation records of *L. catesbeianus* and native anurans in the field were found, probably due to some difficulty



**Fig. 3.** Spatial distribution of *Lithobates catesbeianus* invasive populations and predation reports of native anurans in Brazil. White circles: American Bullfrog populations in Brazil (Both et al. 2011; Instituto Horus 2016); yellow stars: predation reports of adult *Boana raniceps* and adult *Phyllomedusa distincta* in southern and southeastern Brazil; light green circles: locations of 41 published predation records.

in recording and quantifying these events in the field (Pombal Jr. 2007).

The predicted potential occurrence of *L. catesbeianus* in Brazil represents its current distribution in the southern and southeastern regions in the Atlantic Forest, with potential areas for colonization remaining in the central and northeastern regions (Giovanelli et al. 2008; Both et al. 2011). The results showed that all predation records occurred at sites in southern and southeastern Brazil, regions with higher numbers of research centers, thus contributing a disproportionately greater number of field studies.

Native anurans recorded as prey of American Bullfrogs share the same breeding sites. Silva et al. (2011) had found a spatial overlap in microhabitat use between native species and American Bullfrogs during the reproductive season. American Bullfrogs may also overlap with native amphibians in diet composition

(Bissattini et al. 2019). This may lead to a potential competition, and may have a direct influence on community composition patterns since the intrinsic ecological properties of organisms determine the niche overlap between species in the communities (Vignoli and Luiselli 2012; Vignoli et al. 2017). Additionally, the predation on other anuran species by *L. catesbeianus* can represent an example of intraguild predation (Polis et al. 1989), a process that may facilitate the establishment of the American Bullfrog (Bissattini et al. 2018), as found in other disparate introduced taxa, such as ladybird beetles (Snyder et al. 2004) and fish (Pereira et al. 2015). Intraguild predation can benefit the establishment of *L. catesbeianus* by reducing the competitive pressure by direct predation of the other anuran species.

The number of prey species had a positive correlation with the number of species per genus, in which the family Hylidae had the highest number of species as

**Table 1.** Native anuran species as prey of *Lithobates catesbeianus* in Brazil based on 41 records obtained from 11 publications, and from two predation events reported here from southern and southeastern Brazil.

| Anuran prey species   | Body Size <sup>a</sup> (cm) |       | Locality <sup>b</sup>  | Biomes <sup>c</sup>       | Reference   |
|---|-----------------------------|-------|--|---------------------------|---|
|   | SVL ♂                       | SVL ♀ |  |                           |   |
| <b>Bufonidae</b>  |                             |       |  |                           |   |
| <i>Rhinella abei</i> (Baldissera-Jr, Caramaschi and Haddad, 2004) | 6.1                         | 9.3   | Quatro Barras, Campina Grande do Sul, and Bocaíva do Sul, PR                       | AF                        | Leivas et al. 2012  |
| <i>Rhinella ornata</i> (Spix, 1824)                               | 6.4                         | 7.9   | Viçosa, MG<br>Viçosa, MG   | AF, Ce                    | Reis et al. 2007<br>Silva et al. 2010   |
| <b>Hylidae</b>  |                             |       |  |                           |   |
| <i>Dendropsophus elegans</i> (Wied-Neuwied, 1824)                 | 2.9                         | 3.1   | Viçosa, MG<br>Viçosa, MG<br>Viçosa, MG   | AF, Ca                    | Silva et al. 2009<br>Silva et al. 2011<br>Silva et al. 2016                         |
| <i>Dendropsophus minutus</i> (Peters, 1872)                       | 2.3                         | 2.5   | Agudo and Nova Palma, RS<br>Agudo and Nova Palma, RS<br>Viçosa, MG<br>Botucatu, SP | AF, Am, Ca, Ce,<br>Pt, Pp | Boelter and Cechin 2007<br>Boelter et al. 2012<br>Silva et al. 2009<br>Almeida 2010 |
| <i>Dendropsophus nanus</i> (Boulenger, 1889)                      | 2.0                         | 2.3   | Botucatu, SP   | AF, Am, Ca, Ce,<br>Pt, Pp | Almeida 2010  |
| <i>Boana albomarginata</i> (Spix, 1824)                           | 5.1                         | 5.8   | Blumenau, SC   | AF, Ca, Ce                | Dallacorte 2010   |
| <i>Boana bischoffi</i> (Boulenger, 1887)                          | 4.5                         | 5.9   | Quatro Barras, Campina Grande do Sul, and Bocaíva do Sul, PR                       | AF, Ce, Pp                | Leivas et al. 2012  |
| <i>Boana faber</i> (Wied-Neuwied, 1821)                           | 8.8                         | 8.9   | Viçosa, MG<br>Blumenau, SC   | AF, Ca, ce                | Silva et al. 2009<br>Dallacorte 2010  |
| <i>Boana pulchella</i> (Duméril and Bibron, 1841)                 | 4.2                         | 4.5   | Agudo and Nova Palma, RS<br>Agudo and Nova Palma, RS                               | AF, Pp                    | Boelter and Cechin 2007<br>Boelter et al. 2012                                      |
| <i>Boana raniceps</i> Cope, 1862                                  | 7.1                         | 6.8   | Maringá, PR  | AF, Am, Ca,<br>Ce, Pt     | This study  |
| <i>Scinax crosopodophilus</i> (A. Lutz, 1925)                     | 3.0                         | 3.1   | Viçosa, MG   | AF                        | Silva et al. 2009   |
| <i>Scinax eurydice</i> (Bokermann, 1968)                          | 5.3                         | 4.8   | Viçosa, MG<br>Agudo and Nova Palma, RS<br>Agudo and Nova Palma, RS                 | AF, Am, Ca, Ce            | Silva et al. 2009<br>Boelter and Cechin 2007<br>Boelter et al. 2012                 |
| <i>Scinax fuscovarius</i> (A. Lutz, 1925)                         | 4.2                         | 4.5   | Botucatu, SP<br>Viçosa, MG<br>Vieiras, MG<br>Vieiras, MG                           | AF                        | Almeida 2010<br>Silva et al. 2010<br>Silva et al. 2011<br>Silva et al. 2016         |

**Table 1 (continued).** Native anuran species as prey of *Lithobates catesbeianus* in Brazil based on 41 records obtained from 11 publications, and from two predation events reported here from southern and southeastern Brazil.

| Anuran prey species  | Body Size <sup>a</sup> (cm) |       | Locality <sup>b</sup>   | Biomes <sup>c</sup>       | Reference  |
|--|-----------------------------|-------|---|---------------------------|--|
|  | SVL ♂                       | SVL ♀ |   |                           |  |
| <b>Leptodactylidae</b>                                     |                             |       |   |                           |  |
| <i>Leptodactylus fuscus</i> (Schneider, 1799)              | 4.4                         | 4.4   | Agudo, and Nova Palma, RS<br>Agudo, and Nova Palma, RS                                      | AF, Am, Ca, Ce,<br>Pp, Pt | Boelter and Cechin 2007<br>Boelter et al. 2012               |
| <i>Leptodactylus latrans</i> (Steffen, 1815)               | 9.6                         | 9.2   | Quatro Barras, Campina Grande do Sul, and Bociuíva do Sul, PR<br>Agudo and Nova Palma, RS   | AF, Am, Ca, Ce,<br>Pp, Pt | Leivas et al. 2012<br>Boelter and Cechin 2007                |
| <i>Physalaemus cuvieri</i> Fitzinger, 1826                 | 2.8                         | 3.1   | Agudo and Nova Palma, RS<br>Botucatu, SP  | AF, Am, Ca, Ce,<br>Pp, Pt | Boelter et al. 2012<br>Almeida 2010                          |
| <i>Pseudopaludicola mystacalis</i> (Cope, 1887)            | 1.7                         | 2.1   | Quatro Barras, Campina Grande do Sul, and Bociuíva do Sul, PR<br>Vieiras, MG<br>Vieiras, MG | AF, Am, Ca, Ce,<br>Pp, Pt | Leivas et al. 2012<br>Silva et al. 2011<br>Silva et al. 2016 |
| <b>Microhylidae</b>  |                             |       |   |                           |  |
| <i>Elachistocleis cesarii</i> (Miranda-Ribeiro, 1920)      | 2.9                         | 3.2   | Vieiras, MG<br>Vieiras, MG  | AF, Ca, Ce                | Silva et al. 2011<br>Silva et al. 2016                       |
| <i>Elachistocleis bicolor</i> (Guérin-Ménéville, 1838)     | 2.8                         | 3.8   | Agudo and Nova Palma, RS  | AF, Ce, Pp                | Boelter et al. 2012  |
| <b>Odontophrynidae</b>                                     |                             |       |   |                           |  |
| <i>Odontophrynus americanus</i> (Duméril and Bibron, 1841) | 4.5                         | 4.9   | Agudo and Nova Palma, RS<br>Passo Fundo, RS   | AF, Ca, Ce, Pp            | Boelter and Cechin 2007<br>Leivas et al. 2013                |
| <b>Phyllomedusidae</b>                                     |                             |       |   |                           |  |
| <i>Phyllomedusa distincta</i> Lutz, 1950                   | 5.6                         | 6.0   | Iporanga, SP  | AF, Ce, Pp                | This study   |

<sup>a</sup> Values of snout-vent length (SVL) available in Uetanabaro et al. (2008) and Haddad et al. (2013).<sup>b</sup> Brazilian states: MG = Minas Gerais; PR = Paraná; RS = Rio Grande do Sul; SC = Santa Catarina; SP = São Paulo.<sup>c</sup> Occurrence of anuran species within Brazilian biomes: AF = Atlantic Forest; Am = Amazon; Ca = Caatinga; Ce = Cerrado; Pp = Pampa; Pt = Pantanal.

prey of *L. catesbeianus*. Boelter et al. (2012) found that 60% of the prey records corresponded to Hylidae species, suggesting that this group suffers higher predation pressure. At least two non-mutually exclusive hypotheses may explain these patterns by relating family (i.e., richness, abundance) and species traits (i.e., body size) to *L. catesbeianus* predation rates, involving species traits that are often phylogenetically correlated (Martins and Hansen 1997). Firstly, it is possible that Hylidae species are often preyed upon due to their higher species richness in comparison to other families, which is a plausible hypothesis if we assume that predation rates can be proportional to prey abundance or richness (i.e., higher predation rates in higher resource availability conditions; Jacobsen et al. 2014; Madahi et al. 2015). Secondly, Hylidae species may have a higher predation rate because of their smaller size relative to species from other families (e.g., Bufonidae). Predators that feed on whole animals, such as the American Bullfrog, are limited by the prey's body size. Experimental evidence indicates that larger specimens of *L. catesbeianus* feed preferentially on smaller, rather than on large-sized, native anurans (Wang et al. 2007), suggesting a size-based selection of prey species. Partially related to this hypothesis, our results suggest a higher amount of small and medium-sized species as American Bullfrog prey. Therefore, the preference for prey of a certain body size may be proportionally related to the body size of the predators, as observed in previous studies (Quiroga et al. 2015; Silva et al. 2011, 2009; Wang et al. 2007).

This survey found that all anuran species preyed upon by *L. catesbeianus* have large geographic distributions, occurring in various Brazilian states and biomes (Frost 2019; Toledo and Batista 2012). Both et al. (2014) found that American Bullfrog abundance had a positive relationship with communities that consisted of generalist species (e.g., *Physalaemus cuvieri*, *Dendropsophus minutus*), that were anthropogenically adapted and broadly distributed in South America. Native anurans with large geographic ranges (e.g., *Rhinella diptycha*, *Dendropsophus minutus*, *Boana faber*, *B. raniceps*, *Scinax fuscovarius*, and *Physalaemus cuvieri*) have also been found in sympatry with *Lithobates catesbeianus* elsewhere (Affonso et al. 2014).

## Conclusions

This study indicated *L. catesbeianus* preys on at least 21 native anuran species in Brazil. Predation is one of the major negative effects of invasive species on native communities. The quality of being a generalist feeder, preying on many anuran species, has benefited the successful colonization, establishment, and permanence of the American Bullfrog in Brazil (and even worldwide; e.g., Li et al. 2011; Monello et al. 2006; Quiroga et al. 2015). Native species of the family Hylidae may be more susceptible to American Bullfrog predation because of

their higher abundance and richness, and/or due to a higher representation of small- to medium-sized species relative to other anuran families. Knowledge on the species most vulnerable to predation by the American Bullfrog can enable better prediction of the negative impacts of such an invasive species on native anuran communities.

**Acknowledgements.**—The authors would like to thank Centro Universitário de Maringá for having granted us access to the Fazenda Escola UniCesumar, and Bruno Barreto for assisting with Fig. 3. Fabrício H. Oda received a postdoctoral fellowship from Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico/Coordenação de Aperfeiçoamento Pessoal de Nível Superior – CAPES (Grant number 88887.162751/2018-00). Vinicius Guerra received fellowships from CAPES, and Helen C. Proença and Jean Carlo G. Ortega received fellowships from Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq. Ricardo M. Takemoto received a CNPq grant of research productivity fellowship as well. Finally, the authors would like to thank the Instituto Chico Mendes de Conservação da Biodiversidade/Sistema de Autorização e Informação em Biodiversidade for providing us with the collecting permit (process #23866-1).

## Literature Cited

- Almeida SC. 2010. Ecologia de *Lithobates catesbeianus* (Shaw, 1801) e relações com os anfíbios da região de Botucatu, SP (Amphibia, Anura). Ph.D. Dissertation, Universidade Estadual Paulista, Botucatu, SP, Brazil. 79 p.
- Affonso IP, Cafofo EG, Delariva RL, Oda FH, Karling LK, Lourenço-de-Moraes R. 2014. List of anurans (Amphibia: Anura) from the rural zone of the municipality of Maringá, Paraná state, southern Brazil. *Check List* 10(4): 878–882.
- Batista CG. 2002. *Rana catesbeiana* (Bullfrog). Effects on native anuran community. *Herpetological Review* 33(2): 131.
- Batista M, Silva M, Barreto C. 2015. Effects of introduction and decline of a bullfrog population (*Lithobates catesbeianus*) in a community of amphibians in the Cerrado from Central Brazil. *Herpetology Notes* 8: 263–265.
- Bissattini A, Vignoli L. 2017. Let's eat out, there's crayfish for dinner: American Bullfrog niche shifts inside and outside native ranges and the effect of introduced crayfish. *Biological Invasions* 19(9): 2,633–2,646.
- Bissattini A, Buono V, Vignoli L. 2018. Field data and worldwide literature review reveal that alien crayfish mitigate the predation impact of the American Bullfrog on native amphibians. *Aquatic Conservation: Marine and Freshwater Ecosystems* 28(6): 1,465–1,475.

- Bissattini A, Buono V, Vignoli L. 2019. Disentangling the trophic interactions between American Bullfrogs and native anurans: complications due to post-metamorphic ontogenetic niche shifts. *Aquatic Conservation* 29: 270–281.
- Boelter RA, Cechin SZ. 2007. Impact of the bullfrog diet *Lithobates catesbeianus* (Anura, Ranidae) on native fauna: case study from the region of Agudo, RS, Brazil. *Natureza & Conservação* 5(2): 115–123.
- Boelter RA, Kaefer IL, Both C, Cechin S. 2012. Invasive bullfrogs as predators in a Neotropical assemblage: What frog species do they eat? *Animal Biology* 62(4): 397–408.
- Both C, Lingnau R, Santos Jr A, Madalozzo B, Lima LP, Grant T. 2011. Widespread occurrence of the American Bullfrog, *Lithobates catesbeianus* (Shaw, 1802) (Anura: Ranidae), in Brazil. *South American Journal of Herpetology* 6(2): 127–134.
- Both C, Madalozzo B, Lingnau R, Grant T. 2014. Amphibian richness patterns in Atlantic Forest areas invaded by American Bullfrogs. *Austral Ecology* 39(7): 864–874.
- Dallacorte F. 2010. Impacto da Rã-touro-gigante (*Lithobates catesbeianus*) sobre a fauna nativa na zona de amortecimento e interior do Parque Nacional da Serra do Itajaí (PNSI), Blumenau – SC. M.S. Thesis, Universidade Regional de Blumenau, Blumenau, SC, Brazil. 106 p.
- Ficetola GF, Thuller W, Miaud C. 2007. Prediction and validation of the potential global distribution of a problematic alien invasive species – the American Bullfrog. *Diversity and Distributions* 13(4): 476–485.
- Frost DR. 2019. *Amphibian Species of the World: an Online Reference*. Version 6.0. Available: <http://research.amnh.org/herpetology/amphibia/index.html> [Accessed: 12 February 2019].
- Giovanelli JGR, Haddad CFB, Alexandrino J. 2008. Predicting the potential distribution of the alien invasive American Bullfrog (*Lithobates catesbeianus*) in Brazil. *Biological Invasions* 10(5): 585–590.
- Haddad CFB, Toledo LF, Prado CPA, Loebmann D, Gasparini JL, Sazima I. 2013. *Guia dos Anfíbios da Mata Atlântica – Diversidade e Biologia*. Anolis Books, São Paulo, Brazil. 544 p.
- Instituto Hórus. 2016. Base de dados de espécies exóticas invasoras no Brasil. Instituto Hórus, Florianópolis, SC, Brazil. Available: <http://www.institutohorus.org.br> [Accessed: 20 October 2016].
- IUCN. 2012. *Biological Invasions: a Growing Threat to Biodiversity, Human Health and Food Security. Policy Recommendation for the Rio+20 Process Drafted by IUCN Species Survival Commission Invasive Species Specialist Group and Invasive Species Initiative*. International Union for Conservation of Nature (IUCN), Gland, Switzerland. Available: [http://cmsdata.iucn.org/downloads/policy\\_brief\\_on\\_invasive\\_and\\_alien\\_species\\_3.pdf/](http://cmsdata.iucn.org/downloads/policy_brief_on_invasive_and_alien_species_3.pdf/) [Accessed: 13 October 2016].
- Jacobsen L, Berg S, Baktoft H, Nilsson PA, Skov C. 2014. The effect of turbidity and prey fish density on consumption rates of piscivorous Eurasian perch *Perca fluviatilis*. *Journal of Limnology* 73(1): 187–190.
- Kiesecker JM, Blaustein AR. 1998. Effects of introduced bullfrogs and small-mouth bass on the microhabitat use, growth, and survival of native red-legged frogs. *Conservation Biology* 12(4): 776–787.
- Kraus F. 2009. *Alien Reptiles and Amphibians: a Scientific Compendium and Analysis*. Springer, New York, New York, USA. 563 p.
- Leivas PT, Leivas FWT, Moura MO. 2012. Diet and trophic niche of *Lithobates catesbeianus* (Amphibia: Anura). *Zoologia* 29(5): 405–412.
- Leivas PT, Savaris M, Lampert S, Lucas EM. 2013. Predation of *Odontophrynus americanus* (Anura: Odontophrynidae) by the invasive species *Lithobates catesbeianus* (Anura: Ranidae) in an Araucaria Forest remnant in Southern Brazil. *Herpetology Notes* 6: 603–606.
- Li YM, Ke ZW, Wang YH, Blackburn TM. 2011. Frog community responses to recent American Bullfrog invasions. *Current Zoology* 57(1): 83–92.
- Lima SL, Agostinho CA. 1988. *A Criação de Rãs*. Editora Globo, Rio de Janeiro, Brazil. 187 p.
- Liu X, Wang S, Ke Z, Cheng C, Wang Y, Zhang F, Xu F, Li X, Gao X, Jin C, et al. 2018. The invaders do not result in heavier impacts: the effects of non-native bullfrogs on native anurans are mitigated by high densities of non-native crayfish. *Journal of Animal Ecology* 87(3): 850–862.
- Lowe S, Browne M, Boudjelas S, De Poorter M. 2004. *100 of the World's Worst Invasive Alien Species: a Selection from the Global Invasive Species Database*. The Invasive Species Specialist Group (ISSG), a specialist group of the Species Survival Commission (SSC) of the World Conservation Union (IUCN), Gland, Switzerland. Available: [http://www.issg.org/database/species/reference\\_files/100english.pdf/](http://www.issg.org/database/species/reference_files/100english.pdf/) [Accessed: 13 October 2016].
- Loyola RD, Nabout JC, Trindade-Filho J, Lemes P, Urbina-Cardona JN, Dobrovolski R, Sagnori MD, Diniz-Filho JAF. 2012. Climate change might drive species into reserves: a case study of the American Bullfrog in the Atlantic Forest biodiversity hotspot. *Alytes* 29(1-4): 61–74.
- Madahi K, Sahragard A, Hosseini R. 2015. Predation rate and numerical response of *Aphidoletes aphidimyza* feeding on different densities of *Aphis craccivora*. *Biocontrol Science and Technology* 25(1): 72–83.
- Martins EP, Hansen TF. 1997. Phylogenies and the comparative method: A general approach to incorporating phylogenetic information into the analysis of interspecific data. *American Naturalist* 149(4): 646–667.



- Monello RJ, Dennehy JJ, Murray DL, Wirsing AJ. 2006. Growth and behavioral responses of tadpoles of two native frogs to an exotic competitor, *Rana catesbeiana*. *Journal of Herpetology* 40(3): 403–407.
- Nori J, Urbina-Cardona JN, Loyola RD, Lescano JN, Leynaud GC. 2011. Climate change and American Bullfrog invasion: What could we expect in South America? *PLoS ONE* 60: e25718.
- Pereira LS, Agostinho AA, Gomes LC. 2015. Eating the competitor: a mechanism of invasion. *Hydrobiologia* 746(1): 223–231.
- Polis GA, Myers CA, Holt RD. 1989. The ecology and evolution of intraguild predation: potential competitors that eat each other. *Annual Review of Ecology, Evolution, and Systematics* 20: 297–330.
- Pombal Jr JP. 2007. Notas sobre predação em uma taxocenose de anfíbios anuros no sudeste do Brasil. *Revista Brasileira de Zoologia* 24(3): 841–843.
- Quiroga L, Moreno MD, Cataldo AA, Aragón-Traverso JH, Pantano MV, Olivares JPS, Sanabria EA. 2015. Diet composition of an invasive population of *Lithobates catesbeianus* (American Bullfrog) from Argentina. *Journal of Natural History* 49(24–28): 1,703–1,716.
- Reis EP, Silva ET, Feio RN, Filho OPR. 2007. *Chaunus pombali* (Pombali's toad). Predation. *Herpetological Review* 38(3): 321.
- Silva ET, Both C, Filho OPR. 2016. Food habits of invasive bullfrogs and native thin-toed frogs occurring in sympatry in southeastern Brazil. *South American Journal of Herpetology* 11(1): 25–33.
- Silva ET, Ribeiro-Filho OP, Feio RN. 2011. Predation of native anurans by invasive bullfrogs in southeastern Brazil: spatial variation and effect of microhabitat use by prey. *South American Journal of Herpetology* 6(1): 1–10.
- Silva ET, Reis EP, Feio RE, Filho OPR. 2009. Diet of the invasive frog *Lithobates catesbeianus* (Shaw, 1802) (Anura: Ranidae) in Viçosa, Minas Gerais state, Brazil. *South American Journal of Herpetology* 4(3): 286–294.
- Silva ET, Reis EP, Santos PS, Feio RN. 2010. *Lithobates catesbeianus* (American Bullfrog). Diet. *Herpetological Review* 41(4): 475–476.
- Snyder WE, Clevenger GM, Eigenbrode SD. 2004. Intraguild predation and successful invasion by introduced ladybird beetles. *Oecologia* 140(4): 559–565.
- Toledo LF, Batista RF. 2012. Integrative study of Brazilian anurans: geographic distribution, size, environment, taxonomy, and conservation. *Biotropica* 44(6): 785–792.
- Toledo LF, Ribeiro RS, Haddad CFB. 2007. Anurans as prey: an exploratory analysis and size relationships between predators and their prey. *Journal of Zoology* 271(2): 170–177.
- Uetanabaro M, Prado CPA, Rodrigues DJ, Gordo M, Campos Z. 2008. *Guia de Campo dos Anuros do Pantanal Sul e Planaltos de Entorno*. Editora UFMS/UFMT, Campo Grande, MS, Brazil. 196 p.
- Vizotto LD. 1984. Ranicultura. *Ciência e Cultura* 36(1): 42–45.
- Wang YP, Guo ZW, Pearl CA, Li YM. 2007. Body size affects the predatory interactions between introduced American Bullfrogs (*Rana catesbeiana*) and native anurans in China: an experimental study. *Journal of Herpetology* 41(3): 514–520.



**Fabrício Hiroiuki Oda** is a postdoctoral research fellow in the Programa de Pós-graduação em Bioprospecção Molecular (URCA – Universidade Regional do Cariri, Brazil) and research collaborator in the Laboratório de Ictioparasitologia do Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura (UEM – Universidade Estadual de Maringá, Brazil). His primary areas of interest are the natural history, ecology, and parasitology of fish, amphibians, and reptiles.



**Vinicius Guerra** is a postdoctoral researcher in the Laboratório de Herpetologia e Comportamento Animal (UFG – Universidade Federal de Goiás, Brazil). His primary areas of interest are the study of community ecology, animal behavior, natural history, and bioacoustics, with a particular focus on amphibians.



**Eduardo Grou** is a volunteer researcher in the Laboratório de Ictioparasitologia do Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura (UEM – Universidade Estadual de Maringá, Brazil). His primary areas of interest are the ecology and parasitology of chelonians.

## Impact of bullfrogs on native anurans in Brazil



**Lucas Duarte de Lima** is a Master's student fellow in the Programa de Pós-graduação em Biologia Comparada (UEM – Universidade Estadual de Maringá, Brazil). His primary area of interest is amphibian diversity.



**Helen Cássia Proença** is a Doctoral student fellow in the Programa de Pós-graduação em Biologia Comparada (UEM – Universidade Estadual de Maringá, Brazil). Her primary area of interest is snake diversity.



**Priscilla Guedes Gambale** is a researcher in the Departamento de Biologia (UEMS – Universidade Estadual de Mato Grosso do Sul, Brazil). Her primary area of interest is the ecology of amphibians, with a particular focus on bioacoustics.



**Ricardo Massato Takemoto** is a researcher in the Laboratório de Ictioparasitologia do Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura (UEM – Universidade Estadual de Maringá, Brazil) and advisor in the Programa de Pós-graduação em Ecologia de Ambientes Aquáticos Continentais and the Programa de Pós-graduação em Biologia Comparada (UEM – Universidade Estadual de Maringá, Brazil). His primary area of interest is the study of parasites of aquatic organisms, including fish and amphibians.



**Cauê Pinheiro Teixeira** is Master's researcher in the Laboratório de Ecologia de Interações Antagonistas (UFPR – Universidade Federal do Paraná, Brazil). His primary areas of interest are ecology, herpetology, parasitology, and biological invasion.



**Karla Magalhães Campião** is a researcher and coordinator of the Laboratório de Ecologia de Interações Antagonistas (UFPR – Universidade Federal do Paraná, Brazil), and she supervises graduate students in Ecology and Zoology. Her primary areas of interest are amphibian parasites and disease ecology.



**Jean Carlo Gonçalves Ortega** is a postdoctoral researcher in the Programa de Pós-graduação em Ecologia e Evolução (UFG – Universidade Federal de Goiás, Brazil). His primary areas of interest are community ecology and biological invasions.