



Biology of snakes of the genus *Tretanorhinus*: an integrative review

^{1,2,*}Marco D. Barquero and ^{1,3}Viviana Arguedas

¹Asociación para la Conservación y el Estudio de la Biodiversidad (ACEBIO), San José, COSTA RICA ²Sede del Caribe, Universidad de Costa Rica, Montes de Oca, San José, 2060, COSTA RICA ³Sede de Occidente, Universidad de Costa Rica, Montes de Oca, San José, 2060, COSTA RICA

Abstract.—Many aspects of the biology of various snake species remain unknown, and the extent of this lack of information is not always clear. As new research usually depends upon previous findings, the gaps in our knowledge and the accuracy of published information are of major importance. Therefore, an analysis of all available information on snakes of the genus *Tretanorhinus*, from both the literature and museum specimens, is presented here to illuminate existing knowledge gaps. The database compiled from 87 documents referring to snakes of this genus and 755 specimens held in scientific collections revealed major gaps and contradictory information for all four species of this genus. Data on morphology, ecology, and natural history are completely absent for *T. mocquardi* and *T. taeniatus*, whereas confusing distribution reports exist for *T. nigroluteus*. The potential consequences of these problems were determined, and some suggestions for correcting them are addressed. Specifically, we consider that focused efforts on the validation of current species and subspecies, field and lab studies of ecology and behavior, and estimations of population dynamics, are necessary.

Keywords. Colubridae, Dipsadidae, ecology, literature review, morphology, museum specimens, natural history, Reptilia, Serpentes, taxonomy

Citation: Barquero MD, Arguedas V. 2019. Biology of snakes of the genus *Tretanorhinus*: an integrative review. *Amphibian & Reptile Conservation* 13(2) [General Section]: 227–238 (e208).

Copyright: © 2019 Barquero and Arguedas. This is an open access article distributed under the terms of the Creative Commons Attribution License [Attribution 4.0 International (CC BY 4.0): <https://creativecommons.org/licenses/by/4.0/>], which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website: amphibian-reptile-conservation.org.

Received: 24 June 2018; **Accepted:** 6 December 2018; **Published:** 10 December 2019

Introduction

Fieldwork with snakes usually poses several challenges and more attention has focused on snake species from temperate areas than those from the tropics (Avila et al. 2006), despite the latter having higher diversity (Greene 1997). This bias has resulted in a lack of key information on essential aspects of the ecology, natural history, and behavior of many species. The collection of such data is a time-consuming and difficult task, as many snake species have cryptic habits and occur in low densities (Greene 1997). However, the lack of basic information for many snake species is a major concern, because this information is crucial for making general interspecific comparisons, establishing proper phylogenetic relationships, determining population trends and dynamics, and recognizing differences in behavior. Therefore, reviewing the current state of knowledge for a given species is imperative to identify the areas that require more attention and to guide research and conservation efforts in the proper direction.

Snakes of the genus *Tretanorhinus* are nocturnal, aquatic species that have intrigued biologists for more than a century (Cope 1861; Duméril et al. 1854). Their

habits and secretive behavior make them difficult to study, so their biology remains largely unknown (Savage 2002; Schwartz and Henderson 1991). Currently, four species are recognized in the genus: *T. mocquardi* (Bocourt 1891), *T. nigroluteus* (Cope 1861), *T. taeniatus* (Boulenger 1903), and *T. variabilis* (Duméril et al. 1854). Although a handful of studies have increased our knowledge about these species (e.g., Barquero et al. 2005; Dunn 1939; Henderson and Hoevers 1979; Villa 1970), the genus remains poorly studied.

Here we summarize and integrate all available published information, highlight the gaps in our knowledge of the biology of *Tretanorhinus* species, and make suggestions to direct future research in order to fill in these gaps. To achieve this, searches were conducted to find all published material referring directly (i.e., studies focused specifically on one or more *Tretanorhinus* species) or indirectly (i.e., studies focused on many taxa that mentioned one or more *Tretanorhinus* species as part of the topic) to snakes of this genus and a database of key information was compiled (Appendix 1). The information mentioned in each study was used to determine missing data for each species. Each publication was classified using the following categories:

Correspondence. ^{1,2,*}marco.barquero_a@ucr.ac.cr, ^{1,3}viviarguedas@gmail.com




Fig. 1. Number of specimens of the four species of *Tretanorhinus* according to (A) decade and (B) month of collection.

natural history, morphology, taxonomy, systematics, ecology, distribution, biogeography, and reproduction. In addition, data were obtained for specimens held in scientific collections (Appendix 2) from the HerpNet2 data portal (<http://www.herpNet.org>), Global Biodiversity Information Facility (<http://www.gbif.org>), Data Research Warehouse Information Network (<http://darwin.naturalsciences.be>), and by directly contacting collection curators. When available, the following data were extracted: taxonomic classification, type status, sex and age classes, country and locality of the point of capture, latitude and longitude of the collection site, date of collection, and remarks about the individual collected.

Literature Review and Scientific Collections

The search for publications referring to *Tretanorhinus* produced a total of 87 documents, although only 16 focused directly on species of the genus (Appendix 1). Most of the documents focused on general aspects of natural history (31%), taxonomy (21.8%), and biogeography (20.7%), and the vast majority referred to *T. nigroluteus* or *T. variabilis*, with only a handful of studies mentioning *T. mocquardi* ($n = 12$) or *T. taeniatus* ($n = 8$).

A database with information for a total of 755 specimens of *Tretanorhinus* held in 31 scientific collections was compiled (Appendix 2). Most records corresponded to *T. nigroluteus* ($n = 350$) or *T. variabilis* ($n = 357$), and only a few were available for *T. mocquardi* ($n = 25$) or *T. taeniatus* ($n = 9$). In addition, eight records

were identified only to the genus level, and six were mistakenly identified as *Tretanorhinus agassizi*. Most of the collections occurred between the 1940s and 1970s (Fig. 1A) and seasonally between June and August (Fig. 1B), although many records did not include either the year or month of collection.

Distribution and Habitat

Tretanorhinus is a genus exclusive to Middle and South America, and the West Indies. *Tretanorhinus variabilis* is the only one of the species with an unambiguous distribution, inhabiting Cuba, Isla de la Juventud (Isle of Youth), and the Cayman Islands (Fig. 2). *Tretanorhinus nigroluteus* has been frequently recorded along the Atlantic coast from southern Mexico to Colombia, including some islands such as the Bay Islands of Honduras and Great Corn Island of Nicaragua (Fig. 2). However, records of this species in Colombia are controversial. Alarcón-Pardo (1978) reported finding *T. nigroluteus* on the Atlantic coast of Colombia, but we were unable to locate any specimens of this species in the scientific collections consulted which matched the locality mentioned by that author. We found only one specimen of *T. nigroluteus* from the Pacific coast of Colombia as a disjunct point of the species range (Table 1). The distributions of *T. mocquardi* and *T. taeniatus* are the least clear among the species of the genus. *Tretanorhinus mocquardi* ranges from the Canal Zone in Panama through the Pacific coast of Colombia and Ecuador, although it has been reported from only two locations in Colombia and one in Ecuador (Fig. 2, Table 1). *Tretanorhinus taeniatus* could be endemic to Ecuador (Table 1), despite previous reports of this species in Colombia (Castaño-M. et al. 2004; Daniel 1949). Based on these distributions, *T. mocquardi* and *T. taeniatus* could be sympatric in Esmeraldas province, northwestern Ecuador, whereas *T. mocquardi* and *T. nigroluteus* seem to be sympatric in Chocó department on the Pacific coast of Colombia (Fig. 2).

Most information on the preferred habitat of the genus comes from studies conducted on *T. nigroluteus* and *T. variabilis*. *Tretanorhinus* is a fully aquatic genus, inhabiting all kinds of fresh and brackish water bodies such as rivers, streams, lagoons, estuaries, mangroves (Cisneros-Heredia 2005; Neill 1958; Villa 1970), and even cow wells (Seidel and Franz 1994). The species require a muddy or rocky bottom with aquatic vegetation where they can hide and rest (Neill 1965; Villa 1970). Although there are reports of individuals found out of the water (e.g., crossing roads), these events occur after flooding that forces the snakes to search for another water body (Barquero et al. 2005; Villa 1970). The distributions of *T. nigroluteus* and *T. variabilis* also confirm the ability of these snakes to disperse and survive in salt water (Barbour and Amaral 1924; Neill 1958), since they have colonized several islands. In that regard, we observed




Fig. 2. Map showing the current distribution of the four species of *Tretanorhinus* based on specimens from scientific collections.

Table 1. Number of specimens of *Tretanorhinus* species collected from different countries and held in several scientific collections. Specimens with a doubtful or missing location were excluded (n = 38).

Country	<i>T. mocquardi</i>	<i>T. nigroluteus</i>	<i>T. taeniatus</i>	<i>T. variabilis</i>	Total
West Indies					
Cayman Islands	-	-	-	58	58
Cuba	-	-	-	280	280
Middle America					
Belize	-	52	-	-	52
Costa Rica	-	11	-	-	11
Guatemala	-	11	-	-	11
Honduras	-	157	-	-	157
Mexico	-	30	-	-	30
Nicaragua	-	65	-	-	65
Panama	14	19	-	-	33
South America					
Colombia	4	1	-	-	5
Ecuador	6	-	9	-	15
Total	24	346	9	338	717

Biology of *Tretanorhinus* species

Table 2. Morphological information available in the literature for each species of *Tretanorhinus*.

Trait	<i>T. mocquardi</i>	<i>T. nigroluteus</i>	<i>T. taeniatus</i>	<i>T. variabilis</i>
SVL (mm)				
Adults	-	242–656	440	500–800
Juveniles	-	140–168	-	143–145
Tail length (mm)				
Adults	-	142–206	130	145–160
Juveniles	-	51–77	-	-
No. of loreals	1	1–2	1	1–3
No. of prefrontals	1–2	2	3	2
No. of preoculars	2	2–3	2	1–3
No. of postoculars	-	2	2	2
No. of temporals	-	1+2	1+2, 2+3	1+2
No. of upper labials	-	7–9	8	8–9
No. of lower labials	-	9–12	4–5	9–11
No. of ventrals	166–177	127–151	168–175	152–168
No. of caudals	69–85	56–81	74–81	48–81
Posterior chin-shields	In contact	In contact	Separated	Separated
No. of dorsal rows	19	19, 21	21	19, 21
Ventral color	Fuliginous yellow	Orange, light red, yellow, cream	-	Bluish gray, dark brown
Ventral pattern	-	With or without dots or spots	3 dark stripes	Light dots or spots
Dorsal color	-	Olive, grayish brown, light brown, black	Grayish olive	Grayish olive, dark brown
Dorsal pattern	3 longitudinal stripes	With or without dark spots	3 longitudinal stripes	Blackish cross bands

an individual of *T. nigroluteus* resting on a beach on the Caribbean coast of Costa Rica after a heavy rain, suggesting that the snake had been washed out to sea, surviving until it was returned to land by the tide.

Morphology

Morphological information is limited and incomplete for *T. mocquardi* and *T. taeniatus*, despite both species being described more than 110 yrs ago by Bocourt (1891) and Boulenger (1903), respectively. Table 2 provides a summary of some morphological features that were extracted from the literature review. Overall, *Tretanorhinus* are relatively small snakes; *T. variabilis* is the largest species with an SVL of up to 800 mm. All four species in the genus exhibit a grayish dorsal coloration with stripes, spots, or bands, and a yellow, orange, or gray ventral coloration. Juveniles have been found in the wild only infrequently, and the sex of most collected individuals was not identified (Table 3). Juveniles of each species resemble the adults in pattern and coloration (Barquero et al. 2005; Petzold 1967).

Previous attempts to produce a key to the species of the genus have never included all four species. Therefore, the following key is presented to unify the previous efforts (Bocourt 1891; Boulenger 1893; Dunn 1939; Peters and Orejas-Miranda 1970; Köhler 2008) and incorporate more recent data.

Key to the species of *Tretanorhinus*:

- 1a. Dorsum without stripes, more than one loreal can be present.....2
- 1b. Dorsum with three longitudinal stripes, only one loreal3
- 2a. Posterior chin-shields in contact, ventrals fewer than 152.....*nigroluteus*
- 2b. Posterior chin-shields separated, ventrals 152 or more.....*variabilis*
- 3a. Less than three prefrontals, posterior chin-shields in contact.....*mocquardi*
- 3b. Three prefrontals, posterior chin-shields separated*taeniatus*

Taxonomy and Systematics

The evolutionary relationships of the genus are enigmatic and controversial, so that *Tretanorhinus* has been placed within different former subfamilies of Colubridae (e.g., Natricinae, Xenodontinae, and Dipsadinae) by various authors (Crother 1999; Dowling et al. 1996; Pinou and Dowling 1994; Villa 1969). The general consensus is that the genus should be placed within Dipsadidae (formerly Dipsadinae, Grazziotin et al. 2012; Pinou and Dowling 1994) and it is commonly referred to as a xenodontine (Cadle 1985; Minton 1976; Vidal et al. 2000). Moreover, the phylogenetic relationships with other genera are still ambiguous. Crother (1999) placed *Tretanorhinus* as the sister taxon of *Sibon* or the *Sibon-Manolepis* clade. In two studies that included more taxa, Grazziotin et al. (2012) noted that *Tretanorhinus* could be the sister taxon of *Hypsiglena*, and both genera were placed in a clade containing *Trimetopon*, *Geophis*, and *Atractus*; whereas Pyron et al. (2013) placed *Tretanorhinus* as the sister taxon of the *Leptodeira-Imantodes* clade. However, in all these studies the relationships of *Tretanorhinus* with other taxa were poorly supported and the genus is considered as Dipsadidae *incertae sedis* (Grazziotin et al. 2012).

The affinities of each species within *Tretanorhinus* are also enigmatic, and this is caused by the lack of information on the basic biological aspects of some species. Detailed morphological information is only available for *T. nigroluteus* and *T. variabilis* (Table 2). No author has given a complete morphological description of *T. mocquardi*, such that even basic phenotypic traits (e.g., most head scutellation and dorsal coloration) are missing (Table 2). In the case of *T. taeniatus*, morphological information has only been reported for females, as males have never been deposited or identified as such in scientific collections (Table 3). Despite these deficiencies, morphological information compiled from the literature review here suggests that *T. taeniatus* and *T. mocquardi* are more closely related to each other than to *T. nigroluteus* and *T. variabilis*.

Problems have also arisen with the subspecies described for both *T. nigroluteus* (*dichromaticus*, *lateralis*, *mertensi*, *nigroluteus*, and *obscurus*) and *T. variabilis* (*binghami*, *insulaepinorum*, *lewisi*, *variabilis*, and *wagleri*). The morphological variables used to define the subspecies have been chosen arbitrarily (Wilson and Hahn 1973). For example, differences in coloration have been used by some authors to differentiate among subspecies with only superficial descriptions (Schwartz and Ogren 1956; Smith 1965; Villa 1969). In addition, no genetic analyses have yet been reported to confirm the validity of these subspecies.

Ecology and Natural History

Information on the ecology and natural history of *T. mocquardi* and *T. taeniatus* is virtually non-existent

Table 3. Number of specimens of *Tretanorhinus* held in scientific collections that have been identified as female, male, or juvenile.

Species	Female	Male	Juvenile	Total
<i>T. mocquardi</i>	2	1	-	3
<i>T. nigroluteus</i>	14	16	2	32
<i>T. taeniatus</i>	2	-	-	2
<i>T. variabilis</i>	19	17	4	40
Total	37	34	6	77

and they remain poorly studied for *T. nigroluteus* and *T. variabilis*. Snakes of *Tretanorhinus* are nocturnal and seem to hide during the day in water bodies amongst roots and rock crevices (Barbour and Ramsden 1919; Stuart 1937; Villa 1970). Some unusual features have been identified for the genus. For example, individuals of *Tretanorhinus* feed upon fishes, tadpoles, and frogs by either actively chasing prey or remaining motionless with the tail and body attached to a supporting surface (e.g., branch or rock) and striking at passing prey (Barquero et al. 2005; Neill 1965; Petzold 1967; Wilson and Hahn 1973). In addition, these snakes demonstrate shy behavior, such as fleeing to the bottom of water bodies when disturbed (Stuart 1937) and rolling up the body like a ball when caught (Petzold 1967; Seidel and Franz 1994). Known natural predators of *Tretanorhinus* include, but are probably not limited to, turtles (e.g., *Kinosternon* [Villa 1973]) and wading birds (e.g., *Tigrissoma* and *Cochlearius* [Villa 1970]). A specimen from Costa Rica was collected from a crab, thus confirming the assumption of Villa (1970) that some species of crabs can be predators of *Tretanorhinus*.

Information on reproduction is scarce for all four species of *Tretanorhinus*, although observations in captivity demonstrate that *T. nigroluteus* and *T. variabilis* are oviparous, laying 6–9 adherent eggs out of water (Barquero et al. 2005; Petzold 1967). *Tretanorhinus variabilis* lays larger eggs (35[L] x 16.75[W] mm on average) that hatch earlier (35 d) than *T. nigroluteus* (21.5[L] x 10[W] mm, 42 d). Villa (1970) found gravid females of *T. nigroluteus* during both the dry and wet seasons, suggesting that reproduction could occur year-round in this species. Most gravid females of *T. variabilis* have been found during the wet season in July and August (Petzold 1967; Seidel and Franz 1994). Both *T. nigroluteus* and *T. variabilis* are sexually dimorphic, with females being larger than males and only males possessing tubercles on scales of the head (Henderson and Hoevers 1979; Petzold 1967).

The capacity to survive in salt water likely contributed to the colonization of Caribbean islands and northeastern South America from a Central American ancestor (Cisneros-Heredia 2005; Hedges 1996) and allowed *T. nigroluteus* and *T. variabilis* to become fairly abundant in some parts of their ranges (Henderson and Hoevers 1977; Schwartz and Henderson 1991). Henderson and Hoevers (1977) reported that *T. nigroluteus* was more frequently

found during the dry season (December to May) than during the wet season (June to November) of Belize, a difference explained by the overflowing of the river system studied. However, historical collections of this species show that more individuals have been collected during the wet season, a pattern shared with *T. variabilis* (Fig. 1), and in accordance with reports by Wilson and Hahn (1973) for Roatan Island, Honduras. *Tretanorhinus mocquardi* and *T. taeniatus* are less abundant than the other two species and no pattern of variation in abundance can be elucidated from available data.

General Considerations and Future Research

This study has summarized information about the species of the genus *Tretanorhinus* published from 1854 (Duméril et al. 1854) to the present (Estrella-Morales and Piedra-Castro 2018). Information was also incorporated on all collected specimens of this genus that could be identified as preserved in scientific collections throughout the world. This integrative approach allowed the identification of gaps in our knowledge about these snakes. For example, it is surprising that (1) a complete morphological description is not available in the literature for *T. mocquardi*, (2) only a few specimens have been collected for *T. mocquardi* and *T. taeniatus*, and (3) most natural history and ecological information simply has never been reported for any of the species. In addition to the lack of key data, much available information is contradictory, such as the reported occurrence of *T. nigroluteus* and *T. taeniatus* in Colombia, which can cause several problems. Therefore, one can ask how a reliable identification of individuals in the field can be made when such basic data are missing. This is particularly problematic for sympatric species, such as *T. nigroluteus* and *T. mocquardi* in Panama, and for *T. mocquardi* and *T. taeniatus* in Ecuador.

In order to fill in the gaps in our knowledge of *Tretanorhinus*, future research should focus on at least three areas. First, validation of the currently accepted species and subspecies is absolutely urgent. Barbour and Amaral (1924) have questioned the validity of *T. mocquardi* (although see Dunn 1939), while Wilson and Hahn (1973) refused to recognize *T. n. dichromaticus*. Therefore, a comparative study of all species and subspecies that includes both morphological and genetic data and produces a phylogeny of the genus is necessary. Previous attempts have failed to include all species or have used only morphological or genetic data. Second, both field and lab studies are needed to increase our knowledge about these secretive and, in some areas, elusive species. *Tretanorhinus mocquardi* and *T. taeniatus* require extensive work on morphological variation, ecological habits, distribution patterns, and natural history traits. Breeding behavior is completely unknown for these two species and males of *T. taeniatus* have yet to be measured and described. Although

there is significantly more information available for *T. nigroluteus* and *T. variabilis*, nothing is known about their courtship, sexual selection, development, and many other aspects of basic biology. Third, demographic variation and population dynamics need to be quantified to understand the movement of individuals among populations, sex ratios, and population sizes. These types of data are essential for determining the conservation status of species. Some efforts have already been made to identify areas of high and low abundance across the ranges of *T. nigroluteus* and *T. variabilis*. However, long-term studies which monitor changes in populations are yet to be done.

The problems mentioned above are not restricted only to *Tretanorhinus* species, as they also apply to many other snakes (Dorcas and Willson 2009), and it is alarming that we rely so heavily on unconfirmed or erroneous information. Snakes in particular require special attention due to the intrinsic difficulties in generating accurate information. These difficulties arise from certain aspects of their biology, including low densities, great mobility, and cryptic habits. Integrative studies, such as this one, are important for identifying the gaps in our knowledge of different taxa and guiding future efforts in the right direction.

Acknowledgements.—We are grateful to all the curators at institutions that kindly provided us with the information on specimens of *Tretanorhinus*. We are also thankful to Pablo Allen, Guido Saborío, and Rowan McGinley for their helpful comments on the manuscript.

Literature Cited

- Alarcón-Pardo H. 1978. Primer registro de *Tretanorhinus nigroluteus nigroluteus* Cope (Reptilia: Serpentes: Colubridae) para Colombia. *Lozania* 27: 1–4.
- Avila RW, Ferreira VL, Arruda JAO. 2006. Natural history of the South American water snake *Helicops leopardinus* (Colubridae: Hydropsini) in the Pantanal, Central Brazil. *Journal of Herpetology* 40: 274–279.
- Barbour T, Amaral AD. 1924. Notes on some Central American snakes. *Occasional Papers of the Boston Society of Natural History* 5: 129–132.
- Barbour T, Ramsden CT. 1919. The herpetology of Cuba. *Memoirs of the Museum of Comparative Zoölogy at Harvard College* 47: 69–213.
- Barquero MD, Campos-Chinchilla P, Valverde R. 2005. *Tretanorhinus nigroluteus*: biology and captive maintenance of the Orangebelly Swamp Snake. *Reptilia* 42: 47–52.
- Bocourt MF. 1891. Sur quelques ophidiens de l’Amérique intertropicale: appartenant au genre *Tretanorhinus*. *Le Naturaliste* 101: 121–122.
- Boulenger GA. 1893. *Catalogue of the Snakes in the British Museum (Natural History)*. Volume I. Taylor and Francis, London, United Kingdom. 504 p.

- Boulenger GA. 1903. Descriptions of new snakes in the collection of the British Museum. *Annals and Magazine of Natural History; Zoology, Botany, and Geology Series 7*, 12: 350–354.
- Cadle JE. 1985. The Neotropical colubrid snake fauna (Serpentes: Colubridae): lineage components and biogeography. *Systematic Zoology* 34: 1–20.
- Castaño-M OV, Cárdenas-A G, Hernández-R EJ, Castro-H F. 2004. Reptiles en el Chocó biogeográfico. Pp. 599–632 In: *Diversidad Biótica IV. El Chocó Biogeográfico/Costa Pacífica*. Editor, Rangel-Ch JO. Universidad Nacional de Colombia, Instituto de Ciencias Naturales, Conservation International, Bogotá, Colombia. 997 p.
- Cisneros-Heredia DF. 2005. *Tretanorhinus mocquardi* (Swamp Snake). *Herpetological Review* 36: 340.
- Cope ED. 1861. Contributions to the ophiology of Lower California, Mexico, and Central America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 13: 292–306.
- Crother BI. 1999. Phylogenetic relationships among West Indian xenodontine snakes (Serpentes; Colubridae) with comments on the phylogeny of some mainland xenodontines. *Contemporary Herpetology* 2: 7–30.
- Daniel H. 1949. Las serpientes de Colombia. *Revista Facultad Nacional de Agronomía Medellín* 10: 301–333.
- Dorcas ME, Willson JD. 2009. Innovative methods for studies of snake ecology and conservation. Pp. 5–37 In: *Snakes: Ecology and Conservation*. Editors, Mullin SJ, Seigel RA. Cornell University Press, Ithaca, New York, USA. 365 p.
- Dowling HG, Hass CA, Hedges SB, Highton R. 1996. Snake relationships revealed by slow-evolving proteins: a preliminary survey. *Journal of Zoology* 240: 1–28.
- Duméril AMC, Bibron G, Duméril A. 1854. *Erpétologie Générale ou Histoire Naturelle Complète des Reptiles*. Volume 7 (partie 1). Librairie Encyclopédique de Roret, Paris, France. 780 p.
- Dunn ER. 1939. Mainland forms of the snake genus *Tretanorhinus*. *Copeia* 1939: 212–217.
- Estrella-Morales J, Piedra-Castro L. 2018. Anfíbios y reptiles (herpetofauna) en las asociaciones vegetales de la Laguna de Gandoca, Limón, Costa Rica. *Tecnología en Marcha* 31: 127–135.
- Greene HW. 1997. *Snakes: the Evolution of Mystery in Nature*. University of California Press, Berkeley, California, USA. 365 p.
- Grazziotin FG, Zaher H, Murphy RW, Scrocchi G, Benavides MA, Zhang YP, Bonatto SL. 2012. Molecular phylogeny of the New World Dipsadidae (Serpentes: Colubroidea): a reappraisal. *Cladistics* 28: 437–459.
- Hedges SB. 1996. The origin of West Indian amphibians and reptiles. Pp. 95–128 In: *Contributions to West Indian Herpetology: a Tribute to Albert Schwartz*. Editors, Powell R, Henderson RW. Society for the Study of Amphibians and Reptiles, Ithaca, New York, USA. 457 p.
- Henderson RW, Hoevers LG. 1977. The seasonal incidence of snakes at a locality in northern Belize. *Copeia* 1977: 349–355.
- Henderson RW, Hoevers LG. 1979. Variation in the snake *Tretanorhinus nigroluteus lateralis* in Belize with notes on breeding tubercles. *Herpetologica* 35: 245–248.
- Köhler G. 2008. *Reptiles of Central America*. 2nd Edition. Herpeton-Verlag, Offenbach, Germany. 400 p.
- Minton SA Jr. 1976. Serological relationships among some congeneric North American and Eurasian colubrid snakes. *Copeia* 1976: 672–678.
- Neill WT. 1958. The occurrence of amphibians and reptiles in saltwater areas, and a bibliography. *Bulletin of Marine Science of the Gulf and Caribbean* 8: 1–97.
- Neill WT. 1965. Notes on aquatic snakes, *Natrix* and *Tretanorhinus*, in Cuba. *Herpetologica* 21: 172–181.
- Peters JA, Orejas-Miranda B. 1970. Catalogue of the Neotropical Squamata. Part I. Snakes. *Bulletin United States National Museum* 297: 1–347.
- Petzold HG. 1967. Some remarks on the breeding biology and the keeping of *Tretanorhinus variabilis*, a water snake of Cuba. *Herpetologica* 23: 242–246.
- Pinou T, Dowling HG. 1994. The phylogenetic relationships of the Central American snake *Tretanorhinus*: data from morphology and karyology. *Amphibia-Reptilia* 15: 297–305.
- Pyron RA, Burbrink FT, Wiens JJ. 2013. A phylogeny and revised classification of Squamata, including 4,161 species of lizards and snakes. *BMC Evolutionary Biology* 13: 93.
- Savage JM. 2002. *The Amphibians and Reptiles of Costa Rica: a Herpetofauna between Two Continents, between Two Seas*. University of Chicago Press, Chicago, Illinois, USA. 954 p.
- Schwartz A, Henderson RW. 1991. *Amphibians and Reptiles of the West Indies: Descriptions, Distributions, and Natural History*. University of Florida Press, Gainesville, Florida, USA. 720 p.
- Schwartz A, Ogren LH. 1956. A collection of reptiles and amphibians from Cuba, with the descriptions of two new forms. *Herpetologica* 12: 91–110.
- Seidel ME, Franz R. 1994. Amphibians and reptiles (exclusive of marine turtles) of the Cayman Islands. Pp. 407–433 In: *The Cayman Islands: Natural History and Biogeography*. Editors, Brunt MA, Davies JE. Kluwer Academic Publishers, Dordrecht, Netherlands. 604 p.
- Smith HM. 1965. Two new colubrid snakes from the United States and Mexico. *Journal of the Ohio Herpetological Society* 5: 1–4.
- Stuart LC. 1937. Some further notes on the amphibians and reptiles of the Peten forest of northern Guatemala. *Copeia* 1937: 67–70.

Biology of *Tretanorhinus* species

- Vidal N, Kindl SG, Wong A, Hedges SB. 2000. Phylogenetic relationships of xenodontine snakes inferred from 12S and 16S ribosomal RNA sequences. *Molecular Phylogenetics and Evolution* 14: 389–402.
- Villa J. 1969. Two new insular subspecies of the natricid snake *Tretanorhinus nigroluteus* Cope from Honduras and Nicaragua. *Journal of Herpetology* 3: 145–150.
- Villa J. 1970. Notas sobre la historia natural de la serpiente de los pantanos, *Tretanorhinus nigroluteus*. *Revista de Biología Tropical* 17: 97–104.
- Villa J. 1973. A snake in the diet of a kinosternid turtle. *Journal of Herpetology* 7: 380–381.
- Wilson LD, Hahn DE. 1973. The herpetofauna of the Islas de la Bahia, Honduras. *Bulletin of the Florida State Museum, Biological Sciences* 17: 93–150.



Marco D. Barquero is a Costa Rican biologist with a Doctorate degree from Macquarie University, Sydney, Australia. Marco has dedicated the last 18 years to the study of amphibians and reptiles in different parts of the world, but especially in Costa Rica.



Viviana Arguedas is a Costa Rican biologist with a Master's degree from the University of Costa Rica. Viviana has studied different taxa over the last 14 years, with an emphasis on herpetofauna.

Appendix 1. List of studies referring to *Tretanorhinus* species.

1. Alarcón-Pardo H. 1978. Primer registro de *Tretanorhinus nigroluteus nigroluteus* Cope (Reptilia: Serpentes: Colubridae) para Colombia. *Lozania* 27: 1–4.
2. Auth DL. 1994. Checklist and bibliography of the amphibians and reptiles of Panama. *Smithsonian Herpetological Information Service* 98: 1–59.
3. Barbour T. 1914. A contribution to the zoogeography of the West Indies, with special reference to amphibians and reptiles. *Memoirs of the Museum of Comparative Zoölogy at Harvard College* 44: 207–359.
4. Barbour T. 1916. The reptiles and amphibians of the Isle of Pines. *Annals of Carnegie Museum* 10: 207–308.
5. Barbour T; Amaral AD. 1924. Notes on some Central American snakes. *Occasional Papers of the Boston Society of Natural History* 5: 129–132.
6. Barbour T; Ramsden CT. 1916. Catalogo de los reptiles y anfibios de la Isla de Cuba. *Memorias de la Sociedad Cubana de Historia Natural "Felipe Poey" Habana* 2: 124–143.
7. Barbour T; Ramsden CT. 1919. The herpetology of Cuba. *Memoirs of the Museum of Comparative Zoölogy at Harvard College* 47: 69–213.
8. Barquero MD; Campos-Chinchilla P; Valverde R. 2005. *Tretanorhinus nigroluteus*: biology and captive maintenance of the Orangebelly Swamp Snake. *Reptilia* 42: 47–52.
9. Bocourt MF. 1891. Sur quelques ophidiens de l'Amerique intertropicale: appartenant au genre *Tretanorhinus*. *Le Naturaliste* 101: 121–122.
10. Boulenger GA. 1893. *Catalogue of the Snakes in the British Museum (Natural History)*. Volume I. Taylor and Francis, London, United Kingdom. 504 p.
11. Boulenger GA. 1903. Descriptions of new snakes in the collection of the British Museum. *Annals and Magazine of*

Natural History: Zoology, Botany, and Geology Series 7, 12: 350–354.

12. Cadle JE. 1984. Molecular systematics of Neotropical xenodontine snakes: II. Central American xenodontines. *Herpetologica* 40: 21–30.
13. Cadle JE. 1985. The Neotropical colubrid snake fauna (Serpentes: Colubridae): Lineage components and biogeography. *Systematic Zoology* 34: 1–20.
14. Campbell JA. 1999. *Amphibians and Reptiles of Northern Guatemala, the Yucatan, and Belize*. University of Oklahoma Press, Norman, Oklahoma, USA. 400 p.
15. Castaño-M OV; Cárdenas-A G; Hernández-R EJ; Castro-H F. 2004. Reptiles en el Chocó biogeográfico. Pp. 599–632 In: *Diversidad Biótica IV. El Chocó Biogeográfico/Costa Pacífica*. Editor, Rangel-Ch JO. Universidad Nacional de Colombia, Instituto de Ciencias Naturales, Conservation International, Bogotá, Colombia. 997 p.
16. Cisneros-Heredia DF. 2005. *Tretanorhinus mocquardi* (Swamp Snake). *Herpetological Review* 36: 340.
17. Conant R. 1965. Miscellaneous notes and comments on toads, lizards, and snakes from Mexico. *American Museum Novitates* 2205: 1–38.
18. Cope ED. 1861. Contributions to the ophiology of Lower California, Mexico, and Central America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 13: 292–306.
19. Crother BI. 1989. A redescription of the hemipenis of *Hydromorphus concolor* (Colubridae) with comments on its tribal allocation. *Copeia* 1989: 227–229.
20. Crother BI. 1999. Phylogenetic relationships among West Indian xenodontine snakes (Serpentes; Colubridae) with comments on the phylogeny of some mainland xenodontines. *Contemporary Herpetology* 2: 7–30.
21. Daniel H. 1949. Las serpientes de Colombia. *Revista Facultad Nacional de Agronomía Medellín* 10: 301–333.
22. Dowling HG; Hass CA; Hedges SB; Highton R. 1996. Snake relationships revealed by slow-evolving proteins: a preliminary survey. *Journal of Zoology* 240: 1–28.
23. Duellman WE. 1963. Amphibians and reptiles of the rainforests of southern El Petén, Guatemala. *University of Kansas Publications, Museum of Natural History* 15: 205–249.
24. Duméril AMC; Bibron G; Duméril A. 1854. *Erpétologie Générale ou Histoire Naturelle Complète des Reptiles*. Volume 7 (partie 1). Librairie Encyclopédique de Roret, Paris, France. 780 p.
25. Dunn ER. 1932. The colubrid snakes of the Greater Antilles. *Copeia* 1932: 89–92.
26. Dunn ER. 1939. Mainland forms of the snake genus *Tretanorhinus*. *Copeia* 1939: 212–217.
27. Estrella-Morales J; Piedra-Castro L. 2018. Anfíbios y reptiles (herpetofauna) en las asociaciones vegetales de la Laguna de Gandoca, Limón, Costa Rica. *Tecnología en Marcha* 31: 127–135.
28. García-Balderas CM; Carbajal-Márquez RA; Cedeño-Vázquez JR. 2016. Distribution notes. *Tretanorhinus nigroluteus* Cope, 1861. *Mesoamerican Herpetology* 3: 200.
29. Garman S. 1883. The reptiles and batrachians of North America. *Memoirs of the Museum of Comparative Zoology at Harvard College* 3: 1–219.
30. González-C A; Rodríguez-S L; Rivalta-G V. 2014. Colección herpetológica del Instituto de Ecología y Sistemática, La Habana, Cuba: Familia Dipsadidae, géneros *Caraiba* y *Tretanorhinus*; familia Natricidae, género *Nerodia*. *Revista Colombiana de Ciencia Animal* 6: 38–54.
31. González-Sánchez VH; Johnson JD; García-Padilla E; Mata-Silva V; DeSantis DL; Wilson LD. 2017. The herpetofauna of the Mexican Yucatan Peninsula: composition, distribution, and conservation status. *Mesoamerican Herpetology* 4: 263–380.
32. Grant C. 1941. The herpetology of the Cayman Islands. *Bulletin of the Institute of Jamaica, Science Series* 2: 1–56.
33. Grant C. 1949. Notes on *Tretanorhinus* in Cuba and the Isle of Pines. *Journal of Agriculture of the University of Puerto Rico* 30: 102–117.
34. Grazziotin FG; Zaher H; Murphy RW; Scrocchi G; Benavides MA; Zhang YP; Bonatto SL. 2012. Molecular phylogeny of the New World Dipsadidae (Serpentes: Colubroidea): a reappraisal. *Cladistics* 28: 437–459.
35. Gundlach J. 1880. *Contribucion a la Erpétologia Cubana*. G. Montiel y Ca, Habana, Cuba. 99 p.
36. Günther ACLG. 1885–1902. *Biologia Centrali-Americana. Reptilia and Batrachia*. F. Du Cane Godman and Osbert Savin, London, United Kingdom. 7 Volumes.
37. Hedges SB. 1996. The origin of West Indian amphibians and reptiles. Pp. 95–128 In: *Contributions to West Indian Herpetology: a Tribute to Albert Schwartz*. Editors, Powell R, Henderson RW. Society for the Study of Amphibians and Reptiles, Ithaca, New York, USA. 457 p.
38. Henderson RW; Hoevers LG. 1977. The seasonal incidence of snakes at a locality in northern Belize. *Copeia* 1977: 349–355.
39. Henderson RW; Hoevers LG. 1979. Variation in the snake *Tretanorhinus nigroluteus lateralis* in Belize with notes on breeding tubercles. *Herpetologica* 35: 245–248.
40. Hernández-Ordóñez O; Arroyo-Rodríguez V; González-Hernández A; Russildi G; Luna-Reyes R; Martínez-Ramos M; Reynoso VH. 2015. Range extensions of amphibians and reptiles in the southeastern part of the Lacandona rainforest, Mexico. *Revista Mexicana de Biodiversidad* 86: 457–468.
41. Highton R; Hedges SB; Hass CA; Dowling HG. 2002. Snake relationships revealed by slowly-evolving proteins: further analysis and a reply. *Herpetologica* 58: 270–275.
42. Jan G. 1865. Enumerazione sistematica degli ofidi appartenenti ai gruppo Potamophilidae. *Archive per la Zoologia, l'Anatomia et la Fisiologia* 3: 201–265.

43. Johnson JD; Mata-Silva V; García Padilla E; Wilson LD. 2015. The herpetofauna of Chiapas, Mexico: composition, distribution, and conservation. *Mesoamerican Herpetology* 2: 271–329.
44. Köhler G. 2008. *Reptiles of Central America*. 2nd edition. Herpeton-Verlag, Offenbach, Germany. 400 p.
45. Köhler G; Cedeño-Vázquez JR; Beutelspacher-García PM. 2016. The Chetumal Snake Census: generating biological data from road-killed snakes. Part 1. Introduction and identification key to the snakes of southern Quintana Roo, Mexico. *Mesoamerican Herpetology* 3: 669–687.
46. Lee JC. 2000. *A Field Guide to the Amphibians and Reptiles of the Maya World*. Cornell University Press, Ithaca, New York, USA. 416 p.
47. Malnate EV. 1972. Observations on the vertebral hypapophyses and associated musculature in some snakes, with special reference to the Colubridae. *Zoologische Mededelingen* 47: 225–239.
48. Marx H; Rabb GB. 1972. Phyletic analysis of fifty characters of advanced snakes. *Fieldiana Zoology* 63: 1–321.
49. Mata-Silva V; Johnson JD; Wilson LD; García-Padilla E. 2015. The herpetofauna of Oaxaca, Mexico: composition, physiographic distribution, and conservation status. *Mesoamerican Herpetology* 2: 5–62.
50. McCoy CJ. 1970. The snake fauna of Middlesex, British Honduras. *Journal of Herpetology* 4: 135–140.
51. McCranie JR. 2011. *The Snakes of Honduras. Systematics, Distribution, and Conservation*. Society for the Study of Amphibians and Reptiles, Ithaca, New York, USA. 714 p.
52. Mead JI; Steadman DW. 2017. Late Pleistocene snakes (Squamata: Serpentes) from Abaco, The Bahamas. *Geobios* 50: 431–440.
53. Meyer JR. 1966. Records and observations on some amphibians and reptiles from Honduras. *Herpetologica* 22: 172–181.
54. Minton SA Jr. 1976. Serological relationships among some congeneric North American and Eurasian colubrid snakes. *Copeia* 1976: 672–678.
55. Neill WT. 1958. The occurrence of amphibians and reptiles in saltwater areas, and a bibliography. *Bulletin of Marine Science of the Gulf and Caribbean* 8: 1–97.
56. Neill WT. 1965. Notes on aquatic snakes, *Natrix* and *Tretanorhinus*, in Cuba. *Herpetologica* 21: 172–181.
57. Peters JA. 1960. The snakes of Ecuador; check list and key. *Bulletin of the Museum of Comparative Zoölogy at Harvard College* 122: 489–541.
58. Peters JA; Orejas-Miranda B. 1970. Catalogue of the Neotropical Squamata. Part I. Snakes. *Bulletin United States National Museum* 297: 1–347.
59. Peters JL; Allen GM; Barbour T; Loveridge A. 1929. Vertebrates from the Corn Islands. *Bulletin of the Museum of Comparative Zoölogy at Harvard College* 69: 125–146.
60. Petzold HG. 1967. Some remarks on the breeding biology and the keeping of *Tretanorhinus variabilis*, a water snake of Cuba. *Herpetologica* 23: 242–246.
61. Pinou T; Dowling HG. 1994. The phylogenetic relationships of the Central American snake *Tretanorhinus*: data from morphology and karyology. *Amphibia-Reptilia* 15: 297–305.
62. Ray JM; Ruback P. 2015. Updated checklists of snakes for the provinces of Panamá and Panamá Oeste, Republic of Panama. *Mesoamerican Herpetology* 2: 167–188.
63. Rodríguez Schettino L; Mancina CA; Rivalta González V. 2013. Reptiles of Cuba: checklist and geographic distributions. *Smithsonian Herpetological Information Service* 144: 1–96.
64. Rosén N. 1905. List of the snakes in the zoological museums of Lund and Malmö, with descriptions of new species and a new genus. *Annals and Magazine of Natural History; Zoology, Botany, and Geology Series* 7, 15: 168–181.
65. Sánchez-C H; Castaño-M O; Cárdenas-A G. 1995. Diversidad de los reptiles en Colombia. Pp. 277–326 In: *Colombia Diversidad Biótica I*. Editor, Rangel-Ch JO. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Inderena, Bogotá, Colombia. 442 p.
66. Savage JM. 2002. *The Amphibians and Reptiles of Costa Rica: a Herpetofauna between Two Continents, between Two Seas*. University of Chicago Press, Chicago, Illinois, USA. 954 p.
67. Schmidt KP. 1935. Amphibians and reptiles collected by the Smithsonian Biological Survey of the Panama Canal Zone. *Smithsonian Miscellaneous Collections* 89: 1–20.
68. Schmidt KP. 1941. The amphibians and reptiles of British Honduras. *Field Museum of Natural History, Zoological Series* 22: 475–510.
69. Schwartz A; Henderson RW. 1991. *Amphibians and Reptiles of the West Indies: Descriptions, Distributions, and Natural History*. University of Florida Press, Gainesville, Florida, USA. 720 p.
70. Schwartz A; Ogren LH. 1956. A collection of reptiles and amphibians from Cuba, with the descriptions of two new forms. *Herpetologica* 12: 91–110.
71. Seidel ME; Franz R. 1994. Amphibians and reptiles (exclusive of marine turtles) of the Cayman Islands. Pp. 407–433 In: *The Cayman Islands: Natural History and Biogeography*. Editors, Brunt MA, Davies JE. Kluwer Academic Publishers, Dordrecht, Netherlands. 604 p.
72. Smith HM. 1965. Two new colubrid snakes from the United States and Mexico. *Journal of the Ohio Herpetological Society* 5: 1–4.
73. Smith HM; Taylor EH. 1945. An annotated checklist and key to the snakes of Mexico. *Bulletin United States National Museum* 187: 1–239.
74. Solís JM; Wilson LD; Townsend JH. 2014. An updated list of the amphibians and reptiles of Honduras, with

- comments on their nomenclature. *Mesoamerican Herpetology* 1: 122–144.
75. Solórzano A. 2004. *Serpientes de Costa Rica: Distribución, Taxonomía, e Historia Natural = Snakes of Costa Rica: Distribution, Taxonomy, and Natural History*. Instituto Nacional de Biodiversidad, Santo Domingo de Heredia, Costa Rica. 791 p.
 76. Stuart LC. 1937. Some further notes on the amphibians and reptiles of the Peten forest of northern Guatemala. *Copeia* 1937: 67–70.
 77. Sunyer J. 2014. An updated checklist of the amphibians and reptiles of Nicaragua. *Mesoamerican Herpetology* 1: 185–202.
 78. Taylor EH. 1951. A brief review of the snakes of Costa Rica. *The University of Kansas Science Bulletin* 34: 3–188.
 79. Torres-Carvajal O; Salazar-Valenzuela D; Merino-Viteri A. 2014. ReptiliaWebEcuador, Versión 2014.0. Museo de Zoología QCAZ, Pontificia Universidad Católica del Ecuador, Quito, Ecuador. Available: <http://zoologia.puce.edu.ec/Vertebrados/reptiles/reptilesEcuador> [Accessed: 4 March 2015].
 80. Vaeth RH; Rossman DA; Shoop W. 1985. Observations of tooth surface morphology in snakes. *Journal of Herpetology* 19: 20–26.
 81. Vidal N; Kindl SG; Wong A; Hedges SB. 2000. Phylogenetic relationships of xenodontine snakes inferred from 12S and 16S ribosomal RNA sequences. *Molecular Phylogenetics and Evolution* 14: 389–402.
 82. Villa J. 1969. Two new insular subspecies of the natricid snake *Tretanorhinus nigroluteus* Cope from Honduras and Nicaragua. *Journal of Herpetology* 3: 145–150.
 83. Villa J. 1970. Notas sobre la historia natural de la serpiente de los pantanos, *Tretanorhinus nigroluteus*. *Revista de Biología Tropical* 17: 97–104.
 84. Villa J. 1973. A snake in the diet of a kinosternid turtle. *Journal of Herpetology* 7: 380–381.
 85. Wilson LD; Hahn DE. 1973. The herpetofauna of the Islas de la Bahía, Honduras. *Bulletin of the Florida State Museum, Biological Sciences* 17: 93–150.
 86. Wood GC. 1939. The genus *Tretanorhinus* in Cuba and the Isle of Pines. *Proceedings of the New England Zoological Club* 18: 5–11.
 87. Zaher H; de Oliveira L; Grazziotin FG; Campagner M; Jared C; Antoniazzi MM; Prudente AL. 2014. Consuming viscous prey: a novel protein-secreting delivery system in neotropical snail-eating snakes. *BMC Evolutionary Biology* 14: 58–86.

Biology of *Tretanorhinus* species

Appendix 2. Number of specimens of *Tretanorhinus* held in each scientific collection from which data were obtained.

Specimens	Abbrev	Institution name
7	ANSP	Academy of Natural Sciences, Philadelphia
143	AMNH	American Museum of Natural History
1	BYU	Bingham Young University, Monte L. Bean Life Science Museum
59	CAS	California Academy of Sciences
12	CM	Carnegie Museum of Natural History
2	CHP	Círculo Herpetológico de Panamá
12	FMNH	Field Museum of Natural History
82	FLMNH	Florida Museum of Natural History
6	FHGO	Fundación Herpetológica Gustavo Orcés
12	INHS	Illinois Natural History Survey, University of Illinois (formerly University of Illinois Museum of Natural History [UIMNH])
5	IBUNAM	Instituto de Biología, Universidad Nacional Autónoma de México
8	IAvH	Instituto de Investigación de Recursos Biológicos Alexander von Humboldt
35	LACM	Los Angeles County Museum of Natural History
64	LSUMZ	Louisiana Museum of Natural History (formerly Louisiana State University, Museum of Zoology)
45	MPM	Milwaukee Public Museum
15	MNHN	Muséum National d'Histoire Naturelle
60	MCZ	Museum of Comparative Zoology, Harvard University
3	MVZ	Museum of Vertebrate Zoology, University of California at Berkeley
52	USNM	National Museum of Natural History, Smithsonian Institution (formerly United States National Museum)
11	BMNH	Natural History Museum (formerly British Museum of Natural History)
5	RBINS	Royal Belgian Institute of Natural Sciences
3	ROM	Royal Ontario Museum
3	TCWC	Texas Cooperative Wildlife Collection
2	MHUA	Universidad de Antioquia, Museo de Herpetología
12	MZUCR	Universidad de Costa Rica, Museo de Zoología
2	UVC	Universidad del Valle
1	UTCH	Universidad Tecnológica del Chocó
2	UCM	University of Colorado, Museum of Natural History
77	KU	University of Kansas, Biodiversity Institute
10	UMMZ	University of Michigan, Museum of Zoology
4	UTA	University of Texas at Arlington
755	Total	